


AOP 212: Histone deacetylase inhibition leading to testicular atrophy

Short Title: Histone deacetylase inhibition leading to testicular atrophy

Graphical Representation

AOP212:
Histone deacetylase inhibition leading to testicular atrophy

**Authors**

Shihori Tanabe, Akihiko Hirose, Takashi Yamada

Division of Risk Assessment, Biological Safety Research Center, National Institute of Health Sciences

Status

Author status	OECD status	OECD project	SAAOP status
Under development: Not open for comment. Do not cite	EAGMST Under Review	1.52	Included in OECD Work Plan

Abstract

Testicular toxicity is of interest for human health risk assessment especially in terms of reproductive and developmental toxicity, however, the testicular toxicity has not fully elucidated. Histone deacetylase inhibitors (HDIs) are approved as anti-cancer drugs, since HDIs have apoptotic effect in cancer cells. HDIs includes the short chain fatty acids, hydroxamic acids, benzamides and epoxides. The intracellular mechanisms of induction of the spermatocyte apoptosis by HDIs are suggested as histone deacetylase (HDAC) inhibition as MIE, histone acetylation increase, disrupted cell cycle, apoptosis, and spermatocyte depletion as KEs. Adverse outcome has been defined as testicular atrophy. The HDIs inhibit deacetylation of the histone, leading to the increase in histone acetylation. The apoptosis induced by disrupted cell cycle leads to spermatocyte depletion and testis atrophy. This AOP may be one of the pathways induced by HDIs, which suggests the pathway networks of protein hyperacetylations.

Abbreviation: AOP: adverse outcome pathway, HDAC: histone deacetylase, HDI: HDAC inhibitor, KE: key event, MIE: molecular initiating event

Summary of the AOP

Events

Molecular Initiating Events (MIE), Key Events (KE), Adverse Outcomes (AO)

Sequence	Type	Event ID	Title	Short name
1	MIE	1502	Histone deacetylase inhibition (https://aopwiki.org/events/1502)	Histone deacetylase inhibition
2	KE	1503	Histone acetylation, increase (https://aopwiki.org/events/1503)	Histone acetylation, increase
3	KE	1505	Cell cycle, disrupted (https://aopwiki.org/events/1505)	Cell cycle, disrupted
4	KE	1262	Apoptosis (https://aopwiki.org/events/1262)	Apoptosis
5	KE	1515	Spermatocyte depletion (https://aopwiki.org/events/1515)	Spermatocyte depletion
6	AO	1506	Testicular atrophy (https://aopwiki.org/events/1506)	Testicular atrophy

Key Event Relationships

Upstream Event	Relationship Type	Downstream Event	Evidence	Quantitative Understanding
Histone deacetylase inhibition (https://aopwiki.org/relationships/1709)	adjacent	Histone acetylation, increase	High	Moderate
Histone acetylation, increase (https://aopwiki.org/relationships/1997)	adjacent	Cell cycle, disrupted	Moderate	Moderate
Cell cycle, disrupted (https://aopwiki.org/relationships/1712)	adjacent	Apoptosis	Moderate	Moderate
Apoptosis (https://aopwiki.org/relationships/1735)	adjacent	Spermatocyte depletion	High	Not Specified
Spermatocyte depletion (https://aopwiki.org/relationships/1734)	adjacent	Testicular atrophy	High	Not Specified
Histone deacetylase inhibition (https://aopwiki.org/relationships/1715)	non-adjacent	Cell cycle, disrupted	High	Moderate
Histone deacetylase inhibition (https://aopwiki.org/relationships/1716)	non-adjacent	Apoptosis	Moderate	Moderate
Histone deacetylase inhibition (https://aopwiki.org/relationships/2010)	non-adjacent	Spermatocyte depletion	Moderate	Moderate
Histone deacetylase inhibition (https://aopwiki.org/relationships/1717)	non-adjacent	Testicular atrophy	Moderate	Moderate

Stressors

Name	Evidence
Methoxyacetic acid	High
Butyrate	High

Name	Evidence
Trichostatin A	High
Valproate	Moderate

Overall Assessment of the AOP

1. Support for Biological Plausibility of KERs	
MIE => KE1: Histone deacetylase inhibition leads to histone acetylation increase	Biological Plausibility of the MIE => KE1 is high. Rationale: Upon the inhibition of HDAC by HDIs, the acetylation of lysine in histone remains and it leads to transcriptional activation or repression, changes in DNA replication and DNA damage repair. Activity of histone acetyltransferase (HAT) in testis nuclear protein was increased with MAA addition [Wade et al., 2008].
KE1 => KE2: Histone acetylation, increase leads to cell cycle, disrupted	Biological Plausibility of the KE1 => KE2 is moderate. Rationale: Gene transcription is regulated by histone acetylation [Struhl, 1998]. Acetylation of histones neutralizes the positive charge of the histones. Thus, less compacted DNA can be bound more easily by transcription factors and transcribed. In the models proposed for the relationship between histone acetylation and transcription, histone acetylation can be untargeted and occur at both promoter and nonpromoter regions, targeted generally to promoter regions, or targeted to specific promoters by gene-specific activator proteins [Richon et al., 2000; Struhl, 1998].
KE2 => KE3: Cell cycle, disrupted leads to apoptosis	Biological Plausibility of the KE2 => KE3 is moderate. Rationale: Prolonged cell cycle arrest will lead to either senescence or apoptosis. Especially for fast dividing and still differentiating cells, such an arrest will most certainly induce apoptosis as the normal cellular program cannot be followed.
KE3 => KE4: Apoptosis leads to spermatocyte depletion	Biological Plausibility of the KE3 => KE4 is moderate. Rationale: During development and in tissue homeostasis, apoptosis is needed to control organ size. If apoptosis is induced at a higher rate, one can assume it leading to atrophy of the target organ. Especially when target organ / target cells are fast replicating, abnormal levels of apoptosis will lead to depletion.
KE4 => AO: Spermatocyte depletion leads to testicular atrophy	Biological Plausibility of the KE4 => AO is moderate. Rationale: Spermatocyte depletion is one of the main characteristics of testicular atrophy.
2. Support for essentiality of KEs	
KE2: Cell cycle, disrupted	Essentiality of the KE2 is moderate. Rationale for Essentiality of KEs in the AOP: HDAC1-deficient embryonic stem cells showed reduced proliferation rates, which correlates with decreased cyclin-associated kinase activities and elevated levels of the cyclin-dependent kinase inhibitor 1A, a cell cycle regulator p21 [Lagger et al., 2002]. Loss of HDAC1 leads to significantly reduced overall deacetylase activity, hyperacetylation of a subset of histones H3 and H4 [Lagger et al., 2002].
3. Empirical support for KERs	

MIE => KE1: Histone deacetylase inhibition leads to histone acetylation, increase	Empirical Support of the MIE => KE1 is high. Rationale: HDAC inhibitors increase histone acetylation in brain [Schroeder et al., 2013]. The major empirical evidence came from the incubation of cell culture cells with small molecule compounds that inhibit HDAC enzymes followed by western blots or acid urea gel analysis. The first evidence was shown by Riggs et al. who showed that incubation of HeLa cells with <i>n</i> -butyrate leads to an accumulation of acetylated histone proteins [Riggs et al., 1977]. Later, it was shown that <i>n</i> -butyrate specifically increases the acetylation of histone by the incorporation of radioactive [³ H]acetate and analysis of the histones on acid urea gels that allow the detection of acetylated histones [Cousens et al., 1979]. TSA was shown to be an HDAC inhibitor by acid urea gel analysis in 1990 [Yoshida et al., 1990] and good evidence for VPA as an HDAC inhibitor in vitro and in vivo was shown using acetyl-specific antibodies and western blot [Gottlicher et al., 2001].
KE1 => KE2: Histone acetylation, increase leads to cell cycle, disrupted	Empirical Support of the KE1 => KE2 is moderate. Rationale: Increase in histone acetylation by HDAC inhibition induces the cell cycle regulator expression and inhibits progression through the cell cycle. Histone acetylation regulates the gene transcriptional mechanism [Struhl, 1998]. Acetylation of histones promotes the RNA polymerase reaction [Allfrey et al., 1964; Pogo et al., 1966]. Since several results supported a model in which increased histone acetylation is targeted to specific gene and occurs throughout the entire genome, not just the promoter regions, histone acetylation may lead to gene transcription of the cell cycle regulator [Richon et al., 2000].
KE2 => KE3: Cell cycle, disrupted leads to apoptosis	Empirical Support of the KE2 => KE3 is moderate. Rationale: Cell cycle arrest such as G ₁ arrest and G ₁ /S arrest are observed in apoptosis [Li et al., 2012; Dong et al., 2010]. microRNA-1 and microRNA-206 represses CCND2, while microRNA-29 represses CCND2 and induces G ₁ arrest and apoptosis in rhabdomyosarcoma [Li et al., 2012].
KE3 => KE4: Apoptosis leads to spermatocyte depletion	Empirical Support of the KE3 => KE4 is high. Rationale: MicroRNA-21 regulates the spermatogonial stem cell homeostasis, in which suppression of microRNA-21 with anti-miR-21 oligonucleotides led to apoptosis of spermatogonial stem cell-enriched germ cell cultures and the decrease in the number of spermatogonial stem cells [Niu et al., 2011].
KE4 => AO: Spermatocyte depletion leads to testicular atrophy	Empirical Support of the KE4 => AO is high. Rationale: The testicular atrophy seen in 2-methoxyethanol (2-ME), or its major metabolite MAA, treated rats in vivo and in human, and rat in vitro culture was associated with spermatocyte depletion [Beattie et al., 1984].

Domain of Applicability

Life Stage Applicability

Life Stage	Evidence
Adult, reproductively mature	High

Taxonomic Applicability

Term	Scientific Term	Evidence	Links
human	<i>Homo sapiens</i>	Moderate	NCBI (http://www.ncbi.nlm.nih.gov/Taxonomy/Browser/wwwtax.cgi?mode=Info&id=9606)

Term	Scientific Term	Evidence	Links
mouse	<i>Mus musculus</i>	Moderate	NCBI (http://www.ncbi.nlm.nih.gov/Taxonomy/Browser/wwwtax.cgi?mode=Info&id=10090)
rat	<i>Rattus norvegicus</i>	High	NCBI (http://www.ncbi.nlm.nih.gov/Taxonomy/Browser/wwwtax.cgi?mode=Info&id=10116)

Sex Applicability

Sex	Evidence
Male	High

The AOP is applicable to the reproductively mature males in rats, mice and humans. The administration route or doses of HDAC inhibitors may affect the intensity of the toxicity.

Essentiality of the Key Events

Key Event	Direct/Indirect Evidence
MIE: Histone deacetylase inhibition	HDAC inhibition induced testicular toxicity including testis atrophy [Miller et al., 1982]. HDAC inhibition in cell culture resulted in the testicular toxicity including germ cell apoptosis and cell morphology change [Li et al., 1996].
KE1: Histone acetylation, increase	The HDAC inhibition induced cell death in spermatocytes in both rat and human seminiferous tubules [Li et al., 1996].
KE2: Cell cycle, disrupted	In HDAC1-/- fibroblast lines, increase in the amount of cells in G1 phase and decrease in the amount of cells in S phase were observed, which indicates the importance of HDAC inhibition in cell cycle regulation [Zupkovitz et al., 2010].
KE3: Apoptosis	HDAC inhibition leads to cell death through the apoptotic pathways [Falkenberg et al., 2014].
KE4: Spermatocyte depletion	The HDAC inhibition induced cell death in spermatocytes in both rat and human seminiferous tubules [Li et al., 1996]. The HDAC inhibitor treatment resulted in degeneration in spermatocytes in rat seminiferous tubules [Li et al., 1996]. The HDAC inhibition induced the germ cell apoptosis in human testicular tissues [Li et al., 1996].

Weight of Evidence Summary

Biological plausibility, coherence, and consistency of the experimental evidence

The available data supporting the AOP are logic, coherent and consistent with established biological knowledge, whereas there are possibilities for alternative pathways.

Alternative mechanism(s) that logically present themselves and the extent to which they may distract from the postulated AOP

There are some other important apoptotic pathways that are involved in cell death, as well as other important spermatocyte signaling or mechanism influences testicular toxicity.

p53 pathway

The study in which *in vivo* administration of trichostatin A (TSA), a HDI, in mice resulted in male meiosis impairment showed the involvement of p53-noxa-caspase-3 apoptotic pathway in TSA-induced spermatocyte apoptosis [Fenic et al., 2008]. Other study showed that MAA induced up-regulation of p21 expression is mediated through histone hyperacetylation and independent of p53/p63/p73 [Parajuli et al., 2014].

NF-kappaB pathway

The present AOP focuses on p21 pathway leading to apoptosis, however, the alternative pathway such as NF-kappaB signaling pathways may be involved in apoptosis of spermatocytes [Wang et al., 2017].

Communication with Sertoli cells

The present AOP focuses on testicular atrophy by HDAC inhibition-induced apoptosis in spermatocytes, however, the signaling in Sertoli cells may be involved in testicular atrophy. Sertoli cell secretes GDNF, FGF2, CXCL12 or Ccl9 molecules, which results in the activation of RET, FGFR, CXCR4 or CCR1 signaling in spermatogonial stem cells, respectively [Chen and Liu, 2015].

Decrease in deoxynucleotide pool by MAA

MAA induces decrease in deoxynucleotide pool, resulting apoptosis, which may be an alternative pathway other than p21-mediated pathway [Yamazoe et al., 2015]. Inhibition of 5,10-CH₂-THF production by MAA may decreases deoxynucleotide pool in spermatocytes [Yamazoe et al., 2015].

Spermatocyte depletion by necrosis

Spermatocyte may be decreased by necrosis. Cell death mechanism other than apoptosis, such as necrosis, may be considered for spermatocyte depletion.

Quantitative Consideration

Concordance of dose-response relationships

This is a quantitative description on dose-response relationships from MIE to AOP. But some KE relationships individually are not fully supported with dose-response relationships, while there is empirical evidence to support that a change in KEup leads to an appropriate change in the respective KEdown.

Temporal concordance among the key events and adverse outcome

Temporal concordance between MIE and AOP has been described with in vivo experimental data. Empirical evidences show temporal concordance between MIE and the individual KEs, however, the temporal concordance among the individual KEs and AO is not fully elucidated.

Strength, consistency, and specificity of association of adverse outcome and initiating event

The scientific evidence on the linkage between MIE and AO has been described.

The quantitative understanding of the AOP in terms of indirect relations between HDAC inhibition and testicular atrophy was examined in in vivo experiments [Foster et al., 1983; Miller et al., 1982].

Considerations for Potential Applications of the AOP (optional)

The present AOP can be used in risk assessment of several types of the molecules including HDAC inhibitors as the anti-cancer drugs, as well as another types of the chemicals, in terms of testicular toxicity. HDAC inhibitors nowadays have been utilized as therapeutics for cancer or neurology disease, and the adverse effects of HDAC inhibitors should be evaluated. This AOP elucidating the pathway from HDAC inhibition to testicular atrophy may provides important insights for potential toxicity of HDAC inhibitors. It also provides a basis for the HDAC inhibition-induced epigenetic alteration and cell death. HDAC inhibitors such as rocicilostat are clinically evaluated as anti-cancer drugs in clinical trial.

References

Allfrey, V. et al. (1964), "Acetylation and methylation of histones and their possible role in the regulation of RNA synthesis", Proc Natl Acad Sci 51:786-794

Beattie, P.J. et al. (1984), "The effect of 2-methoxyethanol and methoxyacetic acid on Sertoli cell lactate production and protein synthesis in vitro", Toxicol Appl Pharmacol 76:56-61

Chen, S. and Liu, Y. (2015), "Regulation of spermatogonial stem cell self-renewal and spermatocyte meiosis by Sertoli cell signaling", Reproduction 149:R159-R167

Cousens, L.S., et al. (1979), "Different accessibilities in chromatin to histone acetylase", J Biol Chem 254:1716-1723

Dong, Q. et al. (2010), "microRNA let-7a inhibits proliferation of human prostate cancer cells in vitro and in vivo by targeting E2F2 and CCND2", PLoS One 5:e10147

Fenic, I. et al. (2008), "In vivo application of histone deacetylase inhibitor trichostatin-A impairs murine male meiosis", J Andro 29:172-185

Foster, P.M. et al. (1983), "Testicular toxicity of ethylene glycol monomethyl and monoethyl ethers in the rat", Toxicol Appl Pharmacol 69:385-39

Gottlicher, M. et al. (2001), "Valproic acid defines a novel class of HDAC inhibitors inducing differentiation of transformed cells", EMBO J

20:6969-6978

Lagger, G. et al. (2002), "Essential function of histone deacetylase 1 in proliferation control and CDK inhibitor repression", *EMBO J* 21:2672-2681

Li, L. et al. (2012), "Downregulation of microRNAs miR-1, -206 and -29 stabilizes PAX3 and CCND2 expression in rhabdomyosarcoma", *Lab Invest* 92:571-583

Li, L.H. et al. (1996), "2-Methoxyacetic acid (MAA)-induced spermatocyte apoptosis in human and rat testes: an in vitro comparison", *J Androl* 17:538-549

Miller, R.R. et al. (1982), "Toxicity of methoxyacetic acid in rats", *Fundam Appl Toxicol* 2:158-160

Niu, Z. et al. (2011), "microRNA-21 regulates the self-renewal of mouse spermatogonial stem cells", *Proc Natl Acad Sci* 108:12740-12745

Parajuli, K.R. et al. (2014), "Methoxyacetic acid suppresses prostate cancer cell growth by inducing growth arrest and apoptosis", *Am J Clin Exp Urol* 2:300-312

Pogo, B. et al. (1966), "RNA synthesis and histone acetylation during the course of gene activation in lymphocytes", *Proc Natl Acad Sci* 55:805-812

Richon, V.M. et al. (2000), "Histone deacetylase inhibitor selectively induces p21WAF1 expression and gene-associated histone acetylation", *Proc Natl Acad Sci* 97:10014-10019

Riggs, M.G. et al. (1977), "N-butyrate causes histone modification in HeLa and friend erythroleukaemia cells", *Nature* 268:462-464

Schroeder, F.A. et al. (2013), "A selective HDAC 1/2 inhibitor modulates chromatin and gene expression in brain and alters mouse behavior in two mood-related tests", *PLoS One* 8:e71323

Struhl, K. (1998), "Histone acetylation and transcriptional regulatory mechanisms", *Gene Dev* 12:599-606

Wade, M.G. et al. (2008), "Methoxyacetic acid-induced spermatocyte death is associated with histone hyperacetylation in rats", *Biol Reprod* 78:822-831

Wang, C. et al. (2017), "CD147 regulates extrinsic apoptosis in spermatocytes by modulating NFkB signaling pathways", *Oncotarget* 8:3132-3143

Yamazoe, Y. et al. (2015), "Embryo- and testicular-toxicities of methoxyacetate and the related: a review on possible roles of one-carbon transfer and histone modification", *Food Safety* 3:92-107

Yoshida, M. et al. (1990), "Potent and specific inhibition of mammalian histone deacetylase both in vivo and in vitro trichostatin A", *J Biol Chem* 265:17174-17179

Zupkovitz, G. et al. (2010), "The cyclin-dependent kinase inhibitor p21 is a crucial target for histone deacetylase 1 as a regulator of cellular proliferation", *Mol Cell Biol* 30:1171-1181

Appendix 1

List of MIEs in this AOP

Event: 1502: Histone deacetylase inhibition (<https://aopwiki.org/events/1502>)

Short Name: Histone deacetylase inhibition

Key Event Component

Process	Object	Action
enzyme inhibitor activity	histone deacetylase 1	decreased

AOPs Including This Key Event

AOP ID and Name	Event Type
Aop:212 - Histone deacetylase inhibition leading to testicular atrophy (https://aopwiki.org/aops/212)	MolecularInitiatingEvent
Aop:274 - Histone deacetylase inhibition leads to impeded craniofacial development (https://aopwiki.org/aops/274)	MolecularInitiatingEvent
Aop:275 - Histone deacetylase inhibition leads to neural tube defects (https://aopwiki.org/aops/275)	MolecularInitiatingEvent

Stressors

Name
Methoxyacetic acid
Butyrate
Trichostatin A
Valproic acid
Suberoylanilide hydroxamic acid
MS-275
Apicidin

Biological Context

Level of Biological Organization
Molecular
Cell term
Cell term
cell

Organ term

Organ term
organ

Evidence for Perturbation by Stressor

Overview for Molecular Initiating Event

HDIs are classified according to chemical nature and mode of mechanism: the short chain fatty acids (e.g., butyrate, valproate), hydroxamic acids (e.g., suberoylanilide hydroxamic acid or SAHA, Trichostatin A or TSA), cyclic tetrapeptides (e.g., FK-228), benzamides (e.g., N-acetylinaline and MS-275) and epoxides (depeudecin, trapoxin A) [Richon et al., 2003; Ropero and Esteller, 2007; Villar-Garea et al., 2004]. There is a report showing that TSA and butyrate competitively inhibits HDAC activity [Sekhavat et al., 2007]. HDIs inhibit preferentially HDACs with some selectiveness [Hu et al., 2003]. TSA inhibits HDAC1, HDAC3 and HDAC8, whereas MS-27-275 has inhibitory effect for HDAC1 and HDAC3 (IC₅₀ value of ~0.2 mM and ~8 mM, respectively), but no effect for HDAC8 (IC₅₀ value >10 mM) [Hu et al., 2003]. TSA inhibits HDAC1, 2, 3 of class I HDACs. [Damaskos et al., 2016].

Domain of Applicability

Taxonomic Applicability

Term	Scientific Term	Evidence	Links
rat	Rattus norvegicus	High	NCBI (http://www.ncbi.nlm.nih.gov/Taxonomy/Browser/wwwtax.cgi?mode=Info&id=10116)
human	Homo sapiens	High	NCBI (http://www.ncbi.nlm.nih.gov/Taxonomy/Browser/wwwtax.cgi?mode=Info&id=9606)
mouse	Mus musculus	High	NCBI (http://www.ncbi.nlm.nih.gov/Taxonomy/Browser/wwwtax.cgi?mode=Info&id=10090)

Life Stage Applicability

Life Stage	Evidence
All life stages	Moderate

Sex Applicability

Sex	Evidence
Unspecific	High

The inhibition of HDAC by HDIs is well conserved between species from lower organism to mammals.

- HDAC inhibition restores the rate of resorption of subretinal blebs in hyper glycemia in brown Norway rat and HDAC activity was inhibited with HDIs in human ARPE19 cells [Desjardins et al., 2016].
- Treatment of HDIs inducing HDAC inhibition showed anti-tumor effects in human non-small cell lung cancer cells [Ansari et al., 2016; Miyanaga et al., 2008].
- HDAC acetylation level was increased by HDIs in MRL-lpr/lpr murine model of lupus splenocytes [Mishra et al., 2003].
- SAHA increased histone acetylation in brain and spleen of mice [Hockly et al., 2003].
- MAA inhibits HDAC activity in HeLa cells and spleens from C57BL/6 mice [Jansen et al., 2004].
- It is also reported that MAA inhibits HDAC activity in testis cytosolic and nuclear extract of juvenile rats (27 days old) [Wade et al., 2008].
- VPA and TSA inhibit HDAC enzymatic activity in mouse embryo and human HeLa cell nuclear extract [Di Renzo et al., 2007].
- The treatment with HDAC inhibitors, phenylbutyrate (PB) (2 mM) and TSA (200 nM), inhibits HDAC in adjuvant arthritis synovial cells derived from rats, causing higher acetylated histone [Chung et al., 2003].

Key Event Description

Nucleosomes consist of eight core histones, two of each histone H2A, H2B, H3 and H4 [Damaskos et al., 2017]. DNA strands (about 200 bp) wind around the core histones, which can be modified on their N-terminal ends. One possible modification is the acetylation of lysine residues, which decreases the binding strength between DNA and the core histone. Histone deacetylases (HDACs) hydrolyze the acetyl residues [Damaskos et al., 2017]. HDACs remove the acetyl groups from the lysine residues leading to the formation of a condensed and transcriptionally silenced chromatin. Thus, the inhibition of HDAC blocks this action and can result in hyperacetylation of histones associated mostly with increases in transcriptional activation. Histone deacetylase inhibitor (HDI) inhibits HDAC, causing increased acetylation of the histones and thereby facilitating binding of transcription factors [Taunton et al., 1996].

It is known that eukaryotic HDAC isoforms are classified into four classes: class I HDACs (isoforms 1, 2, 3, 8), class II HDACs (isoforms 4, 5, 6, 7, 9, 10), class III HDACs (the sirtuins), and HDAC11 [Gregoretti et al., 2004; Weichert, 2009; Barneda-Zahonero and Parra, 2012]. HDACs 1, 2 and 3 are ubiquitously expressed, whereas HDAC8 is predominantly expressed in cells with smooth muscle/myoepithelial differentiation [Weichert, 2009]. HDAC6 is not observed to be expressed in lymphocytes, stromal cells and vascular endothelial cells [Weichert, 2009]. Class III HDACs, sirtuins, are widely expressed and localized in different cellular compartments [Barneda-Zahonero and Parra, 2012]. SirT1 is highly expressed in testis, thymus and multiple types of germ cells [Bell et al., 2014]. HDAC11 expression is enriched in kidney, brain, testis, heart and skeletal muscle [Barneda-Zahonero and Parra, 2012]. The members of groups 1,2 and 4 are dependent on a zinc ion and a water molecule at their active sites, for their deacetylase function. The Sirtuins of class 3 depend on NAD+, and are considered impervious to most known HDAC inhibitors [Drummond et al., 2005].

Several structurally distinct groups of compounds have been found to inhibit HDACs of class 1, 2 and 4, among others short-chain fatty acids (e.g. butyrate and VPA), hydroxamic acids (e.g. TSA and SAHA) and epoxyketones (e.g. Trichostatin A) [Drummond et al., 2005]. The hydroxamic acids seem to exert their inhibitory action by mimicking the acetyl-lysine target of HDACs, chelating the zinc ion in the active site and displacing the water molecule [Finnin et al., 1999]. Several recent high resolution crystal structures support this mode of inhibition [Decroos et al., 2015; Luckhurst et al., 2016]. The mode of inhibition of epoxyketones seems to function the formation of a stable transition state analog with the zinc ion in the active site [Porter and Christianson, 2017]. The inhibitory actions of the short-chain fatty acids are less well defined, but it has been speculated that VPA blocks access to the binding pocket [Göttlicher et al., 2001]. It has been shown that VPA exert similar gene regulatory effects to TSA, on a panel of migration related transcripts in neural crest cells [Dreser et al., 2015] supporting a mode of action similar to hydroxamic acid type HDAC inhibitors.

How it is Measured or Detected

The measurement of HDAC inhibition monitors changes in histone acetylation. HDAC inhibition can be detected directly by the measurement of HDAC activity using commercially available colorimetric or fluorimetric kits or indirectly by increase of histone acetylation as the detection of global histone acetylation changes by Western blot or mass spectrometry (MS)-based proteomic methods or as detection of site-specific histone acetylation changes using chromatin immunoprecipitation (ChIP) or ChIP-on-Chip. The measurement methods include the immunological detection of histone acetylation with anti-acetylated histone antibodies [Richon et al., 2004]. The histones are isolated from pellets of cells treated with HDIs, followed by acid-urea-triton gel electrophoresis, western blotting, and immunohistochemistry [Richon et al., 2003]. The HDAC activity is measured directly with ultra high performance liquid chromatography-electrospray ionization-tandem mass spectrometry (UHPLC-ESI-MS/MS) by calculating the ratio of deacetylated peptide and acetylated peptide [Zwick et al., 2016]. Some *in silico* methods including molecular modelling, virtual screening and molecular dynamics are used to find the common HDAC inhibitor structures [Huang et al., 2016; Yanuar et al. 2016].

References

Ansari, J. et al. (2016), "Epigenetics in non-small cell lung cancer: from basics to therapeutics", *Transl Lung Cancer Res* 5:155-171

Barneda-Zahonero, B. and Parra, M. (2012), "Histone deacetylases and cancer", *Mol Oncol* 6:579-589

Bell, E.L. et al. (2014), "SirT1 is required in the male germ cell for differentiation and fecundity in mice", *Development* 141:3495-3504

Chung, Y.L. et al. (2003), "A therapeutic strategy uses histone deacetylase inhibitors to modulate the expression of genes involved in the pathogenesis of rheumatoid arthritis", *Mol Ther* 8:707-717

Damaskos, C. et al. (2016), "Histone deacetylase inhibitors: a novel therapeutic weapon against medullary thyroid cancer?", *Anticancer Res* 36:5019-5024

Damaskos, C. et al. (2017), "Histone deacetylase inhibitors: an attractive therapeutic strategy against breast cancer", *Anticancer Research* 37: 35-46.

Decroos, C. et al. (2015), "Biochemical and structural characterization of HDAC8 mutants associated with cornelia de lange syndrome spectrum disorders", *Biochemistry* 54: 6501-6513.

Desjardins, D. et al. (2016), "Histone deacetylase inhibition restores retinal pigment epithelium function in hyperglycemia", *PLoS ONE* 11: e0162596

Di Renzo, F. et al. (2007), "Boric acid inhibits embryonic histone deacetylases: A suggested mechanism to explain boric acid-related teratogenicity", *Toxicol and Appl Pharmacol* 220:178-185

Dreser, N. et al. (2015), "Grouping of histone deacetylase inhibitors and other toxicants disturbing neural crest migration by transcriptional profiling", *Neurotoxicology* 50: 56-70

Drummond, D.C. et al. (2005), "Clinical development of histone deacetylase inhibitors as anticancer agents", *Annu Rev Pharmacol Toxicol* 45: 495-528

Finnin, M.S. et al. (1999), "Structures of a histone deacetylase homologue bound to the TSA and SAHA inhibitors", *Nature* 401: 188-193

Göttlicher, M. et al. (2001), "Valproic acid defines a novel class of HDAC inhibitors inducing differentiation of transformed cells", *EMBO J* 20: 6969-6978

Gregoretti, I.V. et al. (2004), "Molecular evolution of the histone deacetylase family: functional implications of phylogenetic analysis", *J Mol Biol* 338: 17-31

Hockly, E. et al. (2003), "Suberoylanilide hydroxamic acid, a histone deacetylase inhibitor, ameliorates motor deficits in a mouse model of Huntington's disease", *Proc Nat Acad Sci* 100:2041-2046

Hu, E. et al. (2003), "Identification of novel isoform-selective inhibitors within class I histone deacetylases", *J Pharmacol Exp Ther* 307:720-728

Huang, Y.X. et al. (2016), "Virtual screening and experimental validation of novel histone deacetylase inhibitors", *BMC Pharmacol Toxicol* 17(1):32

Jansen, M.S. et al. (2014), "Short-chain fatty acids enhance nuclear receptor activity through mitogen-activated protein kinase activation and histone deacetylase inhibition", *Proc Natl Acad Sci USA* 101(18):7199-7204

Luckhurst, C.A. et al. (2016), "Potent, Selective, and CNS-Penetrant Tetrasubstituted Cyclopropane Class IIa Histone Deacetylase (HDAC) Inhibitors", *ACS Med Chem Lett* 7:34-39

Mishra, N. et al. (2003), "Histone deacetylase inhibitors modulate renal disease in the MRL-lpr/lpr mouse", *J Clin Invest* 111:539-552

Miyanaga, A. et al. (2008), "Antitumor activity of histone deacetylase inhibitors in non-small cell lung cancer cells: development of a molecular predictive model", *Mol Cancer Ther* 7:1923-1930

Porter, N.J., and Christianson, D.W. (2017), "Binding of the microbial cyclic tetrapeptide trapoxin A to the Class I histone deacetylase HDAC8", *ACS Chem Biol* 12:2281-2286

Richon, V.M. et al. (2003), "Histone deacetylase inhibitors: assays to assess effectiveness in vitro and in vivo", *Methods Enzymol.* 376:199-205

Ropero, S. and Esteller, M. (2007), "The role of histone deacetylases (HDACs) in human cancer", *Mol Oncol* 1:19-25

Sekhavat, A. et al. (2007), "Competitive inhibition of histone deacetylase activity by trichostatin A and butyrate", *Biochemistry and Cell Biology* 85:751-758

Taunton, J. et al. (1996), "A mammalian histone deacetylase related to the Yeast transcriptional regulator Rpd3p", *Science* 272:408-411

Villar-Garea, A. and Esteller, M. (2004), "Histone deacetylase inhibitors: understanding a new wave of anticancer agents", *Int J Cancer* 112:171-178

Wade, M.G. et al. (2008), "Methoxyacetic acid-induced spermatocyte death is associated with histone hyperacetylation in rats", *Biol Reprod* 78:822-831

Weichert, W. (2009) "HDAC expression and clinical prognosis in human malignancies", *Cancer Letters* 280:168-176

Yanuar, A. et al. (2016), "In silico approach to finding new active compounds from histone deacetylase (HDAC) family", *Curr Pharm Des* 22: 3488-3497

Zwick, V. et al. (2016), "Cell-based multi-substrate assay coupled to UHPLC-ESI-MS/MS for a quick identification of class-specific HDAC inhibitors", *J Enzyme Inhib Med Chem* 31: 209-214

List of Key Events in the AOP

Event: 1503: Histone acetylation, increase (<https://aopwiki.org/events/1503>)

Short Name: Histone acetylation, increase

Key Event Component

Process	Object	Action
regulation of protein modification process	histone	increased

AOPs Including This Key Event

AOP ID and Name	Event Type
Aop:212 - Histone deacetylase inhibition leading to testicular atrophy (https://aopwiki.org/aops/212)	KeyEvent
Aop:275 - Histone deacetylase inhibition leads to neural tube defects (https://aopwiki.org/aops/275)	KeyEvent

Biological Context

Level of Biological Organization
Cellular

Cell term

Cell term
cell

Organ term

Organ term
organ

Domain of Applicability

Life Stage Applicability

Life Stage	Evidence
Not Otherwise Specified	Moderate

Sex Applicability

Sex	Evidence
Unspecific	High

The histone acetylation increase by HDIs is well conserved between species from lower organism to mammals.

MAA, a HDAC inhibitor, induces acetylation of histones H3 and H4 in Sprague-Dawley (*Rattus norvegicus*) [Wade et al., 2008].

It is also reported that MAA promotes acetylation of H4 in HeLa cells (*Homo sapiens*) and spleens from C57BL/6 mice (*Mus musculus*) treated with MAA [Jansen et al., 2014].

VPA, a HDAC inhibitor, induces hyperacetylation of histone H4 in protein extract of mouse embryos (*Mus musculus*) exposed *in utero* for 1h to VPA [Di Renzo et al., 2007a].

Apicidin, MS-275 and sodium butyrate, HDAC inhibitors, induce hyperacetylation of histone H4 in homogenates from mouse embryos (*Mus musculus*) after the treatments [Di Renzo et al., 2007b].

MAA acetylates histones H3K9 and H4K12 in limbs of CD1 mice (*Mus musculus*) [Dayan and Hales, 2014].

Key Event Description

Gene transcription is regulated with the balance between acetylation and deacetylation. A dynamic balance of histone acetylation and histone deacetylation are typically catalyzed by enzymes with histone acetyltransferase (HAT) and HDAC activities. Histone acetylation relaxes chromatin and makes it accessible to transcription factors, whereas deacetylation is associated with gene silencing. The balance can be disturbed also by targeting HAT, not only HDACs. At least theoretically, an activation of HAT might lead to increase in histone acetylation. The acetylation and deacetylation are modulated on the NH3+ groups of lysine amino acid residues in histones. DNA in acetylated histones is more accessible for transcription factors, leading to increase in gene expression. HDAC inhibition promotes the hyperacetylation by inhibiting deacetylation of histones with classes of H2A, H2B, H3 and H4 in nucleosomes. [Wade et al., 2008]. The inhibition of HDACs result in an accumulation of acetylated proteins such as tubulin or histones.

How it is Measured or Detected

Histone acetylation is measured by the immunological detection of histone acetylation with anti-acetylated histone antibodies [Richon et al., 2004]. Histone acetylation on chromatin can be measured using labeling method with sodium [³H]acetate [Gunjan et al., 2001]. The histone acetylation increase is detected as global histone acetylation changes by Western blot or mass spectrometry (MS)-based proteomic methods or as site specific histone acetylation changes using chromatin immunoprecipitation (ChIP) or ChIP-on-Chip.

References

Dayan, C. and Hales, B.F. (2014), "Effects of ethylene glycol monomethyl ether and its metabolite, 2-methoxyacetic acid, on organogenesis stage mouse limbs in vitro", Birth Defects Res (Part B) 101:254-261

Di Renzo, F. et al. (2007a), "Boric acid inhibits embryonic histone deacetylases: A suggested mechanism to explain boric acid-related teratogenicity", Toxicol and Appl Pharmacol 220:178-185

Di Renzo, F. et al. (2007b), "Relationship between embryonic histone hyperacetylation and axial skeletal defects in mouse exposed to the three HDAC inhibitors apicidin, MS-275, and sodium butyrate", Toxicol Sci 98:582-588

Gunjan, A. et al. (2001), "Core histone acetylation is regulated by linker histone stoichiometry in vivo", J Biol Chem 276:3635-3640

Jansen, M.S. et al. (2014), "Short-chain fatty acids enhance nuclear receptor activity through mitogen-activated protein kinase activation and histone deacetylase inhibition", Proc Natl Acad Sci USA 101:7199-7204

Richon, V.M. et al. (2004), "Histone deacetylase inhibitors: assays to assess effectiveness in vitro and in vivo", Methods Enzymol 376:199-205

Wade, M.G. et al. (2008), "Methoxyacetic acid-induced spermatocyte death is associated with histone hyperacetylation in rats", Biol Reprod 78:822-831

Event: 1505: Cell cycle, disrupted (<https://aopwiki.org/events/1505>)

Short Name: Cell cycle, disrupted

Key Event Component

Process	Object	Action
regulation of cell cycle	cell cycle-related cyclin	disrupted

AOPs Including This Key Event

AOP ID and Name	Event Type
Aop:212 - Histone deacetylase inhibition leading to testicular atrophy (https://aopwiki.org/aops/212)	KeyEvent

Biological Context

Level of Biological Organization
Cellular

Cell term

Cell term
cell

Organ term

Organ term
organ

Domain of Applicability

Taxonomic Applicability

Term	Scientific Term	Evidence	Links
Homo sapiens	Homo sapiens	High	NCBI (http://www.ncbi.nlm.nih.gov/Taxonomy/Browser/wwwtax.cgi?mode=Info&id=9606)
Mus musculus	Mus musculus	High	NCBI (http://www.ncbi.nlm.nih.gov/Taxonomy/Browser/wwwtax.cgi?mode=Info&id=10090)

Life Stage Applicability

Life Stage	Evidence
Not Otherwise Specified	Moderate

Sex Applicability

Sex	Evidence
Unspecific	High

The histone gene expression alters in each phase of cell cycle in human HeLa cell (*Homo sapiens*) [Heintz et al., 1982].

Key Event Description

The disruption of the cell cycle leads to a decrease in the cell number. The cell cycle consists of G₁, S, G₂, M, and G₀ phase. The cell cycle regulation is disrupted by the cell cycle arrest in certain cell cycle phase. The histone gene expression is regulated in cell cycle phases [Heintz et al., 1983].

How it is Measured or Detected

The percentage of cells at G₁, G₀, S, and G₂/M phases can be detected by flow cytometry [Li et al., 2013]. Cell cycle distribution was analyzed by fluorescence-activated cell sorter (FACS) analysis with a Partec PAS-II sorter [Zupkovitz et al., 2010]. The four cell cycle phases in living cells can be measured with four-color fluorescent proteins using live cell imaging [Bajar et al., 2016]. The incorporation of [³H]deoxycytidine or [³H]thymidine into cell DNA during S phase can be monitored as DNA synthesis [Heintz et al., 1982].

References

Bajar, B.T. et al. (2016), "Fluorescent indicators for simultaneous reporting of all four cell cycle phases", *Nat Methods* 13:993-996

Heintz, N. et al. (1983), "Regulation of human histone gene expression: Kinetics of accumulation and changes in the rate of synthesis and in the half-lives of individual histone mRNAs during the HeLa cell cycle", *Molecular and Cellular Biology* 3:539-550

Li, Q. et al. (2013), "Glyphosate and AMPA inhibit cancer cell growth through inhibiting intracellular glycine synthesis", *Drug Des Devel Ther* 7:635-643

Event: 1262: Apoptosis (<https://aopwiki.org/events/1262>)

Short Name: Apoptosis

Key Event Component

Process	Object	Action
apoptotic process		increased

AOPs Including This Key Event

AOP ID and Name	Event Type
Aop:205 - AOP from chemical insult to cell death (https://aopwiki.org/aops/205)	AdverseOutcome
Aop:207 - NADPH oxidase and P38 MAPK activation leading to reproductive failure in <i>Caenorhabditis elegans</i> (https://aopwiki.org/aops/207)	KeyEvent
Aop:212 - Histone deacetylase inhibition leading to testicular atrophy (https://aopwiki.org/aops/212)	KeyEvent
Aop:285 - Inhibition of N-linked glycosylation leads to liver injury (https://aopwiki.org/aops/285)	KeyEvent
Aop:268 - Uncoupling of oxidative phosphorylation leading to growth inhibition (6) (https://aopwiki.org/aops/268)	KeyEvent

Biological Context

Level of Biological Organization
Cellular

Cell term

Cell term
cell

Organ term

Organ term
organ

Domain of Applicability

Life Stage Applicability

Life Stage	Evidence
Not Otherwise Specified	High

Sex Applicability

Sex	Evidence
Unspecific	High

Apoptosis is induced in human prostate cancer cell lines (*Homo sapiens*) [Parajuli et al., 2014].

Apoptosis occurs in B6C3F1 mouse (*Mus musculus*) [Elmore, 2007].

Apoptosis occurs in Sprague-Dawley rat (*Rattus norvegicus*) [Elmore, 2007].

Apoptosis occurs in nematode (*Caenorhabditis elegans*) [Elmore, 2007].

Key Event Description

Apoptosis, the process of programmed cell death, is characterized by distinct morphology with DNA fragmentation and energy dependency [Elmore, 2007]. Apoptosis, also called as “physiological cell death”, is involved in cell turnover, physiological involution and atrophy of various tissues and organs [Kerr et al., 1972]. The formation of apoptotic bodies involves marked condensation of both nucleus and cytoplasm, nuclear fragmentation, and separation of protuberances [Kerr et al., 1972]. Apoptosis is characterized by DNA ladder and chromatin condensation. Several stimuli such as hypoxia, nucleotides deprivation, chemotherapeutical drugs, DNA damage, and mitotic spindle damage

induce p53 activation, leading to p21 activation and cell cycle arrest [Pucci et al., 2000]. The SAHA or TSA treatment on neonatal human dermal fibroblasts (NHDFs) for 24 or 72 hrs inhibited proliferation of the NHDF cells [Glaser et al., 2003]. Considering that the acetylation of histone H4 was increased by the treatment of SAHA for 4 hrs, histone deacetylase inhibition may be involved in the inhibition of the cell proliferation [Glaser et al., 2003]. The impaired proliferation was observed in HDAC1^{-/-} ES cells, which was rescued with the reintroduction of HDAC1 [Zupkovitz et al., 2010]. The present AOP focuses on p21 pathway leading to apoptosis, however, the alternative pathway such as NF- κ B signaling pathways may be involved in apoptosis of spermatocytes [Wang et al., 2017].

How it is Measured or Detected

Apoptosis is characterized by many morphological and biochemical changes such as cell shrinkage and pyknosis, cell fragments into apoptotic bodies, DNA fragmentation, enzymatic activation of pro-caspases or phosphatidylserine translocation that can be measured using electron and cytochemical optical microscopy, proteomic and genomic methods, spectroscopic techniques [Martines, 2010, Taatjes, 2008, Archana, 2013]. Some of the methods are described as follows:

The apoptosis is detected with the expression alteration of procaspases 7 and 3 by Western blotting using antibodies [Parajuli, 2014].

The apoptosis is measured with down-regulation of anti-apoptotic gene baculoviral inhibitor of apoptosis protein repeat containing 2 (BIRC2, or cIAP1) [Parajuli, 2014].

Apoptotic nucleosomes are detected using Cell Death Detection ELISA kit, which were calculated as absorbance subtraction at 405 nm and 490 nm [Parajuli, 2014].

Cleavage of PARP is detected with Western blotting [Parajuli, 2014].

Caspase-3 and caspase-9 activity is measured with the enzyme-catalyzed release of p-nitroanilide (pNA) and quantified at 405 nm [Wu, 2016].

Apoptosis is measured with Annexin V-FITC probes, and the relative percentage of Annexin V-FITC-positive/PI-negative cells is analyzed by flow cytometry [Wu et al., 2016].

Apoptosis is detected with the Terminal dUTP Nick End-Labeling (TUNEL) method to assay the endonuclease cleavage products by enzymatically end-labeling the DNA strand breaks [Kressel and Groscurth, 1994].

For the detection of apoptosis, the testes are fixed in neutral buffered formalin, and embedded in paraffin. Germ cell death is visualized in testis sections by Terminal dUTP Nick End-Labeling (TUNEL) staining method [Wade et al., 2008]. The incidence of TUNEL-positive cells is expressed as the number of positive cells per tubule examined for one entire testis section per animal [Wade et al., 2008].

References

Archana, M. et al. (2013), "Various methods available for detection of apoptotic cells", Indian J Cancer 50:274-283

Elmore, S. (2007), "Apoptosis: a review of programmed cell death", Toxicol Pathol 35:495-516

Glaser, K.B. et al. (2003), "Gene expression profiling of multiple histone deacetylase (HDAC) inhibitors: defining a common gene set produced by HDAC inhibition in T24 and MDA carcinoma cell lines", Mol Cancer Ther 2:151-163

Kerr, J.F.R. et al. (1972), "Apoptosis: a basic biological phenomenon with wide-ranging implications in tissue kinetics", Br J Cancer 26:239-257

Kressel, M. and Groscurth, P. (1994), "Distinction of apoptotic and necrotic cell death by in situ labelling of fragmented DNA", Cell Tissue Res 278:549-556

Martinez, M.M. et al. (2010), "Detection of apoptosis: A review of conventional and novel techniques", Anal Methods 2:996-1004

Parajuli, K.R. et al. (2014), "Methoxyacetic acid suppresses prostate cancer cell growth by inducing growth arrest and apoptosis", Am J Clin Exp Urol 2:300-313

Pucci, B. et al. (2000), "Cell cycle and apoptosis", Neoplasia 2:291-299

Taatjes, D.J. et al. (2008), "Morphological and cytochemical determination of cell death by apoptosis", Histochem Cell Biol 129:33-43

Wade, M.G. et al. (2008), "Methoxyacetic acid-induced spermatocyte death is associated with histone hyperacetylation in rats", Biol Reprod 78:822-831

Wang, C. et al. (2017), "CD147 regulates extrinsic apoptosis in spermatocytes by modulating NF κ B signaling pathways", Oncotarget 8:3132-3143

Wu, R. et al. (2016), "microRNA-497 induces apoptosis and suppressed proliferation via the Bcl-2/Bax-caspase9-caspase 3 pathway and cyclin D2 protein in HUVECs", PLoS One 11:e0167052

Zupkovitz, G. et al. (2010), "The cyclin-dependent kinase inhibitor p21 is a crucial target for histone deacetylase 1 as a regulator of cellular proliferation", Mol Cell Biol 30:1171-1181

Event: 1515: Spermatocyte depletion (<https://aopwiki.org/events/1515>)

Short Name: Spermatocyte depletion

Key Event Component

Process	Object	Action
	spermatocyte	decreased

AOPs Including This Key Event

AOP ID and Name	Event Type
Aop:212 - Histone deacetylase inhibition leading to testicular atrophy (https://aopwiki.org/aops/212)	KeyEvent

Biological Context

Level of Biological Organization
Tissue

Organ term

Organ term
testis

Domain of Applicability

Taxonomic Applicability

Term	Scientific Term	Evidence	Links
Rattus norvegicus	Rattus norvegicus	Moderate	NCBI (http://www.ncbi.nlm.nih.gov/Taxonomy/Browser/wwwtax.cgi?mode=Info&id=10116)

Life Stage Applicability

Life Stage	Evidence
Adult	Moderate

Sex Applicability

Sex	Evidence
Male	High

There are evidences of spermatocyte depletion in different species.

Mature sperm counts were decreased and the residual spermatozoa had reduced motility and decreased viability (*Mus musculus*) [Zindy et al., 2001].

The sperm counts in the cauda epididymis of rats were significantly decreased (*Rattus norvegicus*) [Oishi, 2001].

Spermatocyte death can be induced in Sprague-Dawley rats (*Rattus norvegicus*) [Wade et al., 2008].

Key Event Description

Spermatocytes are differentiated from spermatogonial stem cells via random proliferation, differentiation and synchronized mitoses with several stages [Rooij, 2001]. In several steps of the spermatogonial differentiation, different molecular mechanism is activated in testis [Rooij, 2001, de Kretser et al., 2016]. The stem cell factor (SCF) genes are involved in differentiation into A1 spermatogonia. The expression of cyclin D2 is regulated in the epithelial stage VIII when the aligned spermatogonia differentiate into A1 spermatogonia [Rooij, 2001]. Upon the apoptosis of spermatogonia, overexpression of the apoptosis-inhibiting proteins Bcl-2 and Bcl-xL and deficiency of the apoptosis-inducing protein Bax have been shown to cause an accumulation of spermatogonia in the testis, leading to apoptosis of all cells [Rooij, 2001].

How it is Measured or Detected

Traditional spermatocytes assessment includes sperm count and concentration (haemocytometer, automated imaged-based system), morphology

and motility (microscope, automated imaged-based system) and viability (for example propidium iodide staining of necrotic cells, TUNEL assay staining apoptotic cells). In addition, there are functional tests such as assays for genetic integrity (e.g. via measurement of DNA fragmentation/integrity -Halosperm kit or reactive oxygen species) and fertilization defects (through various measures of sperm-zona pellucida (ZP) interaction, such as measurement of ZP-receptor binding).

The sperm-containing fluid was squeezed out of the cauda, and suspended in medium containing HEPES buffer and bovine serum albumin, and incubated at 37°C for 20 min. The number of spermatozoa was determined by hematocytometer [Zindy et al., 2001].

Testicular sperm counts and daily sperm production were determined by counting the total number of spermatids per testis, and divided by the testicular weight to give the results in spermatids per gram of testis [Oishi, 2001].

For the testis cell analysis, fresh testes were dispersed using a two-stage enzymatic digestion and incubated in BSA containing collagenase and DNase I [Wade et al., 2006]. The seminiferous tubules were further digested and cells were fixed in ice-cold 70% ethanol [Wade et al., 2006]. Relative proportions of spermatogenic cell populations were assessed in fixed cells using a flow cytometric method [Wade et al., 2006]. The principle of the test is that spermatogenic cells, as they differentiate from normal diploid spermatogonial stem cells through to mature spermatozoa with a highly condensed haploid complement of DNA, progress through various intermediate stages with differing nuclear DNA content and cellular content of mitochondria. Relative proportions of cells in each population were calculated with WinList software [Wade et al., 2006].

References

de Kretser, D.M. et al. (2016), "Endocrinology: Adult and Pediatric (Seventh Edition)", W.B. Saunders, Chapter 136-Spermatogenesis, pages 2325-2353.e9, Editors: J. Larry Jameson, Leslie J De Groot, David M. de Kretser, Linda C. Giudice, Ashley B. Grossman, Shlomo Melmed, John T. Potts, Gordon C. Weir

Oishi, S. (2001), "Effects of butylparaben on the male reproductive system in rats", *Toxicol Indust Health* 17:31-39

Rooij, D.G. (2001), "Proliferation and differentiation of spermatogonial stem cells", *Reproduction* 121:347-354

Wade, M.G. et al. (2006), "Testicular toxicity of candidate fuel additive 1,6-dimethoxyhexane: comparison with several similar aliphatic ethers", *Toxicol Sci* 89:304-313

Wade, M.G. et al. (2008), "Methoxyacetic acid-induced spermatocyte death is associated with histone hyperacetylation in rats", *Biol Reprod* 78:822-831

Zindy, F. et al. (2001), "Control of spermatogenesis in mice by the cyclin D-dependent kinase inhibitors p18Ink4c and p19Ink4d", *Mol Cell Biol* 21:3244-3255

List of Adverse Outcomes in this AOP

Event: 1506: Testicular atrophy (<https://aopwiki.org/events/1506>)

Short Name: Testicular atrophy

Key Event Component

Process	Object	Action
Testicular atrophy	Testis	increased

AOPs Including This Key Event

AOP ID and Name	Event Type
Aop:212 - Histone deacetylase inhibition leading to testicular atrophy (https://aopwiki.org/aops/212)	AdverseOutcome

Biological Context

Level of Biological Organization
Organ

Organ term

Organ term
testis

Domain of Applicability

Taxonomic Applicability

Term	Scientific Term	Evidence	Links
Rattus norvegicus	Rattus norvegicus	Moderate	NCBI (http://www.ncbi.nlm.nih.gov/Taxonomy/Browser/wwwtax.cgi?mode=Info&id=10116)

Life Stage Applicability

Life Stage	Evidence
Adult, reproductively mature	Moderate

Sex Applicability

Sex	Evidence
Male	High

- The decrease in testis weight associated with testicular cell damage was induced by EGME or MAA treatment in rats (*Rattus norvegicus*) [Foster et al., 1983].
- The number of spermatocytes, principally pachytene cells, is decreased by EGME treatment in CD-1 mice (*Mus musculus*) and CD rats (*Rattus norvegicus*) [Anderson et al., 1987].
- The testicular lesions induced by 2-methoxyethanol (ethylene glycol monomethyl ether; EGME) were observed in rats (*Rattus norvegicus*) and guinea pigs (*Cavia porcellus*), which are different in onset, characteristics and severity [Ku et al., 1984].
- Spermatogenesis was disrupted by EGME treatment in rabbits (*Oryctolagus cuniculus*) [Foote et al., 1995].
- Testicular toxicity such as spermatocyte death in seminiferous tubule stages I-IV and stages XII-XIV was induced by dimethoxyhexane (DMH) treatment in Sprague-Dawley rats (*Rattus norvegicus*) [Wade et al., 2006].

Key Event Description

It is hypothesized that the testicular effects of 1,6-dimethoxyhexane (DMH) are caused by its metabolism to methoxyacetic acid (MAA) [Wade et al., 2006; Poon et al., 2004]. MAA produces testicular and thymic atrophy such as the decrease in size [Miller et al., 1982; Moss et al., 1985]. The spermatogenic stages in which the toxicity of MAA is induced are on the pachytene spermatocytes immediately before and during meiotic division, which are Stages XII-XIV of the cycle in the rat and the early pachytene spermatocytes at stages I-IV of the cycle. Dead germ cells can be seen as soon as 12 hours after the treatment of MAA [Casarett & Doull's, 7th edition].

How it is Measured or Detected

- Testicular atrophy can be assessed by testicular volume measurement using orchidometer, rulers, calipers, and ultrasonography or by testis weighing and histopathologic examination.
- The testis weight is measured to detect the testicular atrophy [Foster et al., 1983].
- The urinary zinc excretion and testicular zinc content are examined, since zinc concentration has been shown to play an important role in the production of testicular injury [Foster et al., 1983].
- The testicular tissue structure is observed whether there are normal germinal epithelial cells and Leydig cells [Mercantepe et al., 2018]. Testis is fixed for observations by light microscopy or transmission electron microscopy [McDowell and Trump, 1976; Mercantepe et al., 2018].
- Changes in sperm are measured by computer-assisted sperm analysis [Foote et al., 1995].
- For the assessment of sperm morphology, eosin-stained sperm collected from the cauda epididymis is observed. At least 200 sperm on each slide were examined for the proportion of sperm with abnormal head (overhooked, blunt hook, banana-shaped, amorphous, or extremely oversized) or tail (twisted, bent, corkscrew, double/multiple) by one individual unaware of animal number or treatment [Wade et al., 2006].
- For the measurement of the total number of condensed spermatids per testis, a weighed portion of the parenchyma from the left testis was homogenized [Wade et al., 2006]. Sperm or homogenization-resistant spermatid nuclei densities were calculated from the average number of nuclei and were expressed as total or as per gram of epididymis or testis weight [Wade et al., 2006].
- For the determination of total LDH and LDH-X in supernatant of the homogenized testis fragment, enzyme activity was measured by monitoring extinction of NAD absorbance [Wade et al., 2006].

Regulatory Significance of the AO

The testicular toxicity assessment is important for assessing the side effects of the medicines such as anti-cancer drugs, as well as hazard and risk of chemicals. The testicular atrophy including decrease in testis weight and sperm count, fertility, decrease in morphology and function of the sperm, can become one of the main endpoints as the adverse effects of the therapeutics. The unexpected effects of the therapeutics may be predicted with this Adverse Outcome.

References

Anderson, D. et al. (1987), "Effect of ethylene glycol monomethyl ether on spermatogenesis, dominant lethality, and F1 abnormalities in the rat and the mouse after treatment of F0 males", *Teratog Carcinog Mutagen* 7:141-158

Casarett & Doull's Toxicology, the Basic Science of Poisons, 7th Edition, Edited by Curtis D. Klaassen, Chapter 20 Toxic responses of the reproductive system

Foote, R.H. et al. (1995), "Ethylene glycol monomethyl ether effects on health and reproduction in male rabbits", *Reprod Toxicol* 9:527-539

Foster, P.M. et al. (1983), "Testicular toxicity of ethylene glycol monomethyl and monoethyl ethers in the rats", *Toxicol Appl Pharmacol* 69:385-399

Ku, W.W. et al. (1994), "Comparison of the testicular effects of 2-methoxyethanol (ME) in rats and guinea pigs", *Exp Mol Pathol* 61:119-133

McDowell, E.M. and Trump, B.F. (1976), "Histologic fixatives suitable for diagnostic light and electron microscopy", *Arch Pathol Lab Med* 100:405-414

Mercantepe, T. et al. (2018), "Protective effects of amifostine, curcumin and caffeic acid phenethyl ester against cisplatin-induced testis tissue damage in rats", *Exp Ther Med* 15:3404-3412

Miller, R. et al. (1982), "Toxicity of methoxyacetic acid in rats", *Fundam Appl Toxicol* 2:158-160

Moss, E.J. et al. (1985), "The role of metabolism in 2-methoxyethanol-induced testicular toxicity", *Toxicol Appl Pharmacol* 79:480-489

Poon, R. et al. (2004), "Short-term oral toxicity of pentyl ether, 1,4-diethoxybutane, and 1,6-dimethoxyhexane in male rats", *Toxicol Sci* 77:142-150

Wade, M.G. et al. (2006), "Testicular toxicity of candidate fuel additive 1,6-dimethoxyhexane: comparison with several similar aliphatic ethers", *Toxicol Sci* 89:304-313

Appendix 2

List of Key Event Relationships in the AOP

List of Adjacent Key Event Relationships

Relationship: 1709: Histone deacetylase inhibition leads to Histone acetylation, increase
(<https://aopwiki.org/relationships/1709>)

AOPs Referencing Relationship

AOP Name	Adjacency	Weight of Evidence	Quantitative Understanding
Histone deacetylase inhibition leading to testicular atrophy (https://aopwiki.org/aops/212)	adjacent	High	Moderate
Histone deacetylase inhibition leads to neural tube defects (https://aopwiki.org/aops/275)	adjacent	Not Specified	Not Specified

Evidence Supporting Applicability of this Relationship

Taxonomic Applicability

Term	Scientific Term	Evidence	Links
Homo sapiens	Homo sapiens	High	NCBI (http://www.ncbi.nlm.nih.gov/Taxonomy/Browser/wwwtax.cgi?mode=Info&id=9606)
Rattus norvegicus	Rattus norvegicus	High	NCBI (http://www.ncbi.nlm.nih.gov/Taxonomy/Browser/wwwtax.cgi?mode=Info&id=10116)
Mus musculus	Mus musculus	High	NCBI (http://www.ncbi.nlm.nih.gov/Taxonomy/Browser/wwwtax.cgi?mode=Info&id=10090)
Oryctolagus cuniculus	Oryctolagus cuniculus	Moderate	NCBI (http://www.ncbi.nlm.nih.gov/Taxonomy/Browser/wwwtax.cgi?mode=Info&id=9986)

Term	Scientific Term	Evidence	Links
Brassica napus	Brassica napus	Moderate	NCBI (http://www.ncbi.nlm.nih.gov/Taxonomy/Browser/wwwtax.cgi?mode=Info&id=3708)

Life Stage Applicability

Life Stage	Evidence
All life stages	Moderate

Sex Applicability

Sex	Evidence
Unspecific	High

The relationship between HDAC inhibition and hyperacetylation is likely well conserved between species from lower organisms to mammals.

- Hyperacetylation by HDIs such as SAHA and Cpd-60 are observed in mouse (*Mus musculus*) [Schroeder et al., 2013].
- TSA induces acetylation of histone H4 in time-dependent manner in mouse cell lines (*Mus musculus*) [Alberts et al., 1998].
- AR-42, a novel HDI, induces the hyperacetylation in human pancreatic cancer cells (*Homo sapiens*) [Henderson et al., 2016].
- SAHA and MS-275 leads to the hyperacetylation of lysine residues of histones in human cell lines of epithelial (A549) and lymphoid origin (Jurkat) (*Homo sapiens*) [Choudhary et al., 2009].
- SAHA treatment induces the H3 and H4 histone acetylation in human corneal fibroblasts and conjunctiva from rabbits after glaucoma filtration surgery (*Homo sapiens, Oryctolagus cuniculus*) [Sharma et al., 2016].
- TSA induces the acetylation of histones H3 and H4 in *Brassica napus* microspore cultures (*Brassica napu*) [Li et al., 2014].

Key Event Relationship Description

The HDAC inhibitors (HDIs) inhibit deacetylation of the histone, leading to the increase in histone acetylation and gene transcription. HDACs deacetylate acetylated histone in epigenetic regulation [Falkenberg and Johnstone, 2014].

Histone acetylation is one of the major epigenetic mechanisms and belongs to the posttranslational modifications of histones. Histone acetyltransferase is setting the mark, and deacetylase (HDAC) is responsible for removing the acetyl group from specific lysine residues of the histones. It has been shown that the inhibition of HDACs leads to a hyperacetylation of histones and in general to an imbalance of other histone modifications.

Evidence Supporting this KER**Biological Plausibility**

HDACs are important proteins in epigenetic regulation of gene transcription. Upon the inhibition of HDAC by HDIs, the acetylation of lysine in histone remains and it leads to transcriptional activation or repression, changes in DNA replication and DNA damage repair. The treatment by HDIs increased histone acetylation [Wade et al., 2008].

In all eukaryotes the DNA containing the genetic information of an organism, is organized in chromatin. The basic unit of chromatin is the nucleosome around which the naked DNA is wrapped. A nucleosome consists of two copies of each of the core histones H2A, H2B, H3 and H4 [Luger et al., 1997]. In order to dynamically regulate this highly complex structure several mechanism are involved, including the posttranslational modifications of histones (reviewed in [Bannister and Kouzarides, 2011; Kouzarides, 2007]). For long time it is known that histones get acetylated and that this reaction is catalyzed by histone acetyl transferases (HAT) and the acetyl groups are removed by histone deacetylases (HDAC). Inhibition of HDACs by small molecule compounds lead to hyperacetylation of the histones as the homeostasis of acetylation and deacetylation is disturbed (reviewed in [Gallinari et al., 2007]).

Empirical Evidence

The major empirical evidence came from the incubation of cell culture cells with small molecule compounds that inhibit HDAC enzymes followed by western blots or acid urea gel analysis. The first evidence was shown by Riggs et al. who showed that incubation of HeLa cells with n-butyrate leads to an accumulation of acetylated histone proteins [Riggs et al., 1977]. Later, it was shown that n-butyrate specifically increases the acetylation of histone by the incorporation of radioactive [H^3] acetate and analysis of the histones on acid urea gels that allow the detection of acetylated histones [Cousens et al., 1979]. TSA was shown to be an HDAC inhibitor by acid urea gel analysis in 1990 [Yoshida et al., 1990] and good evidence for VPA as an HDAC inhibitor *in vitro* and *in vivo* was shown using acetyl-specific antibodies and western blot [Gottlicher et al., 2001].

There exist several evidences showing the link between histone deacetylase inhibition and increase in histone acetylation as follows:

- Exposure of mouse embryos *in utero* to VPA and TSA (two well-known HDAC inhibitors) showed an increased histone acetylation level in whole embryo extracts and was also shown *in situ* immuno stainings [Menegola et al., 2005].
- HDAC inhibition by HDIs leads to hyperacetylation of histone and a large number of cellular proteins such as NF- κ B [Falkenberg and Johnstone, 2014; Chen et al., 2018].
- The concentrations of half-maximum inhibitory effect (IC_{50}) for HDAC enzyme inhibition of 20 valproic acid derivatives correlated with teratogenic potential of the compounds, and HDAC inhibitory compounds showed hyperacetylation of core histone 4 in treated F9 cells [Eikel et al., 2006].
- HDIs increase histone acetylation in brain [Schroeder et al., 2013].

- The HDI selectivity exists, in which more acetylation sites on the histones H3 and H4 are responsive to SAHA than MS-275 [Choudhary et al., 2009].
- HDI AR-42 induces acetylation of histone H3 in dose-response manner in human pancreatic cancer cell lines [Henderson et al., 2016].
- MAA treatment in rats (650 mg/kg, for 4, 8, 12, and 24 hrs) induced hyperacetylation in histones H3 and H4 of testis nuclei [Wade et al., 2008].
- HDAC inhibition induced by valproic acid (VPA) leads to histone hyperacetylation in mouse teratocarcinoma cell line F9 [Eikel et al., 2006].
- Hyperacetylation of histone H3 was observed in HDAC1-deficient ES cells [Lagger et al., 2002].
- The treatment of MAA induced histone acetylation in H3K9Ac and H4K12Ac, as well as p53K379Ac [Dayan and Hales, 2014].

Uncertainties and Inconsistencies

HDACs affect a large number of cellular proteins including histones, which reminds us the HDAC inhibition by HDIs hyperacetylates cellular proteins other than histones and exhibit biological effects. It is also noted that HDAC functions as the catalytic subunits of large protein complex, which suggests that the inhibition of HDAC by HDIs affect the function of the large multiprotein complexes of HDAC [Falkenberg and Johnstone, 2014]. Related-analysis are usually indirect or in purified systems, therefore a direct cause-consequence relation is difficult to obtain.

Quantitative Understanding of the Linkage

To quantify acetylation by HDAC, stable isotope labeling with amino acids in cell culture (SILAC) method is used [Choudhary et al., 2009].

Response-response relationship

SAHA and MS-275 treatment leads to increase in acetylation of specific lysine residues on histones more than two-fold [Choudhary et al., 2009]. Acetylation of the variant histone H2AZ-a mark for DNA damage sites- was upregulated almost 20-fold by SAHA, whereas a number of sites on the core histones H3 and H4 are several times more highly regulated in response to SAHA than by MS-275 [Choudhary et al., 2009].

TSA (100 ng/ml) treatment leads to accumulation of the acetylated histones in a variety of mammalian cell lines, and the partially purified HDAC from wild-type FM3A cells was effectively inhibited by TSA ($K_i = 3.4$ nM) [Yoshida et al., 1990].

References

Alberts, A.S. et al. (1998), "Activation of SRF-regulated chromosomal templates by Rho-family GTPases requires a signal that also induces H4 hyperacetylation", *Cell* 92:475-487

Bannister, A. J. and Kouzarides, T. (2011), "Regulation of chromatin by histone modifications", *Cell Res* 21:381-395

Chen, S. et al. (2018), "Valproic acid attenuates traumatic spinal cord injury-induced inflammation via STAT1 and NF- κ B pathway dependent of HDAC3", *J Neuroinflammation* 15:150

Choudhary, C. et al. (2009), "Lysine acetylation targets protein complexes and co-regulates major cellular functions", *Science* 325:834-840

Cousens, L. S. et al. (1979), "Different accessibilities in chromatin to histone acetylase", *J Biol Chem* 254:1716-1723

Dayan, C. and Hales, B.F. (2014), "Effects of ethylene glycol monomethyl ether and its metabolite, 2-methoxyacetic acid, on organogenesis stage mouse limbs in vitro", *Birth Defects Res (Part B)* 101:254-261

Eikel, D. et al. (2006), "Teratogenic effects mediated by inhibition of histone deacetylases: evidence from quantitative structure activity relationships of 20 valproic acid derivatives", *Chem Res Toxicol* 19:272-278

Falkenberg, K.J. and Johnstone, R.W. (2014), "Histone deacetylases and their inhibitors in cancer, neurological disease and immune disorders", *Nat Rev Drug Discov* 13:673-691

Gallinari, P. et al. (2007), "HDACs, histone deacetylation and gene transcription: From molecular biology to cancer therapeutics", *Cell Res* 17:195-211

Gottlicher, M. et al. (2001), "Valproic acid defines a novel class of HDAC inhibitors inducing differentiation of transformed cells", *EMBO J* 20:6969-6978

Henderson, S.E. et al. (2016), "Suppression of tumor growth and muscle wasting in a transgenic mouse model of pancreatic cancer by the novel histone deacetylase inhibitor AR-42", *Neoplasia* 18:765-774

Kouzarides, T. (2007), "Chromatin modifications and their function", *Cell* 128:693-705

Lagger, G. et al. (2002), "Essential function of histone deacetylase 1 in proliferation control and CDK inhibitor repression", *EMBO J* 21:2672-2681

Li, H. et al. (2014), "The histone deacetylase inhibitor trichostatin A promotes totipotency in the male gametophyte", *Plant Cell* 26:195-209

Luger, K. et al. (1997), "Crystal structure of the nucleosome core particle at 2.8 a resolution", *Nature* 389:251-260

Menegola, E. et al. (2005), "Inhibition of histone deacetylase activity on specific embryonic tissues as a new mechanism for teratogenicity", *Birth Defects Res B Dev Reprod Toxicol* 74:392-398

Riggs, M.G. et al. (1977), "N-butyrate causes histone modification in HeLa and friend erythroleukaemia cells", *Nature* 268:462-464

Schroeder, F.A. et al. (2013), "A selective HDAC 1/2 inhibitor modulates chromatin and gene expression in brain and alters mouse behavior in two mood-related tests", *PLoS One* 8:e71323

Sharma, A. et al. (2016), "Epigenetic modification prevents excessive wound healing and scar formation after glaucoma filtration surgery", *Invest Ophthalmol Vis Sci* 57:3381-3389

Wade, M.G. et al. (2008), "Methoxyacetic acid-induced spermatocyte death is associated with histone hyperacetylation in rats", *Biol Reprod* 78:822-831

Yoshida, M. et al. (1990), "Potent and specific inhibition of mammalian histone deacetylase both in vivo and in vitro by trichostatin A", J Biol Chem 265:17174-17179

Relationship: 1997: Histone acetylation, increase leads to Cell cycle, disrupted (<https://aopwiki.org/relationships/1997>)
AOPs Referencing Relationship

AOP Name	Adjacency	Weight of Evidence	Quantitative Understanding
Histone deacetylase inhibition leading to testicular atrophy (https://aopwiki.org/aops/212)	adjacent	Moderate	Moderate

Evidence Supporting Applicability of this Relationship

Taxonomic Applicability

Term	Scientific Term	Evidence	Links
Homo sapiens	Homo sapiens	High	NCBI (http://www.ncbi.nlm.nih.gov/Taxonomy/Browser/wwwtax.cgi?mode=Info&id=9606)
Mus musculus	Mus musculus	High	NCBI (http://www.ncbi.nlm.nih.gov/Taxonomy/Browser/wwwtax.cgi?mode=Info&id=10090)

Life Stage Applicability

Life Stage	Evidence
All life stages	High

Sex Applicability

Sex	Evidence
Unspecific	High

The relationship between increased histone acetylation and p21 expression increase is likely well conserved between species.

- Chidamide induced histone acetylation and cell cycle arrest in RPMI8226 and U266 human myeloma cells (*Homo sapiens*) [Yuan et al., 2019].
- TSA and sodium butyrate induced cell cycle regulator p21 mRNA expression in HT-29 human colon carcinoma cells (*Homo sapiens*) [Wu et al., 2001].
- VPA increased acetylation of histone H3 from 3 hrs to 72 hrs after the treatment, and increased p21 expression in 24 hrs after the treatment in K562 cells (*Homo sapiens*) [Gurvich et al., 2004].
- Scriptaid, a HDI, up-regulated p21 mRNA expression in mouse embryonic kidney cells (*Mus musculus*) [Chen et al., 2011].

Key Event Relationship Description

Upon histone acetylation increase, cell cycle regulation is disrupted. Acetylation of the promoter region of the coding genes have a close correlation [Gurvich et al., 2004]. Transient histone hyperacetylation was sufficient for the activation of down-stream molecules involving cell cycle regulation [Wu et al., 2001]. Histone hyperacetylating agents butyrate and TSA induced mRNA expression of cell cycle regulator gene [Archer et al., 1998]. SAHA induced the accumulation of acetylated histones in the chromatin of the gene regulating cell cycle [Richon et al., 2000].

Evidence Supporting this KER

Biological Plausibility

Histone deacetylase inhibitors induce histone hyperacetylation and the activation of down-stream molecules leading to the cell cycle arrest, which suggests the close correlation between histone hyperacetylation and cell cycle arrest [Yuan et al., 2019]. The histone acetylation regulates the gene transcription through the promoter region of the coding gene, which may lead to the overexpression of cell cycle regulators [Richon et al., 2000; Struhal, 1998]. Histone deacetylase inhibition leads to acetylation of histone, inducing the expression of cyclin-dependent kinase inhibitors, followed by a cell-cycle arrest [Li and Seto, 2016].

Empirical Evidence

- MAA induced histone acetylation of H4 in prostate cancer cells including LNCaP, C4-2B, PC-3 and DU-145 parallel with cyclin dependent kinase inhibitor p21, a cell cycle regulator, mRNA level increase [Parajuli et al., 2014].
- HDIs accumulated acetylation of histones and induced cell cycle regulator p21 protein and mRNA expression [Richon et al., 2000; Wu et

al., 2001].

Uncertainties and Inconsistencies

The histone acetylation causes cell cycle disruption in several pathways, in which the specific molecule involvement remains uncertain.

Quantitative Understanding of the Linkage

Histone acetylation occurs in dose-dependent manner with the treatment of chidamide for 48 hrs [Yuan et al., 2019]. The expression of proteins related to G₀/G₁ cell cycle arrest, p21 and phosphorylated p53 is increased in dose-dependent manner [Yuan et al., 2019].

Response-response relationship

Dose-response of histone acetylation and expression of p21 and phosphorylated p53 showed that treatment with 0.5, 1, or 2 micro mol/l of chidamide for 48hrs induced histone acetylation in RPMI8226 myeloma cells, while 2, 4, or 8 micro mol/l of chidamide for 48 hrs induced histone acetylation in U266 myeloma cells [Yuan et al., 2019]. Chidamide treatment in 0.5, 1, or 2 micro mol/l in RPMI8226 or 2, 4, or 8 micro mol/l in U266 induced G₀/G₁ arrest in the myeloma cells [Yuan et al., 2019]. Dose-response of valproic acid (VPA) showed that 5, 10, and 20 mM of VPA inhibited HDAC6 and HDAC7 activity in 293T cells, and 0.1-2 mM of VPA induced acetylation of lysine in H3 in U937 cells [Gurvich et al., 2004]. The p21 protein level was induced with the treatment of 0.25-2 mM of VPA in U937 cells [Gurvich et al., 2004].

Time-scale

Time course for histone H4 hyperacetylation in response to repeated doses of TSA every 8 hrs showed that histone hyperacetylation was peaked in 12 hrs in 8-fold increase and showed 5-fold increase in 24 hrs compared to control [Wu et al., 2001]. TSA (0.3 uM) induced cell cycle regulator p21 mRNA expression in 1 hr after stimulation and the induction is returned to the basal level in 24 hrs [Wu et al., 2001]. Sodium butyrate (5 mM) and repetitive doses of TSA (0.3 uM, every 8 hrs) induced the p21 mRNA level in 24 hrs in HT-29 cells [Wu et al., 2001]. Acetylation of p21 promoter and p21 mRNA induction were correlated in treatment of valproic acid and analogs [Gurvich et al., 2004]. MAA-induced acetylation increases in histones H3 and H4 was occurred in 4, 8, 12 hrs and returned to basal level in 24 hrs after the treatment in rat testis [Wade et al., 2008].

References

Archer, S.Y. et al. (1998), "p21WAF1 is required for butyrate-mediated growth inhibition of human colon cancer cells", Proc Natl Acad Sci USA 95:6791-6796

Chen, S. et al. (2011), "Histone deacetylase (HDAC) activity for embryonic kidney gene expression, growth, and differentiation", J Biol Chem 286:32775-32789

Gurvich, N. et al. (2004), "Histone deacetylase is a target of valproic acid-mediated cellular differentiation", Cancer Res 64:1079-1086

Li, Y. and Seto, E. (2016), "HDACs and HDAC inhibitors in cancer development and therapy", Cold Spring Harb Perspect Med 6:a026831

Parajuli, K.R. et al. (2014), "Methoxyacetic acid suppresses prostate cancer cell growth by inducing growth arrest and apoptosis", Am J Clin Exp Urol 2:300-313

Richon, V.M. et al. (2000), "Histone deacetylase inhibitor selectively induces p21WAF1 expression and gene-associated histone acetylation", Proc Natl Acad Sci 97:10014-10019

Struhl, K. (1998), "Histone acetylation and transcriptional regulatory mechanisms", Gene Dev 12:599-606

Wade, M.G. et al. (2008), "Methoxyacetic acid-induced spermatocyte death is associated with histone hyperacetylation in rats", Biol Reprod 78:822-831

Wu, J.T. et al. (2001), "Transient vs prolonged histone hyper acetylation: effects on colon cancer cell growth, differentiation, and apoptosis", Am J Physiol Gastrointest Liver Physiol 280:G482-G490

Yuan, X. et al. (2019), "Chidamide, a histone deacetylase inhibitor, induces growth arrest and apoptosis in multiple myeloma cells in a caspase-dependent manner", Oncol Let 18:411-419

Relationship: 1712: Cell cycle, disrupted leads to Apoptosis (<https://aopwiki.org/relationships/1712>)

AOPs Referencing Relationship

AOP Name	Adjacency	Weight of Evidence	Quantitative Understanding
Histone deacetylase inhibition leading to testicular atrophy (https://aopwiki.org/aops/212)	adjacent	Moderate	Moderate

Evidence Supporting Applicability of this Relationship

Taxonomic Applicability

Term	Scientific Term	Evidence	Links

Term	Scientific Term	Evidence	Links
Homo sapiens	Homo sapiens	High	NCBI (http://www.ncbi.nlm.nih.gov/Taxonomy/Browser/wwwtax.cgi?mode=Info&id=9606)
Mus musculus	Mus musculus	High	NCBI (http://www.ncbi.nlm.nih.gov/Taxonomy/Browser/wwwtax.cgi?mode=Info&id=10090)
Oryctolagus cuniculus	Oryctolagus cuniculus	Moderate	NCBI (http://www.ncbi.nlm.nih.gov/Taxonomy/Browser/wwwtax.cgi?mode=Info&id=9986)

Life Stage Applicability

Life Stage	Evidence
Not Otherwise Specified	High

Sex Applicability

Sex	Evidence
Unspecific	High

The relationship between disrupted cell cycle and apoptosis is likely well conserved between species.

- MicroRNA let-7a induced cell cycle arrest, inhibited CCND2 and proliferation of human prostate cancer cells (*Homo sapiens*) [Dong et al., 2010].
- microRNA-497 down-regulated CCND2 and induced apoptosis via the Bcl-2/Bax-caspase 9- caspase 3 pathway in HUVECs (*Homo sapiens*) [Wu et al., 2016].
- microRNA-26a regulated p53-mediated apoptosis and CCND2 and CCNE2 in mouse hepatocyte (*Mus musculus*) [Zhou et al., 2016].

Key Event Relationship Description

Cell cycle dysregulation may leads to apoptosis. Cell cycles characterized by the DNA content changes regulate cell death and cell proliferation [Lynch et al., 1986].

Evidence Supporting this KER

microRNA-497, potentially targeting Bcl2 and Cyclin D2 (CCND2), induced apoptosis via the Bcl-2/Bax - caspase 9 - caspase 3 pathway and CCND2 protein in human umbilical vein endothelial cells (HUVECs) [Wu, 2016]. The microRNA-497 activated caspases 9 and 3, and decreased Bcl2 and CCND2 [Wu et al., 2016]. CCND2 is an important cell cycle gene that induces G₁ arrest [Li et al., 2012], and deregulated CCND2 is implicated in cell proliferation inhibition [Wu et al., 2016; Mermelstein et al., 2005; Dong et al., 2010].

Biological Plausibility

The incidence of apoptosis was increased in vincristine-treated cells, in which metaphases were arrested, compared to untreated cells, which indicates that cell cycle dysregulation leads to apoptosis [Sarraf and Bowen, 1986]. Cell gain and loss are balanced with mitosis and apoptosis [Cree et al., 1987]. Apoptosis is mediated by caspase activation [Porter and Janicke, 1999]. Caspase-3 is activated in the programmed cell death, and the pathways to caspase-3 activation include caspase-9 and mitochondrial cytochrome c release [Porter and Janicke, 1999]. The activation of caspase-3 leads to apoptotic chromatin condensation and DNA fragmentation [Porter and Janicke, 1999]. Sinularin, a marine natural compound, exhibited DNA damage and induced G₂/M cell cycle arrest, followed by apoptosis in human hepatocellular carcinoma HepG2 cells [Chung et al., 2017]. Sinularin induced caspases 8, 9, and 3, and pro-apoptotic protein Bax, whereas it decrease the anti-apoptotic Bcl-2 protein expression level [Chung et al., 2017].

Empirical Evidence

- Cell cycle arrest such as G₁ arrest and G₁/S arrest are observed in apoptosis [Li et al., 2012; Dong et al., 2010].
- microRNA-1 and microRNA-206 represses CCND2, while microRNA-29 represses CCND2 and induces G₁ arrest and apoptosis in rhabdomyosarcoma [Li et al., 2012].
- The blockade of G₁/S transition of cell cycle and reduction of CDK4 and CDK2, and apoptosis have occurred in HDAC inhibitor treatment [Parajuli et al., 2014].

Uncertainties and Inconsistencies

MAA induces CDK4 and CDK2 decreases, cell cycle arrest and apoptosis, which may be regulated by several pathways [Parajuli et al., 2014].

Quantitative Understanding of the Linkage

Cell proliferation which was determined at daily intervals after a 24-hr pulse of [³H]thymidine changed as the amount of DNA in the cultures changed. Cell death which was measured by lactic dehydrogenase (LDH) activity in the medium changed in parallel with the changes in cell proliferation [Lynch et al., 1986]. The decrease in total DNA was measured, the increase in cell death was observed [Lynch et al., 1986].

Response-response relationship

Treatment with sinularin, a natural product isolated from cultured soft coral possessing antineoplastic activity, at 12.5, 25, 50 microM resulted in

AOP212

cell cycle disruption and apoptosis in dose-dependent manner in hepatocellular carcinoma cells [Chun et al., 2017]. The cell cycle disruption and apoptosis are induced by 30 micromol/L curcumin, a major component extracted from turmeric plants which have anti-cancer effect [Liu et al., 2018].

Time-scale

MAA (5 mM) decreases CDK4, CDK2 expression in 48 hrs after the treatment, which indicates the G₁ arrest [Parajuli et al., 2014]. MAA (5 mM) decreases the protein expression of procaspase 7 and 3 in 24 to 72 hrs after the treatment, indicating the activation of caspases 7 and 3 [Parajuli et al., 2014].

References

Chung, T.W. et al. (2017), "Sinularin induces DNA damage, G2/M phase arrest, and apoptosis in human hepatocellular carcinoma cells", *BMC Complement Altern Med* 17:62

Cree, I.A. et al. (1987), "Cell death in granulomata: the role of apoptosis", *J Clin Pathol* 40:1314-1319

Dong, Q. et al. (2010), "microRNA let-7a inhibits proliferation of human prostate cancer cells in vitro and in vivo by targeting E2F2 and CCND2", *PLoS One* 5:e10147

Kerr, J.F.R. et al. (1972), "Apoptosis: a basic biological phenomenon with wide-ranging implications in tissue kinetics", *Br J Cancer* 26:239-257

Li, L. et al. (2012), "Downregulation of microRNAs miR-1, -206 and -29 stabilizes PAX3 and CCND2 expression in rhabdomyosarcoma", *Lab Invest* 92:571-583

Liu, W. et al. (2018), "Curcumin suppresses gastric cancer biological activity by regulation of miRNA-21: an in vitro study", *Int J Clin Exp Pathol* 11:5820-5289

Lynch, M.P. et al. (1986), "Evidence for soluble factors regulating cell death and cell proliferation in primary cultures of rabbit endometrial cells grown on collagen", *Proc Natl Acad Sci USA* 83:4784-4788

Mermelstein, A. et al. (2005), "Expression of F-type cyclins in colon cancer and in cell lines from colon carcinomas", *Br J Cancer* 93:338-345

Parajuli, K.R. et al. (2014), "Methoxyacetic acid suppresses prostate cancer cell growth by inducing growth arrest and apoptosis", *Am J Clin Exp Urol* 2:300-313

Porter, A.G. and Janicke, R.U. (1999), "Emerging roles of caspase-3 in apoptosis", *Cell Death Differ* 6:99-104

Sarraf, C.E. and Bowen, I.D. (1986), "Kinetic studies on a murine sarcoma and an analysis of apoptosis", *Br J Cancer* 54:989-998

Wu, R. et al. (2016), "microRNA-497 induces apoptosis and suppressed proliferation via the Bcl-2/Bax-caspase9-caspase 3 pathway and cyclin D2 protein in HUVECs", *PLoS One* 11:e0167052

Zhou, J. et al. (2016), "miR-26a regulates mouse hepatocyte proliferation via directly targeting the 3' untranslated region of CCND2 and CCNE2", *Hepatobiliary Pancreat Dis Int* 15:65-72

Relationship: 1735: Apoptosis leads to Spermatocyte depletion (<https://aopwiki.org/relationships/1735>)

AOPs Referencing Relationship

AOP Name	Adjacency	Weight of Evidence	Quantitative Understanding
Histone deacetylase inhibition leading to testicular atrophy (https://aopwiki.org/aops/212)	adjacent	High	Not Specified

Evidence Supporting Applicability of this Relationship

Taxonomic Applicability

Term	Scientific Term	Evidence	Links
Mus musculus	Mus musculus	High	NCBI (http://www.ncbi.nlm.nih.gov/Taxonomy/Browser/wwwtax.cgi?mode=Info&id=10090)
Rattus norvegicus	Rattus norvegicus	High	NCBI (http://www.ncbi.nlm.nih.gov/Taxonomy/Browser/wwwtax.cgi?mode=Info&id=10116)

Life Stage Applicability

Life Stage	Evidence
Adult, reproductively mature	High

Sex Applicability

Sex	Evidence
Male	High

The apoptosis of the cells leads to spermatocyte depletion. The relationship between apoptosis and spermatocyte depletion is likely well conserved between species.

- Spermatogenesis was inhibited by knockdown of Sucl2, a β subunit of succinyl coenzyme A synthase, via apoptosis in the mouse spermatocyte (*Mus musculus*) [Huang et al., 2016].
- The suppression of microRNA-21 led to apoptosis of spermatogonial stem cell-enriched germ cell cultures and the decrease in the number of spermatogonial stem cells in mice (*Mus musculus*) [Niu et al., 2011].
- MAA induced apoptosis and depletion of spermatocytes in adult rats (*Rattus norvegicus*) [Brinkworth et al., 1995].
- The apoptosis and proliferation inhibition induced by MAA, a HDAC inhibitor, was measured in human prostate cancer cell lines (*Homo sapiens*) [Parajuli et al., 2014].
- The cell viability inhibition induced by SAHA or TSA, which are HDAC inhibitors, was observed in NHDFs (*Homo sapiens*) [Glaser et al., 2003].
- The proliferation of the HDAC^{-/-} ES cells was inhibited compared to HDAC^{+/+} ES cells (*Homo sapiens*) [Zupkovitz et al., 2010].
- It has been reported that mice lacking both *Ink4c* and *Ink4d*, cyclin D-dependent kinase inhibitors, produced few mature sperm, and the residual spermatozoa had reduced motility and decreased viability (*Mus musculus*) [Zindy et al., 2001].
- The sperm counts in the cauda epididymis of rats exposed to butylparaben were significantly decreased (*Rattus norvegicus*) [Oishi, 2001].
- MAA treatment induced spermatocyte death in Sprague-Dawley rats (*Rattus norvegicus*) [Wade et al., 2008].

Key Event Relationship Description

Apoptosis results in spermatocyte depletion via cell death. Apoptosis and spermatocyte depletion is correlated, where spermatocyte depletion via apoptosis is a general mechanism [Brinkworth et al., 1995].

Evidence Supporting this KER

Biological Plausibility

Induced apoptosis during development of germ cells results in progressive depletion of spermatocyte [Brinkworth et al., 1995]. A HDAC inhibitor, MAA, induced apoptosis and spermatocyte depletion at stages IX-II [Brinkworth et al., 1995].

Empirical Evidence

In the mouse spermatocyte, spermatogenesis is inhibited by knockdown of Sucl2, a beta subunit of succinyl coenzyme A synthase, which is located in mitochondria and catalyzes the reversible synthesis of succinate and adenosine triphosphate in the tricarboxylic acid cycle [Huang et al., 2016]. The knockdown of Sucl2 induces apoptosis of mouse spermatocytes [Huang et al., 2016]. The prolonged cryptorchidism leads to germ cell apoptosis and testicular sperm count decrease [Barqawi et al., 2004]. CD147 was reported to regulate apoptosis in mouse testis and spermatocyte cell line (GC-2 cells) via NF κ B pathway [Wang et al., 2017]. MicroRNA-21 regulates the spermatogonial stem cell homeostasis, in which suppression of microRNA-21 with anti-miR-21 oligonucleotides led to apoptosis of spermatogonial stem cell-enriched germ cell cultures and the decrease in the number of spermatogonial stem cells [Niu et al., 2011].

Uncertainties and Inconsistencies

The process of apoptosis is necessary for the meiosis of the stem cell differentiation in the testis, which remains in question for the regulation of spermatocyte deletion and testis atrophy/weight loss [Dym, 1994].

References

Barqawi, A. et al. (2004), "Effect of prolonged cryptorchidism on germ cell apoptosis and testicular sperm count", Asian J Androl 6:47-51

Bose, R. et al. (2017), "Ubiquitin ligase Huwe1 modulates spermatogenesis by regulating spermatogonial differentiation and entry into meiosis", Sci Rep 7:17759

Brinkworth, M. et al. (1995), "Identification of male germ cells undergoing apoptosis in adult rats", J Reprod Fertil 105:25-33

Dym, M. (1994), "Spermatogonial stem cells of the testis", Proc Natl Acad Sci USA 91:11287-11289

Glaser, K.B. et al. (2003), "Gene expression profiling of multiple histone deacetylase (HDAC) inhibitors: defining a common gene set produced by HDAC inhibition in T24 and MDA carcinoma cell lines", Mol Cancer Ther 2:151-163

Huang, S. et al. (2016), "Knockdown of Sucl2 decreases the viability of mouse spermatocytes by inducing apoptosis through injury of the mitochondrial function of cells", Folia Histochem Cytopiol 54:134-142

Niu, Z. et al. (2011), "microRNA-21 regulates the self-renewal of mouse spermatogonial stem cells", Proc Natl Acad Sci 108:12740-12745

Oishi, S. (2001), "Effects of butylparaben on the male reproductive system in rats", Toxicol Indust Health 17:31-39

Parajuli, K.R. et al. (2014), "Methoxyacetic acid suppresses prostate cancer cell growth by inducing growth arrest and apoptosis", Am J Clin Exp Urol 2:300-313

Wade, M.G. et al. (2008), "Methoxyacetic acid-induced spermatocyte death is associated with histone hyperacetylation in rats", Biol Reprod 78:822-831

Wang, C. et al. (2017), "CD147 regulates extrinsic apoptosis in spermatocytes by modulating NF κ B signaling pathways", Oncotarget 8:3132-3143

AOP212

Zindy, F. et al. (2001), "Control of spermatogenesis in mice by the cyclin D-dependent kinase inhibitors p18Ink4c and p19Ink4d", *Mol Cell Biol* 21:3244-3255

Zupkovitz, G. et al. (2010), "The cyclin-dependent kinase inhibitor p21 is a crucial target for histone deacetylase 1 as a regulator of cellular proliferation", *Mol Cell Biol* 30:1171-1181

Relationship: 1734: Spermatocyte depletion leads to Testicular atrophy (<https://aopwiki.org/relationships/1734>)

AOPs Referencing Relationship

AOP Name	Adjacency	Weight of Evidence	Quantitative Understanding
Histone deacetylase inhibition leading to testicular atrophy (https://aopwiki.org/aops/212)	adjacent	High	Not Specified

Evidence Supporting Applicability of this Relationship

Taxonomic Applicability

Term	Scientific Term	Evidence	Links
Mus musculus	Mus musculus	High	NCBI (http://www.ncbi.nlm.nih.gov/Taxonomy/Browser/wwwtax.cgi?mode=Info&id=10090)
Rattus norvegicus	Rattus norvegicus	High	NCBI (http://www.ncbi.nlm.nih.gov/Taxonomy/Browser/wwwtax.cgi?mode=Info&id=10116)

Life Stage Applicability

Life Stage	Evidence
Adult, reproductively mature	High

Sex Applicability

Sex	Evidence
Male	High

The relationship between spermatocyte depletion and testicular toxicity is likely well conserved between species.

- ME and MAA induced spermatocyte depletion and testicular atrophy in rat (*Rattus norvegicus*) [Beattie et al., 1984].
- Ethylene glycol monomethyl ether induced depletion of late spermatocytes and testicular atrophy in F344 rat (*Rattus norvegicus*) [Chapin et al., 1984].
- The epididymal tubules of rats with testicular degeneration had low sperm density (*Rattus norvegicus*) [Lee et al., 1993].
- Hydroxyurea induced spermatocyte reduction and testicular atrophy (*Mus musculus*) [Wiger et al., 1995].

Key Event Relationship Description

Spermatocyte depletion leads to testicular toxicity such as testicular atrophy with decrease in size. The spermatocyte depletion is involved in testicular atrophy and testicular toxicity [Chapin et al., 1984]. There are different insults that can induce spermatocyte depletion and consequently testicular toxicity.

Evidence Supporting this KER

Biological Plausibility

Spermatocyte depletion caused by apoptosis leads to the testicular toxicity. Apoptosis is a basic biological phenomenon in which the cells are controlled in the atrophy of various tissues and organs through the deletion and turnover, as well as in tumor regression [Kerr et al., 1972].

Empirical Evidence

2-methoxyethanol (ME) or its major metabolite, methoxyacetic acid (MAA), HDAC inhibitor, induced spermatocyte depletion and testicular atrophy [Beattie et al., 1984].

Uncertainties and Inconsistencies

Spermatogonial stem cell self-renewal and spermatocyte meiosis are regulated by Sertoli cell signaling, which suggests us that various pathways in spermatocytes or spermatogonia are involved in the spermatocyte deletion and testis atrophy/weight loss [Chen et al., 2015].

References

Abedi, N. et al. (2017), "Short and long term effects of different doses of paracetamol on sperm parameters and DNA integrity in mice", Middle East Fertility Society Journal 22:323-328

Beattie, P.J. et al. (1984), "The effect of 2-methoxyethanol and methoxyacetic acid on Sertoli cell lactate production and protein synthesis in vitro", Toxicol Appl Pharmacol 76:56-61

Chapin, R.E. et al. (1984), "The effects of ethylene glycol monomethyl ether on testicular histology in F344 rats", J Andro 5:369-380

Chen, S. and Liu, Y. (2015), "Regulation of spermatogonial stem cell self-renewal and spermatocyte meiosis by Sertoli cell signaling", Reproduction 149:R159-R167

de Rooij, D.G. et al. (2001), "Proliferation and differentiation of spermatogonial stem cells", Reproduction 121:347-354

de Rooij, D.G. (1998), "Stem cells in the testis", Int J Exp Path 79:67-80

Kerr, J.F.R. et al. (1972), "Apoptosis: a basic biological phenomenon with wide-ranging implications in tissue kinetics", Br J Cancer 26:239-257

Lee, K.P. et al. (1993), "Testicular degeneration and spermatid retention in young male rats", Toxicol Pathol 21:292-302

Wiger, R. et al. (1995), "Effects of acetaminophen and hydroxyurea on spermatogenesis and sperm chromatin structure in laboratory mice", Reprod Toxicol 9:21-33

List of Non Adjacent Key Event Relationships

Relationship: 1715: Histone deacetylase inhibition leads to Cell cycle, disrupted (<https://aopwiki.org/relationships/1715>)
AOPs Referencing Relationship

AOP Name	Adjacency	Weight of Evidence	Quantitative Understanding
Histone deacetylase inhibition leading to testicular atrophy (https://aopwiki.org/aops/212)	non-adjacent	High	Moderate

Evidence Supporting Applicability of this Relationship

Taxonomic Applicability

Term	Scientific Term	Evidence	Links
Homo sapiens	Homo sapiens	High	NCBI (http://www.ncbi.nlm.nih.gov/Taxonomy/Browser/wwwtax.cgi?mode=Info&id=9606)
Mus musculus	Mus musculus	Moderate	NCBI (http://www.ncbi.nlm.nih.gov/Taxonomy/Browser/wwwtax.cgi?mode=Info&id=10090)

Life Stage Applicability

Life Stage	Evidence
Not Otherwise Specified	High

Sex Applicability

Sex	Evidence
Unspecific	High

MAA induced G₁ cell cycle arrest in human prostate cancer cells (*Homo sapiens*) [Parajuli et al., 2014].

Apicidin induced G₁ cell cycle arrest in HeLa cells (*Homo sapiens*) [Han et al., 2000].

The change in the amounts of cells in G₁ phase and S phase of cell cycle was detected in mouse HDAC1 knock out fibroblast lines (*Mus musculus*) [Zupkovitz et al., 2010].

Loss of HDAC1 in mouse embryonic stem (ES) cells results in the acetylation of histones H3 and H4, up-regulation of cyclin-dependent kinase inhibitors p21^{WAF1/CIP1} and p27^{KIP1} and inhibition of proliferation (*Mus musculus*) [Lagger et al., 2002].

Key Event Relationship Description

HDAC inhibition leads to cell cycle arrest including G₁/S phase arrest [Falkenberg and Johnstone, 2014]. The HDAC inhibition-induced cell cycle arrest is mediated by transcriptional changes of the CDK inhibitors such as p21 [Falkenberg and Johnstone, 2014].

Evidence Supporting this KER

Biological Plausibility

The knockdown of HDACs may induce antitumor effects such as cell cycle arrest and inhibition of proliferation [Falkenberg and Johnstone, 2014]. In leukemia, an oncogenic fusion protein recruits a variety of proteins including HDACs to repress cell cycle inhibitors, which suggests that HDAC inhibition leads to cell cycle dysregulation [Falkenberg and Johnstone, 2014].

Empirical Evidence

- HDAC inhibition with SAHA, TSA and MS-27-275 induced acetylation of histone H4, up-regulation of cyclin-dependent kinase inhibitor p21, and inhibition of proliferation in human bladder carcinoma cells [Glaser et al., 2003].
- Apicidin [cyclo(N-O-methyl-L-tryptophanyl-L-isoleucinyl-D-pipecolinyl-L-2-amino-8-oxodecanoyl)], a fungal metabolite HDI, inhibits proliferation of tumor cells via p21 induction [Han et al., 2000]. Apicidin induced hyperacetylation of histone H4, up-regulation of p21, and G₀/G₁ cell cycle arrest in HeLa cells [Han et al., 2000].
- Falkenberg and Johnstone (2014) nicely reviewed that HDAC inhibition leads to cell cycle arrest in which G₁/S phase arrest occurs via up-regulation of p21.
- Loss of HDAC1 in mouse embryonic stem (ES) cells has demonstrated the acetylation of histones H3 and H4, up-regulation of cyclin-dependent kinase inhibitors p21^{WAF1/CIP1} and p27^{KIP1} and inhibition of proliferation [Lagger et al., 2002].
- G₁/S transition blockade was observed in methoxyacetic acid (MAA)-treated prostate cancer cells [Parajuli et al., 2014].
- The change in the amounts of cells in G₁ phase and S phase of cell cycle was detected in mouse HDAC1 knock out fibroblast lines [Zupkovitz et al., 2010].
- MAA, a HDI, induced cell cycle arrest and up-regulation of p21 expression, and inhibited prostate cancer cell growth [Parajuli et al., 2014].

Uncertainties and Inconsistencies

The involvement of p53/p63/p73 in up-regulation of p21 induced by HDAC inhibition is not fully elucidated, where time course of the p21 and p53/p63/p73 mRNA expression has demonstrated the cell-line specific differences in the responses in 4 human prostate cancer cell lines LNCaP, C4-2B, PC-3 and DU-145 [Parajuli et al., 2014].

Quantitative Understanding of the Linkage

MAA (20 mM) induced G₁ cell cycle arrest upon the treatment for 24 hrs in LNCaP, C4-2B, PC-3 and DU-145 human prostate cancer cell lines [Parajuli et al., 2014]. Almost 80% of the cells were arrested in G₁ phase upon stimulation of MAA, whereas approximately 40 to 60 % of the cells were in G₁ phase without MAA treatment [Parajuli et al., 2014].

Time-scale

MAA (5 mM) induced p21 up-regulation in 12 to 72 hrs in LNCaP, C4-2B, PC-3 and DU-145 human prostate cancer cell lines [Parajuli et al., 2014].

References

Falkenberg, K.J. and Johnstone, R.W. (2014), "Histone deacetylases and their inhibitors in cancer, neurological disease and immune disorders", *Nat Rev Drug Discov* 13:673-691

Glaser, K.B. et al. (2003), "Gene expression profiling of multiple histone deacetylase (HDAC) inhibitors: defining a common gene set produced by HDAC inhibition in T24 and MDA carcinoma cell lines", *Mol Cancer Ther* 2:151-163

Han, J.W. et al. (2000), "Apicidin, a histone deacetylase inhibitor, inhibits proliferation of tumor cells via induction of p21^{WAF1/Cip1} and gelsolin", *Cancer Res* 60:6068-6074

Lagger, G. et al. (2002), "Essential function of histone deacetylase 1 in proliferation control and CDK inhibitor repression", *EMBO J* 21:2672-2681

Parajuli, K.R. et al. (2014), "Methoxyacetic acid suppresses prostate cancer cell growth by inducing growth arrest and apoptosis", *Am J Clin Exp Urol* 2:300-312

Zupkovitz, G. et al. (2010), "The cyclin-dependent kinase inhibitor p21 is a crucial target for histone deacetylase 1 as a regulator of cellular proliferation", *Mol Cell Biol* 30:1171-1181

Relationship: 1716: Histone deacetylase inhibition leads to Apoptosis (<https://aopwiki.org/relationships/1716>)

AOPs Referencing Relationship

AOP Name	Adjacency	Weight of Evidence	Quantitative Understanding
Histone deacetylase inhibition leading to testicular atrophy (https://aopwiki.org/aops/212)	non-adjacent	Moderate	Moderate

Evidence Supporting Applicability of this Relationship

Taxonomic Applicability

Term	Scientific Term	Evidence	Links
Homo sapiens	Homo sapiens	High	NCBI (http://www.ncbi.nlm.nih.gov/Taxonomy/Browser/wwwtax.cgi?mode=Info&id=9606)
Mus musculus	Mus musculus	High	NCBI (http://www.ncbi.nlm.nih.gov/Taxonomy/Browser/wwwtax.cgi?mode=Info&id=10090)

Life Stage Applicability

Life Stage	Evidence
Not Otherwise Specified	High

Sex Applicability

Sex	Evidence
Unspecific	High

AR-42 inhibited proliferation of human pancreatic cancer cells (*Homo sapiens*) [Henderson et al., 2016].

MAA induced apoptosis in human prostate cancer cell lines. The apoptosis and proliferation inhibition induced by MAA, a HDAC inhibitor, was measured in human prostate cancer cell lines (*Homo sapiens*) [Parajuli et al., 2014].

SAHA or TSA, which are HDAC inhibitors, reduced cell viability in NHDFs (*Homo sapiens*) [Glaser et al., 2003].

The proliferation of the HDAC^{-/-} ES cells was inhibited compared to HDAC^{+/+} ES cells (*Homo sapiens*) [Zupkovitz et al., 2010].

Key Event Relationship Description

HDAC inhibition leads to cell death through the apoptotic pathways [Falkenberg and Johnstone, 2014]. Intrinsic apoptosis pathway requires BH3-only proteins, and BCL-2 protein overexpression inhibits apoptosis [Falkenberg and Johnstone, 2014]. Administration of methoxyacetic acid (MAA), a HDAC inhibitor, causes apoptosis with DNA ladder in male germ cells [Brinkworth et al., 1995]. MAA induces the apoptosis of spermatocytes at spermatogenic cycle stage IX-II [Brinkworth et al., 1995].

Evidence Supporting this KER

Biological Plausibility

HDAC inhibition in cancer results in apoptosis with the up-regulation of pro-apoptotic B cell lymphoma 2 (BCL-2) family genes and down-regulation of pro-survival BCL-2 genes [Falkenberg, 2014]. The antitumor effect of HDAC inhibition includes cell death and apoptosis [Falkenberg and Johnstone, 2014].

Empirical Evidence

- MAA-induced spermatocyte death is associated with histone acetylation increase [Wade et al., 2008].
- The HDAC inhibition induced apoptosis markers such as BAK overexpression and suppression of phosphorylated AKT [Henderson et al., 2016].
- The administration of MAA can cause apoptosis in the germ cells of adult male rats [Brinkworth et al., 1995].

Uncertainties and Inconsistencies

It is uncertain through which pathway the HDAC inhibition induces apoptosis.

Quantitative Understanding of the Linkage

MAA (5 mM) induced apoptosis in prostate cancer cell lines, LNCaP, C4-2B, PC-3 and DU-145, in which apoptotic nucleosomes were calculated as absorbance at 405 nm – absorbance at 490 nm [Parajuli et al., 2014].

Time-scale

MAA (5 mM) decreased protein expression of BIRC2 and activated caspases 7 and 3 within 72 hrs [Parajuli et al., 2014].

References

Brinkworth, M.H. et al. (1995), "Identification of male germ cells undergoing apoptosis in adult rats", J Reprod Fertil 105:25-33

Falkenberg, K.J. and Johnstone, R.W. (2014), "Histone deacetylases and their inhibitors in cancer, neurological disease and immune disorders", Nat Rev Drug Discov 13:673-691

Glaser, K.B. et al. (2003), "Gene expression profiling of multiple histone deacetylase (HDAC) inhibitors: defining a common gene set produced by HDAC inhibition in T24 and MDA carcinoma cell lines", Mol Cancer Ther 2:151-163

Henderson, S.E. et al. (2016), "Suppression of tumor growth and muscle wasting in a transgenic mouse model of pancreatic cancer by the novel histone deacetylase inhibitor AR-42", Neoplasia 18:765-774

Parajuli, K.R. et al. (2014), "Methoxyacetic acid suppresses prostate cancer cell growth by inducing growth arrest and apoptosis", Am J Clin Exp Urol 2:300-312

AOP212

Wade, M.G. et al. (2008), "Methoxyacetic acid-induced spermatocyte death is associated with histone hyperacetylation in rats", *Biol Reprod* 78:822-831

Zupkovitz, G. et al. (2010), "The cyclin-dependent kinase inhibitor p21 is a crucial target for histone deacetylase 1 as a regulator of cellular proliferation", *Mol Cell Biol* 30:1171-1181

Relationship: 2010: Histone deacetylase inhibition leads to Spermatocyte depletion
(<https://aopwiki.org/relationships/2010>)

AOPs Referencing Relationship

AOP Name	Adjacency	Weight of Evidence	Quantitative Understanding
Histone deacetylase inhibition leading to testicular atrophy (https://aopwiki.org/aops/212)	non-adjacent	Moderate	Moderate

Evidence Supporting Applicability of this Relationship

Taxonomic Applicability

Term	Scientific Term	Evidence	Links
rat	<i>Rattus norvegicus</i>	High	NCBI (http://www.ncbi.nlm.nih.gov/Taxonomy/Browser/wwwtax.cgi?mode=Info&id=10116)

Life Stage Applicability

Life Stage	Evidence
Adult, reproductively mature	Moderate

Sex Applicability

Sex	Evidence
Male	High

Histone deacetylase inhibition by histone deacetylase inhibitors caused spermatocyte death in rats. MAA treatment induced spermatocyte death in Sprague-Dawley rats (*Rattus norvegicus*) [Wade et al., 2008].

VPA exposure caused decrease in sperm count in human (*Homo sapiens*) [Yerby and McCoy, 1999; Kose-Ozlece et al., 2015].

Key Event Relationship Description

Histone deacetylase inhibition triggered by histone deacetylase inhibitors such as methoxyacetic acid (MAA) leads to spermatocyte death causing spermatocyte depletion [Wade et al., 2008]. Histone deacetylase inhibition leads to increase in histone acetylation, leading to spermatocyte apoptosis.

Evidence Supporting this KER

MAA administration induces the spermatocyte deaths, which has been revealed by section staining of the germ cell death [Wade et al., 2008].

Biological Plausibility

Histone deacetylase inhibition causes histone acetylation, which increases the gene expression of cell-cycle related protein, followed by spermatocyte apoptosis in testis.

Empirical Evidence

Administration of MAA in rats, a histone deacetylase inhibitor, demonstrated the emergence of TUNEL-positive spermatocytes, which indicates the spermatocyte apoptosis [Wade et al., 2008]. Treatment of valproate (VPA) resulted in a decline in the sperm count [Yerby and McCoy, 1999; Kose-Ozlece et al., 2015].

Quantitative Understanding of the Linkage

The administration of MAA in rats induced spermatocyte depletion which was confirmed with TUNEL-staining of the germ cells [Wade et al., 2008].

Time-scale

TUNEL-positive germ cell were increased after 8, 12, and 24 hrs of MAA exposure (650 mg/kg i.p.) in the rats [Wade et al., 2008]. TUNEL-positive zygotene spermatocytes were emerged in after 12 hrs of MAA exposure in the rats, which was confirmed by the section staining [Wade et al., 2008].

References

Kose-Ozlece, H. et al. (2015), "Alterations in semen parameters in men with epilepsy treated with valproate", *Iran J Neurol* 14:164-167

Wade, M.G. et al. (2008), "Methoxyacetic acid-induced spermatocyte death is associated with histone hyperacetylation in rats", *Biol Reprod* 78:822-831

Yerby, M.S. and McCoy, G.B. (1999), "Male infertility: Possible association with valproate exposure", *Epilepsia* 40:520-521

Relationship: 1717: Histone deacetylase inhibition leads to Testicular atrophy (<https://aopwiki.org/relationships/1717>)
 AOPs Referencing Relationship

AOP Name	Adjacency	Weight of Evidence	Quantitative Understanding
Histone deacetylase inhibition leading to testicular atrophy (https://aopwiki.org/aops/212)	non-adjacent	Moderate	Moderate

Evidence Supporting Applicability of this Relationship

Taxonomic Applicability

Term	Scientific Term	Evidence	Links
Homo sapiens	Homo sapiens	High	NCBI (http://www.ncbi.nlm.nih.gov/Taxonomy/Browser/wwwtax.cgi?mode=Info&id=9606)
Rattus norvegicus	Rattus norvegicus	High	NCBI (http://www.ncbi.nlm.nih.gov/Taxonomy/Browser/wwwtax.cgi?mode=Info&id=10116)

Life Stage Applicability

Life Stage	Evidence
Adult, reproductively mature	High

Sex Applicability

Sex	Evidence
Male	High

MAA induced spermatocyte apoptosis and cell morphology change in human testes (*Homo sapiens*) [Li et al., 1996].

Valproic acid caused the decrease in rat testicular weight (*Rattus norvegicus*) [Kallen, 2004].

Key Event Relationship Description

HDAC inhibition induced testicular toxicity including testis atrophy such as the decrease in size [Miller et al., 1982]. HDAC inhibition in cell culture resulted in the testicular toxicity including germ cell apoptosis and cell morphology change [Li et al., 1996]. Valproic acid, a HDAC inhibitor, caused a reduced testicular weight in the offspring in rats [Kallen, 2004].

Evidence Supporting this KER

Biological Plausibility

The HDAC inhibition induced cell death in spermatocytes in both rat and human seminiferous tubules [Li et al., 1996]. The HDAC inhibitor treatment resulted in degeneration in spermatocytes in rat seminiferous tubules [Li et al., 1996]. The HDAC inhibition induced the germ cell apoptosis in human testicular tissues [Li et al., 1996].

Empirical Evidence

- HDAC inhibitor, methoxyacetic acid (MAA), (300 mg/kg) induced testicular toxicity measured with testis weight loss [Miller et al., 1982].
- MAA induced apoptosis and degeneration in spermatocytes in human testicular tissue and 25-day rat seminiferous tubule cultures [Li et al.,

1996].

- MAA-induced spermatocyte death with an association of histone acetylation increase [Wade et al., 2008].
- MAA-induced apoptosis in male germ cells was modulated by Sertoli cells via P/Q type voltage-operated calcium channels [Barone et al., 2005].
- The *p.o.* administration of ethylene glycol monomethyl (500 mg/kg/day) in rats induced the testis or liver organ weight loss on 2, 4, 7 and 11 days or 24 hrs and 2, 4 and 7 days after treatment, respectively [Foster et al., 1983].
- The investigation of 2-methoxyethanol (2-ME)-induced testicular toxicity has revealed that the conversion of 2-ME to MAA is required in 2-ME-induced testicular toxicity [Moss et al., 1985].

Uncertainties and Inconsistencies

It is reported that HDAC inhibition leads to teratogenic toxicity, whereas the correlation with testicular toxicity and teratogenic toxicity by HDAC inhibition is not fully understood [Menegola et al., 2006]. The oral administration of vorinostat (SAHA), a HDAC inhibitor, in Sprague-Dawley rats showed no indication of reproductive toxicity in drug-treated male rats, which suggested the involvement of some compensation mechanisms or digestion [Wise et al., 2008]. Some studies have demonstrated that the decrease in histone acetylation in spermatids is associated with impaired spermatogenesis corresponding with the well-known reduction of protamine expression in the cells [Sonnack et al., 2002; Li et al., 2014]. It has also been reported that the histological examination of sections revealed no difference between wild-type and HDAC6-deficient testes [Zhang et al., 2008].

Quantitative Understanding of the Linkage

MAA administration (592 mg/kg/day) for 4 days showed testis weight loss in which the relative organ weights were 0.773 ± 0.022 g/100 g body weight, compared to 0.985 ± 0.028 g/100g body weight in control treated with water [Foster et al., 1984].

Time-scale

The relative testicular weight was decreased at day 2 after the treatment of 500 mg/kg/day treatment of ethylene glycol monomethyl ether [Foster et al., 1984]. The treatment of 5 mM MAA for 5 hrs induced the pachytene spermatocyte death in early stage tubules in 19 hrs [Li et al., 1996]. The degeneration in late spermatocytes was observed in late-stage tubules in 19 hrs after 5 mM MAA treatment for 5 hrs [Li et al., 1996].

References

Barone, F. et al. (2005), "Modulation of MAA-induced apoptosis in male germ cells: role of Sertoli cell P/Q-type calcium channels", *Reprod Biol Endocrinol* 3:13

Foster, P.M. et al. (1983), "Testicular toxicity of ethylene glycol monomethyl and monoethyl ethers in the rat", *Toxicol Appl Pharmacol* 69:385-399

Foster, P.M. et al. (1984), "Testicular toxicity produced by ethylene glycol monomethyl and monoethyl esters in the rat", *Environ Health Perspect* 57:207-217

Kallen, B. (2004), "Valproic acid is known to cause hypospadias in man but does not reduce anogenital distance or causes hypospadias in rats", *Basic Clin Pharmacol Toxicol* 94:51-54

Li, L.H. et al. (1996), "2-Methoxyacetic acid (MAA)-induced spermatocyte apoptosis in human and rat testes: an in vitro comparison", *J Androl* 17:538-549

Li, W. et al. (2014), "Chd5 orchestrates chromatin remodeling during sperm development", *Nat Commun* 5:3812

Menegola, E. et al. (2006), "Inhibition of histone deacetylase as a new mechanism of teratogenesis", *Birth Defects Res* 78:345-353

Miller, R.R. et al. (1982), "Toxicity of methoxyacetic acid in rats", *Fundam Appl Toxicol* 2:158-160

Moss, E.J. et al. (1985), "The role of metabolism in 2-methoxyethanol-induced testicular toxicity", *Toxicol Appl Pharmacol* 79:480-489

Sonnack, V. et al. (2002), "Expression of hyperacetylated histone H4 during normal and impaired human spermatogenesis", *Andrologia* 34:384-390

Wade, M.G. et al. (2008), "Methoxyacetic acid-induced spermatocyte death is associated with histone hyperacetylation in rats", *Biol Reprod* 78:822-831

Wise, L.D. et al. (2008), "Assessment of female and male fertility in Sprague-Dawley rats administered vorinostat, a histone deacetylase inhibitor", *Birth Defects Res B Dev Reprod Toxicol* 83:19-26

Zhang, Y. et al. (2008), "Mice lacking histone deacetylase 6 have hyperacetylated tubulin but are viable and develop normally", *Mol Cel Biol* 28:1688-1701