


AOP ID and Title:

AOP 298: Increase in reactive oxygen species (ROS) leading to human treatment-resistant gastric cancer

Short Title: Increase in ROS leading to human treatment-resistant gastric cancer**Graphical Representation****Authors**Shihori Tanabe¹⁾, Sabina Quader²⁾, Ryuichi Ono³⁾, Horacio Cabral⁴⁾, Ed Perkins⁵⁾¹Division of Risk Assessment, Center for Biological Safety and Research, National Institute of Health Sciences, Japan²Innovation Centre of NanoMedicine (iCONM), Kawasaki Institute of Industrial Promotion, Japan³Division of Cellular and Molecular Toxicology, Center for Biological Safety and Research, National Institute of Health Sciences, Japan⁴Department of Bioengineering, Graduate School of Engineering, The University of Tokyo, Japan⁵Environmental Laboratory, US Army Engineer Research and Development Center, United States**Status**

Author status	OECD status	OECD project	SAAOP status
Open for comment. Do not cite	Under Review	1.58	Included in OECD Work Plan

Abstract

This AOP entitled “Increase in reactive oxygen species (ROS) leading to human treatment-resistant gastric cancer” consists of MIE as KE1115 “Increase, ROS” followed by KE1 as KE1753 “Chronic ROS,” KE2 as KE1754 “porcupine-induced Wnt secretion and Wnt signaling activation,” KE3 as KE1755 “beta-catenin activation,” KE4 as KE1457 “epithelial-mesenchymal transition (EMT),” and AO as KE1651 “human treatment-resistant gastric cancer.” ROS has multiple roles in disease, such as the development and progression of cancer or apoptotic induction, causing anti-tumor effects. In this AOP, we focus on sustained levels of chronic reactive oxygen species (ROS) in inducing therapy resistance in human gastric cancer. Epithelial-mesenchymal transition (EMT), a cellular phenotypic change from epithelial to mesenchymal-like features, demonstrates cancer stem cell-like characteristics in human gastric cancer. EMT is induced by Wnt/beta-catenin signaling, providing the rationale to have Wnt secretion and beta-catenin activation as KE1 and KE2 on the AOP, respectively. The AOP might be useful for the development of anti-cancer drugs or the prediction of adverse effects of therapeutics, which are of possible regulatory relevance.

Summary of the AOP**Events****Molecular Initiating Events (MIE), Key Events (KE), Adverse Outcomes (AO)**

Sequence	Type	Event ID	Title	Short name
	MIE	1115	Increase, Reactive oxygen species	Increase, ROS
3	KE	1754	Increase, porcupine-induced Wnt secretion and Wnt signaling activation	Increase, porcupine-induced Wnt secretion and Wnt signaling activation
4	KE	1755	beta-catenin activation	beta-catenin activation
	KE	1457	Epithelial Mesenchymal Transition	EMT
6	AO	1651	Treatment-resistant gastric cancer	Resistant gastric cancer

Key Event Relationships

Upstream Event	Relationship Type	Downstream Event	Evidence	Quantitative Understanding
Increase, Reactive oxygen species	adjacent	Increase, porcupine-induced Wnt secretion and Wnt signaling activation	Moderate	Moderate
Increase, porcupine-induced Wnt secretion and Wnt signaling activation	adjacent	beta-catenin activation	Moderate	Moderate
beta-catenin activation	adjacent	Epithelial Mesenchymal Transition	Moderate	Moderate
Epithelial Mesenchymal Transition	adjacent	Treatment-resistant gastric cancer	Moderate	Moderate

Stressors

Name	Evidence
Wnt	High
WNT2	High
Porcupine	Moderate
Wntless	Moderate
Ionizing Radiation	Moderate
ferric nitrilotriacetate	Not Specified

Wnt

WNT induces EMT (J. Zhang, Tian, & Xing, 2016).

WNT2

WNT2 induces EMT in cervical cancer (Zhou et al., 2016).

Porcupine

Porcupine palmitoleates Wnt and facilitates the secretion of the Wnt ligand([Yu & Virshup, 2014](#)) .

Wntless

Wntless binds to and transport Wnt to the plasma membrane leading to the secretion of Wnt ligand([Yu & Virshup, 2014](#)).

ferric nitrilotriacetate

Carcinogenic iron(III)-nitrilotriacetate induces reactive oxygen species production via transfer of an electron to molecular oxygen to form reactive oxygen species [Tsuchiya K, Akai K, Tokumura A, Abe S, Tamaki T, Takiguchi Y, Fukuzawa K. *Biochim Biophys Acta*. 2005 Aug 30;1725(1):111-9. doi: 10.1016/j.bbagen.2005.05.001, Akai K, Tsuchiya K, Tokumura A, Kogure K, Ueno S, Shibata A, Tamaki T, Fukuzawa K. *Free Radic Res*. 2004 Sep;38(9):951-62. doi: 10.1080/1071576042000261945].

Overall Assessment of the AOP

1. Support for Biological Plausibility of KERs	
MIE => KE1: Increase, ROS leads to porcupine-induced Wnt secretion and Wnt signaling activation	Biological Plausibility of the MIE => KE1 is moderate. Rationale: Sustained ROS caused by/causes DNA damage, which will alter several signaling pathways including Wnt signaling. ROS stimulate inflammatory factor production and Wnt/b-catenin signaling. Macrophages accumulate into injured tissue to recover the tissue damage, which may be followed by porcupine-induced Wnt secretion (Vallée & Lecarpentier, 2018).
KE1 => KE2: Porcupine-induced Wnt secretion and Wnt signaling activation leads to beta-catenin activation	Biological Plausibility of the KE1 => KE2 is moderate. Rationale: Secreted Wnt ligand stimulates Wnt/b-catenin signaling, in which b-catenin is activated. Wnt ligand binds to Frizzled receptor, which leads to GSK3b inactivation. GSK3b inactivation leads to b-catenin dephosphorylation, which avoids the ubiquitination of the b-catenin and stabilize the b-catenin (Clevers & Nusse, 2012).

KE2 => KE3: beta-catenin activation leads to Epithelial-mesenchymal transition (EMT)	Biological Plausibility of the KE2 => KE3 is moderate. Rationale: b-catenin activation, of which mechanism include the stabilization of the dephosphorylated b -catenin and translocation of b-catenin into the nucleus, induce the formation of b-catenin-TCF complex and transcription of transcription factors such as Snail, Zeb and Twist (Clevers & Nusse, 2012) (Ahmad et al., 2012; Pearlman et al., 2017; Sohn et al., 2019; Yang W et al., 2019). EMT-related transcription factors including Snail, ZEB and Twist are up-regulated in cancer cells (Diaz et al., 2014). The transcription factors such as Snail, ZEB and Twist bind to E-cadherin (CDH1) promoter and inhibit the CDH1 transcription via the consensus E-boxes (5'-CACCTG-3' or 5'-CAGGTG-3'), which leads to EMT (Diaz et al., 2014).
KE3 => AO: Epithelial-mesenchymal transition (EMT) leads to treatment-resistant gastric cancer	Biological Plausibility of the KE3 => AO is moderate. Rationale: Some population of the cells exhibiting EMT demonstrates the feature of cancer stem cells (CSCs), which are related to cancer malignancy (Shibue & Weinberg, 2017; Tanabe, 2015a, 2015b; Tanabe et al., 2015). EMT phenomenon is related to cancer metastasis and cancer therapy resistance (Smith & Bhowmick, 2016; Tanabe, 2013). Increase expression of enzymes that degrade the extracellular matrix components and the decrease in adhesion to the basement membrane in EMT induce the cell escape from the basement membrane and metastasis (Smith & Bhowmick, 2016). Morphological changes observed during EMT is associated with therapy resistance (Smith & Bhowmick, 2016).
2. Support for essentiality of KEs	
MIE: Increase, ROS	Essentiality of the MIE is high. Rationale for Essentiality of the MIE in the AOP: Sustained ROS contributes to the initiation and development of human gastric cancer (Gu et al., 2018).
KE1: Porcupine-induced Wnt secretion and Wnt signaling activation	Essentiality of the KE1 is moderate. Rationale for Essentiality of the KE1 in the AOP: The Wnt signaling activation is essential for the subsequent b-catenin activation and cancer resistance (Tanabe, 2018).
KE2: beta-catenin activation	Essentiality of the KE2 is moderate. Rationale for Essentiality of the KE2 in the AOP: b-catenin activation is essential for the Wnt-induced cancer resistance (Tanabe, 2018).
KE3: Epithelial-mesenchymal transition (EMT)	Essentiality of the KE3 is moderate. Rationale for Essentiality of the KE3 in the AOP: EMT is essential for the Wnt-induced cancer promotion and acquisition of resistance to anti-cancer drug (Tanabe, 2018, Tanabe et al, 2020a, 2020b, 2023).

Domain of Applicability

Life Stage Applicability

Life Stage Evidence

All life stages High

Taxonomic Applicability

Term Scientific Term Evidence Links

Homo sapiens Homo sapiens High [NCBI](#)

Sex Applicability

Sex Evidence

Unspecific High

The AOP298 applies to *Homo sapiens* (human), all life stages, and both male and female.

Essentiality of the Key Events

Sustained ROS contributes to the initiation and development of human gastric cancer (Gu et al., 2018).

Wnt signaling is involved in cancer malignancy (Tanabe, 2018).

Upon stimulation with Wnt ligand to the Frizzled receptor, Wnt/beta-catenin signaling is activated. Wnt/beta-catenin consists of GSK3 beta inactivation, beta-catenin activation, and up-regulation of transcription factors such as Zeb, Twist, and Snail. The transcription factors Zeb, Twist and Snail relate to the activation of EMT-related genes. EMT is regulated with various gene networks (Tanabe, 2015c, Tanabe et al, 2020a, 2020b).

Event	Direct Evidence	Indirect Evidence	No experimental evidence
MIE: Increase, ROS	***	*	□
KE1: Increase, porcupine-induced Wnt secretion and Wnt signaling activation	□	*	**
KE2: beta-catenin activation	□	*	**
KE3: Epithelial-Mesenchymal Transition	□	*	**

Weight of Evidence Summary

The Wnt signaling promotes EMT and cancer malignancy in colorectal cancer (Lazarova & Bordonaro, 2017). Although the potential pathways other than Wnt signaling exist in EMT induction and the mechanism underlain cancer malignancy, Wnt signaling is one of the main pathways to induce EMT and cancer malignancy (Polakis, 2012).

MIE => KE1: Increases in cellular ROS leads to porcupine-induced Wnt secretion and Wnt signaling activation	Empirical Support of the MIE => KE1 is moderate. Rationale: Production of ROS and DNA double-strand break cause the tissue damages (Gao et al., 2019). ROS-related signaling induces Wnt/b-catenin pathway activation (Pérez et al., 2017).
KE1 => KE2: Porcupine-induced Wnt secretion and Wnt signaling activation leads to beta-catenin activation	Empirical Support of the KE1 => KE2 is moderate. Rationale: Dishevelled (DVL), a positive regulator of Wnt signaling, form the complex with FZD and lead to trigger the Wnt signaling together with Wnt coreceptor low-density lipoprotein (LDL) receptor-related protein 6 (LRP6) (Clevers & Nusse, 2012; Jiang et al., 2015). Wnt binds to FZD and activate the Wnt signaling (Clevers & Nusse, 2012; Janda et al., 2012; Nile et al., 2017). Wnt binding towards FZD induce the formation of the protein complex with LRP5/6 and DVL, leading to the down-stream signaling activation including beta-catenin (Clevers & Nusse, 2012).
KE2 => KE3: beta-catenin activation leads to Epithelial-mesenchymal transition (EMT)	Empirical Support of the KE2 => KE3 is moderate. Rationale: The inhibition of c-MET, which is overexpressed in diffuse-type gastric cancer, induced increase in phosphorylated b-catenin, decrease in b-catenin and Snail (Sohn et al., 2019). The garcinol, that has anti-cancer effect, increases phosphorylated beta-catenin, decreases b-catenin and ZEB1/ZEB2, and inhibit EMT (Ahmad et al., 2012). The inhibition of sortilin by AF38469 (a sortilin inhibitor) or small interference RNA (siRNA) results in decrease in b-catenin and Twist expression in human glioblastoma cells (Yang W. et al., 2019). Histone deacetylase inhibitors effect on EMT-related transcription factors including ZEB, Twist and Snail (Wawruszak et al., 2019). Snail and Zeb induces EMT and suppress E-cadherin (CDH1) (Batlle et al., 2000; Diaz et al., 2014; Peinado et al., 2007).
KE3 => AO: Epithelial-mesenchymal transition (EMT) leads to Treatment-resistant gastric cancer	Empirical Support of the KE3 => AO is moderate. Rationale: EMT activation induces the expression of multiple members of the ATP-binding cassette (ABC) transporter family, which results in doxorubicin resistance (Saxena et al., 2011; Shibue & Weinberg, 2017). TGFb-1 induced EMT results in the acquisition of cancer stem cell (CSC) like properties (Pirozzi et al., 2011; Shibue & Weinberg, 2017). Snail-induced EMT induces the cancer metastasis and resistance to dendritic cell-mediated immunotherapy (Kudo-Saito et al., 2009). Zinc finger E-box-binding homeobox (ZEB1)-induced EMT results in relief of miR-200-mediated repression of programmed cell death 1 ligand (PD-L1) expression, a major inhibitory ligand for the programmed cell death protein (PD-1) immune-checkpoint protein on CD8+ cytotoxic T lymphocyte (CTL), subsequently the CD8+ T cell immunosuppression and metastasis (Chen et al., 2014).

Quantitative Consideration

Wnt signaling activates the CSCs to promote cancer malignancy (Reya & Clevers, 2005). The responses in KEs related to Wnt signaling, Frizzled activation, GSK3beta inactivation, beta-catenin activation, Snail, Zeb, and Twist activation are dose-dependently related. The quantification of EMT and cancer malignancy would require further investigation.

Considerations for Potential Applications of the AOP (optional)

AOP entitled "Increase in reactive oxygen species (ROS) and chronic ROS leading to human treatment-resistant gastric cancer" might be utilized for the development and risk assessment of anti-cancer drugs. EMT is involved in the acquisition of drug resistance, which is one of the critical features of cancer malignancy. The assessment of the activity of the EMT network would serve as a prediction of the adverse effects of or responsiveness to anti-cancer drugs (Tanabe et al., 2023). The detection methods for increases in ROS in this AOP have future regulatory potentials to assess the human health effects of radiation or ROS-related diseases. The detection methods for human treatment-resistant gastric cancer have future regulatory potentials to diagnose the diseases.

References

- Ahmad, A., Sarkar, S. H., Bitar, B., Ali, S., Aboukameel, A., Sethi, S., . . . Sarkar, F. H. (2012). Garcinol regulates EMT and Wnt signaling pathways in vitro and in vivo, leading to anticancer activity against breast cancer cells. *Mol Cancer Ther*, 11(10), 2193-2201. doi:10.1158/1535-7163.MCT-12-0232-T
- Batlle, E., Sancho, E., Francí, C., Domínguez, D., Monfar, M., Baulida, J., & García de Herreros, A. (2000). The transcription factor Snail is a repressor of E-cadherin gene expression in epithelial tumour cells. *Nature Cell Biology*, 2(2), 84-89. doi:10.1038/35000034
- Chen, L., Gibbons, D. L., Goswami, S., Cortez, M. A., Ahn, Y.-H., Byers, L. A., . . . Qin, F. X.-F. (2014). Metastasis is regulated via microRNA-200/ZEB1 axis control of tumour cell PD-L1 expression and intratumoral immunosuppression. *Nature communications*, 5, 5241-5241. doi:10.1038/ncomms6241
- Clevers, H., & Nusse, R. (2012). Wnt/beta-catenin signaling and disease. *Cell*, 149(6), 1192-1205. doi:10.1016/j.cell.2012.05.012
- Diaz, V. M., Vinas-Castells, R., & Garcia de Herreros, A. (2014). Regulation of the protein stability of EMT transcription factors. *Cell Adh Migr*, 8(4), 418-428. doi:10.4161/19336918.2014.969998
- Gao, Q., Zhou, G., Lin, S.-J., Paus, R., & Yue, Z. (2019). How chemotherapy and radiotherapy damage the tissue: Comparative biology lessons from feather and hair models. *Experimental dermatology*, 28(4), 413-418. doi:10.1111/exd.13846
- Gu, H., Huang, Y. Shen, Y. Liu, F. Zhou, Y. Jin, et al. (2018). Reactive Oxygen Species-Mediated Tumor Microenvironment Transformation: The Mechanism of Radioresistant Gastric Cancer. *Oxid Med Cell Longev* 2018 Vol. 2018 Pages 5801209. doi:10.1155/2018/5801209
- Janda, C. Y., Waghray, D., Levin, A. M., Thomas, C., & Garcia, K. C. (2012). Structural basis of Wnt recognition by Frizzled. *Science*, 337(6090), 59-64. doi:10.1126/science.1222879
- Kudo-Saito, C., Shirako, H., Takeuchi, T., & Kawakami, Y. (2009). Cancer Metastasis Is Accelerated through Immunosuppression during Snail-Induced EMT of Cancer Cells. *Cancer Cell*, 15(3), 195-206. doi: 10.1016/j.ccr.2009.01.023
- Lazarova, D., & Bordonaro, M. (2017). ZEB1 Mediates Drug Resistance and EMT in p300-Deficient CRC. *Journal of Cancer*, 8(8), 1453-1459. doi:10.7150/jca.18762
- Nile, A. H., Mukund, S., Stanger, K., Wang, W., & Hannoush, R. N. (2017). Unsaturated fatty acyl recognition by Frizzled receptors mediates dimerization upon Wnt ligand binding. *Proc Natl Acad Sci U S A*, 114(16), 4147-4152. doi:10.1073/pnas.1618293114
- Pearlman, R. L., Montes de Oca, M. K., Pal, H. C., & Afaq, F. (2017). Potential therapeutic targets of epithelial-mesenchymal transition in melanoma. *Cancer Lett*, 391, 125-140. doi:10.1016/j.canlet.2017.01.029
- Peinado, H., Olmeda, D., & Cano, A. (2007). Snail, Zeb and bHLH factors in tumour progression: an alliance against the epithelial phenotype? *Nat Rev Cancer*, 7(6), 415-428. doi:10.1038/nrc2131
- Pérez, S., Taléns-Visconti, R., Rius-Pérez, S., Finamor, I., & Sastre, J. (2017). Redox signaling in the gastrointestinal tract. *Free radical biology & medicine*, 104, 75-103. doi:10.1016/j.freeradbiomed.2016.12.048
- Pirozzi, G., Tirino, V., Camerlingo, R., Franco, R., La Rocca, A., Liguori, E., . . . Rocco, G. (2011). Epithelial to mesenchymal transition by TGFβ-1 induction increases stemness characteristics in primary non small cell lung cancer cell line. *PLoS One*, 6(6), e21548-e21548. doi:10.1371/journal.pone.0021548
- Polakis, P. (2012). Wnt signaling in cancer. *Cold Spring Harb Perspect Biol*, 4(5). doi:10.1101/cshperspect.a008052
- Reya, T., & Clevers, H. (2005). Wnt signalling in stem cells and cancer. *Nature*, 434(7035), 843-850. doi:10.1038/nature03319
- Saxena, M., Stephens, M. A., Pathak, H., & Rangarajan, A. (2011). Transcription factors that mediate epithelial-mesenchymal transition lead to multidrug resistance by upregulating ABC transporters. *Cell death & disease*, 2(7), e179-e179. doi:10.1038/cddis.2011.61
- Shibue, T., & Weinberg, R. A. (2017). EMT, CSCs, and drug resistance: the mechanistic link and clinical implications. *Nat Rev Clin Oncol*, 14(10), 611-629. doi:10.1038/nrclinonc.2017.44
- Sohn, S. H., Kim, B., Sul, H. J., Kim, Y. J., Kim, H. S., Kim, H., . . . Zang, D. Y. (2019). INC280 inhibits Wnt/beta-catenin and EMT signaling pathways and its induce apoptosis in diffuse gastric cancer positive for c-MET amplification. *BMC Res Notes*, 12(1), 125. doi:10.1186/s13104-019-4163-x
- Tanabe, S. (2013). Perspectives of gene combinations in phenotype presentation. *World journal of stem cells*, 5(3), 61-67. doi:10.4252/wjsc.v5.i3.61
- Tanabe, S. (2015a). Origin of cells and network information. *World journal of stem cells*, 7(3), 535-540. doi:10.4252/wjsc.v7.i3.535
- Tanabe, S. (2015b). Signaling involved in stem cell reprogramming and differentiation. *World journal of stem cells*, 7(7), 992-998. doi:10.4252/wjsc.v7.i7.992
- Tanabe, S. (2015c). Overview of gene regulation in stem cell network to identify therapeutic targets utilizing genome databases. *Insights Stem Cells*, 1(1).
- Tanabe, S. (2018). Wnt Signaling and Epithelial-Mesenchymal Transition Network in Cancer. *Research Journal of Oncology*, 2(2), 3.
- Tanabe, S., Aoyagi, K., Yokozaki, H., & Sasaki, H. (2015). Regulated genes in mesenchymal stem cells and gastric cancer. *World journal of stem cells*, 7(1), 208-222. doi:10.4252/wjsc.v7.i1.208
- Tanabe, S., Quader, S., Cabral, H., & Ono, R. (2020a). Interplay of EMT and CSC in Cancer and the Potential Therapeutic Strategies. *Front. Pharmacol.* 11:904. doi: 10.3389/fphar.2020.00904

Tanabe, S., Quader, S., Ono, R., Cabral, H., Aoyagi, K., Hirose, A., Yokozaki, H., & Sasaki, H. (2020b). Molecular Network Profiling in Intestinal- and Diffuse-Type Gastric Cancer. *Cancers*, 12(12), 3833. <https://doi.org/10.3390/cancers12123833>

Tanabe, S., Quader, S., Ono, R., Cabral, H., Aoyagi, K., Hirose, A., Perkins, E.J., Yokozaki, H., & Sasaki, H. (2023). Regulation of Epithelial-Mesenchymal Transition Pathway and Artificial Intelligence-Based Modeling for Pathway Activity Prediction. *Onco*, 3(1):13-25. doi: 10.3390/onco3010002

Vallée, A., & Lecarpentier, Y. (2018). Crosstalk Between Peroxisome Proliferator-Activated Receptor Gamma and the Canonical WNT/ β -Catenin Pathway in Chronic Inflammation and Oxidative Stress During Carcinogenesis. *Frontiers in immunology*, 9, 745-745. doi:10.3389/fimmu.2018.00745

Wawruszak, A., Kalafut, J., Okon, E., Czapinski, J., Halasa, M., Przybyszewska, A., . . . Stepulak, A. (2019). Histone Deacetylase Inhibitors and Phenotypical Transformation of Cancer Cells. *Cancers (Basel)*, 11(2). doi:10.3390/cancers11020148

Yang, W., Wu, P. F., Ma, J. X., Liao, M. J., Wang, X. H., Xu, L. S., . . . Yi, L. (2019). Sortilin promotes glioblastoma invasion and mesenchymal transition through GSK-3 β /beta-catenin/twist pathway. *Cell Death Dis*, 10(3), 208. doi:10.1038/s41419-019-1449-9

Appendix 1

List of MIEs in this AOP

Event: 1115: Increase, Reactive oxygen species

Short Name: Increase, ROS

Key Event Component

Process	Object	Action
reactive oxygen species biosynthetic process	reactive oxygen species	increased

AOPs Including This Key Event

AOP ID and Name	Event Type
Aop:186 - unknown MIE leading to renal failure and mortality	KeyEvent
Aop:213 - Inhibition of fatty acid beta oxidation leading to nonalcoholic steatohepatitis (NASH)	KeyEvent
Aop:303 - Frustrated phagocytosis-induced lung cancer	KeyEvent
Aop:383 - Inhibition of Angiotensin-converting enzyme 2 leading to liver fibrosis	KeyEvent
Aop:382 - Angiotensin II type 1 receptor (AT1R) agonism leading to lung fibrosis	KeyEvent
Aop:384 - Hyperactivation of ACE/Ang-II/AT1R axis leading to chronic kidney disease	KeyEvent
Aop:396 - Deposition of ionizing energy leads to population decline via impaired meiosis	KeyEvent
Aop:409 - Frustrated phagocytosis leads to malignant mesothelioma	KeyEvent
Aop:413 - Oxidation and antagonism of reduced glutathione leading to mortality via acute renal failure	KeyEvent
Aop:416 - Aryl hydrocarbon receptor activation leading to lung cancer through IL-6 toxicity pathway	KeyEvent
Aop:418 - Aryl hydrocarbon receptor activation leading to impaired lung function through AHR-ARNT toxicity pathway	KeyEvent
Aop:386 - Deposition of ionizing energy leading to population decline via inhibition of photosynthesis	KeyEvent
Aop:387 - Deposition of ionising energy leading to population decline via mitochondrial dysfunction	KeyEvent
Aop:319 - Binding to ACE2 leading to lung fibrosis	KeyEvent
Aop:451 - Interaction with lung resident cell membrane components leads to lung cancer	KeyEvent
Aop:476 - Adverse Outcome Pathways diagram related to PBDEs associated male reproductive toxicity	MolecularInitiatingEvent
Aop:492 - Glutathione conjugation leading to reproductive dysfunction via oxidative stress	KeyEvent
Aop:497 - ERα inactivation alters mitochondrial functions and insulin signalling in skeletal muscle and leads to insulin resistance and metabolic syndrome	KeyEvent
Aop:500 - Activation of MEK-ERK1/2 leads to deficits in learning and cognition via ROS and apoptosis	KeyEvent
Aop:505 - Reactive Oxygen Species (ROS) formation leads to cancer via inflammation pathway	MolecularInitiatingEvent
Aop:513 - Reactive Oxygen (ROS) formation leads to cancer via Peroxisome proliferation-activated receptor (PPAR) pathway	MolecularInitiatingEvent
Aop:521 - Essential element imbalance leads to reproductive failure via oxidative stress	KeyEvent
Aop:540 - Oxidative Stress in the Fish Ovary Leads to Reproductive Impairment via Reduced Vitellogenin Production	MolecularInitiatingEvent
Aop:462 - Activation of reactive oxygen species leading the atherosclerosis	MolecularInitiatingEvent
Aop:299 - Deposition of energy leading to population decline via DNA oxidation and follicular atresia	KeyEvent
Aop:311 - Deposition of energy leading to population decline via DNA oxidation and oocyte apoptosis	KeyEvent
Aop:325 - Excessive reactive oxygen species leading to growth inhibition via lipid peroxidation and cell death	MolecularInitiatingEvent

AOP ID and Name	Event Type
Aop:332 - Excessive reactive oxygen species leading to growth inhibition via lipid peroxidation and reduced cell proliferation	MolecularInitiatingEvent
Aop:324 - Excessive reactive oxygen species leading to growth inhibition via oxidative DNA damage and cell death	MolecularInitiatingEvent
Aop:331 - Excessive reactive oxygen species leading to growth inhibition via oxidative DNA damage and reduced cell proliferation	MolecularInitiatingEvent
Aop:326 - Excessive reactive oxygen species leading to growth inhibition via protein oxidation and cell death	MolecularInitiatingEvent
Aop:333 - Excessive reactive oxygen species leading to growth inhibition via uncoupling of oxidative phosphorylation	MolecularInitiatingEvent
Aop:327 - Excessive reactive oxygen species production leading to mortality (1)	MolecularInitiatingEvent
Aop:328 - Excessive reactive oxygen species production leading to mortality (2)	MolecularInitiatingEvent
Aop:329 - Excessive reactive oxygen species production leading to mortality (3)	MolecularInitiatingEvent
Aop:330 - Excessive reactive oxygen species production leading to mortality (4)	MolecularInitiatingEvent
Aop:26 - Calcium-mediated neuronal ROS production and energy imbalance	KeyEvent
Aop:534 - Succinate dehydrogenase (SDH) inhibition leads to cancer through oxidative stress	KeyEvent
Aop:273 - Mitochondrial complex inhibition leading to liver injury	KeyEvent
Aop:488 - Increased reactive oxygen species production leading to decreased cognitive function	MolecularInitiatingEvent
Aop:298 - Increase in reactive oxygen species (ROS) leading to human treatment-resistant gastric cancer	MolecularInitiatingEvent
Aop:27 - Cholestatic Liver Injury induced by Inhibition of the Bile Salt Export Pump (ABCB11)	KeyEvent
Aop:511 - The AOP framework on ROS-mediated oxidative stress induced vascular disrupting effects	MolecularInitiatingEvent
Aop:207 - NADPH oxidase and P38 MAPK activation leading to reproductive failure in Caenorhabditis elegans	KeyEvent
Aop:423 - Toxicological mechanisms of hepatocyte apoptosis through the PARP1 dependent cell death pathway	MolecularInitiatingEvent
Aop:481 - AOPs of amorphous silica nanoparticles: ROS-mediated oxidative stress increased respiratory dysfunction and diseases.	MolecularInitiatingEvent
Aop:282 - Adverse outcome pathway on photochemical toxicity initiated by light exposure	MolecularInitiatingEvent
Aop:569 - Decreased DNA methylation of FAM50B/PTCHD3 leading to IQ loss of children via PI3K-Akt pathway	KeyEvent

Biological Context

Level of Biological Organization

Cellular

Cell term

Cell term

cell

Organ term

Organ term

organ

Domain of Applicability

Taxonomic Applicability

Term	Scientific Term	Evidence	Links
Vertebrates	Vertebrates	High	NCBI
human	Homo sapiens	Moderate	NCBI
human and other cells in culture	human and other cells in culture	Moderate	NCBI
mouse	Mus musculus	Moderate	NCBI
crustaceans	Daphnia magna	High	NCBI
Lemna minor	Lemna minor	High	NCBI
zebrafish	Danio rerio	High	NCBI

Life Stage Applicability

Life Stage	Evidence
All life stages	High

Sex Applicability

Sex	Evidence
Unspecific	High

Sex Evidence

Mixed High

ROS is a normal constituent found in all organisms, *lifestages*, and *sexes*.

Key Event Description

Biological State: increased reactive oxygen species (ROS)

Biological compartment: an entire cell -- may be cytosolic, may also enter organelles.

Reactive oxygen species (ROS) are O_2^- -derived molecules that can be both free radicals (e.g. superoxide, hydroxyl, peroxy, alkoxy) and non-radicals (hypochlorous acid, ozone and singlet oxygen) (Bedard and Krause 2007; Ozcan and Ogun 2015). ROS production occurs naturally in all kinds of tissues inside various cellular compartments, such as mitochondria and peroxisomes (Drew and Leeuwenburgh 2002; Ozcan and Ogun 2015). Furthermore, these molecules have an important function in the regulation of several biological processes – they might act as antimicrobial agents or triggers of animal gamete activation and capacitation (Goud et al. 2008; Parrish 2010; Bisht et al. 2017). However, in environmental stress situations (exposure to radiation, chemicals, high temperatures) these molecules have its levels drastically increased, and overly interact with macromolecules, namely nucleic acids, proteins, carbohydrates and lipids, causing cell and tissue damage (Brieger et al. 2012; Ozcan and Ogun 2015).

Reactive oxygen species (ROS) refers to the chemical species superoxide, hydrogen peroxide, and their secondary reactive products. In the biological context, ROS are signaling molecules with important roles in cell energy metabolism, cell proliferation, and fate. Therefore, balancing ROS levels at the cellular and tissue level is an important part of many biological processes. Disbalance, mainly an increase in ROS levels, can cause cell dysfunction and irreversible cell damage.

ROS are produced from both exogenous stressors and normal endogenous cellular processes, such as the mitochondrial electron transport chain (ETC). Inhibition of the ETC can result in the accumulation of ROS. Exposure to chemicals, heavy metal ions, or ionizing radiation can also result in increased production of ROS. Chemicals and heavy metal ions can deplete cellular antioxidants reducing the cell's ability to control cellular ROS and resulting in the accumulation of ROS. Cellular antioxidants include glutathione (GSH), protein sulfhydryl groups, superoxide dismutase (SOD).

ROS are radicals, ions, or molecules that have a single unpaired electron in their outermost shell of electrons, which can be categorized into two groups: free oxygen radicals and non-radical ROS [Liou et al., 2010].

<Free oxygen radicals>

superoxide	$O_2^{\cdot -}$
hydroxyl radical	$\cdot OH$
nitric oxide	$NO\cdot$
organic radicals	$R\cdot$
peroxyl radicals	$ROO\cdot$
alkoxyl radicals	$RO\cdot$
thiyl radicals	$RS\cdot$
sulfonyl radicals	$ROS\cdot$
thiyl peroxy radicals	$RSOO\cdot$
disulfides	$RSSR$

<Non-radical ROS>

hydrogen peroxide	H_2O_2
singlet oxygen	1O_2
ozone/trioxygen	O_3
organic hydroperoxides	$ROOH$
hypochlorite	ClO^-
peroxynitrite	$ONOO^-$
nitrosoperoxycarbonate anion	$O=NOOCO_2^-$
nitrocarbonate anion	$O_2NOCO_2^-$
dinitrogen dioxide	N_2O_2
nitronium	NO_2^+
highly reactive lipid- or carbohydrate-derived carbonyl compounds	

Potential sources of ROS include NADPH oxidase, xanthine oxidase, mitochondria, nitric oxide synthase, cytochrome P450, lipoxygenase/cyclooxygenase, and monoamine oxidase [Granger et al., 2015]. ROS are generated through NADPH oxidases consisting of p47^{phox} and p67^{phox}. ROS are generated through xanthine oxidase activation in sepsis [Ramos et al., 2018]. Arsenic produces ROS [Zhang et al., 2011]. Mitochondria-targeted paraquat and metformin mediate ROS production [Chowdhury et al., 2020]. ROS are generated by bleomycin [Lu et al., 2010]. Radiation induces dose-dependent ROS production [Ji et al., 2019].

ROS are generated in the course of cellular respiration, metabolism, cell signaling, and inflammation [Dickinson and Chang 2011; Egea et al. 2017]. Hydrogen peroxide is also made by the endoplasmic reticulum in the course of protein folding. Nitric oxide (NO) is produced at the highest levels by nitric oxide synthase in endothelial cells and phagocytes. NO production is one of the main mechanisms by which phagocytes kill bacteria [Wang et al., 2017]. The other species are produced by reactions with superoxide or peroxide, or by other free radicals or enzymes.

ROS activity is principally local. Most ROS have short half-lives, ranging from nano- to milliseconds, so diffusion is limited, while reactive nitrogen species (RNS) nitric oxide or peroxynitrite can survive long enough to diffuse across membranes [Calcerrada et al. 2011]. Consequently, local concentrations of ROS are much higher than average cellular concentrations, and signaling is typically controlled by colocalization with redox buffers [Dickinson and Chang 2011; Egea et al. 2017].

Although their existence is limited temporally and spatially, ROS interact with other ROS or with other nearby molecules to produce more ROS and participate in a feedback loop to amplify the ROS signal, which can increase RNS. Both ROS and RNS also move into neighboring cells, and ROS can increase intracellular ROS signaling in neighboring cells [Egea et al. 2017].

In the primary event, photoreactive chemicals are excited by the absorption of photon energy. The energy of the photoactivated chemicals transfer to oxygen and then generates the reactive oxygen species (ROS), including superoxide ($O_2^{\cdot -}$) via type I reaction and singlet oxygen (1O_2) via type II reaction, as principal intermediate species in phototoxic reaction (Foote, 1991, Onoue et al., 2009).

How it is Measured or Detected

Photocolorimetric assays (Sharma et al. 2017; Griending et al. 2016) or through commercial kits purchased from specialized companies.

Yuan, Yan, et al., (2013) described ROS monitoring by using H₂-DCF-DA, a redox-sensitive fluorescent dye. Briefly, the harvested cells were incubated with H₂-DCF-DA (50 µmol/L final concentration) for 30 min in the dark at 37°C. After treatment, cells were immediately washed twice, re-suspended in PBS, and analyzed on a BD-FACS Aria flow cytometry. ROS generation was based on fluorescent intensity which was recorded by excitation at 504 nm and emission at 529 nm.

Lipid peroxidation (LPO) can be measured as an indicator of oxidative stress damage Yen, Cheng Chien, et al., (2013).

Chattopadhyay, Sukumar, et al. (2002) assayed the generation of free radicals within the cells and their extracellular release in the medium by addition of yellow NBT salt solution (Park et al., 1968). Extracellular release of ROS converted NBT to a purple colored formazan. The cells were incubated with 100 µl of 1 mg/ml NBT solution for 1 h at 37 °C and the product formed was assayed at 550 nm in an Anthos 2001 plate reader. The observations of the 'cell-free system' were confirmed by cytological examination of parallel set of explants stained with chromogenic reactions for NO and ROS.

On the basis of the pathogenesis of drug-induced phototoxicity, a reactive oxygen species (ROS) assay was proposed to evaluate the phototoxic risk of chemicals. The ROS assay can monitor generation of ROS, such as singlet oxygen and superoxide, from photoirradiated chemicals, and the ROS data can be used to evaluate the photoreactivity of chemicals (Onoue et al. , 2014, Onoue et al. , 2013, Onoue and Tsuda, 2006). The ROS assay is a recommended approach by guidelines to evaluate the phototoxic risk of chemicals (ICH, 2014, PCPC, 2014).

<Direct detection>

Many fluorescent compounds can be used to detect ROS, some of which are specific, and others are less specific.

□ ROS can be detected by fluorescent probes such as *p*-methoxy-phenol derivative [Ashoka et al., 2020].

□ Chemiluminescence analysis can detect the superoxide, where some probes have a wider range for detecting hydroxyl radical, hydrogen peroxide, and peroxynitrite [Fuloria et al., 2021].

□ ROS in the blood can be detected using superparamagnetic iron oxide nanoparticles (SPION)-based biosensor [Lee et al., 2020].

□ Hydrogen peroxide (H₂O₂) can be detected with a colorimetric probe, which reacts with H₂O₂ in a 1:1 stoichiometry to produce a bright pink colored product, followed by the detection with a standard colorimetric microplate reader with a filter in the 540-570 nm range.

□ The levels of ROS can be quantified using multiple-step amperometry using a stainless steel counter electrode and non-leak Ag|AgCl reference node [Flaherty et al., 2017].

□ Singlet oxygen can be measured by monitoring the bleaching of *p*-nitrosodimethylaniline at 440 nm using a spectrophotometer with imidazole as a selective acceptor of singlet oxygen [Onoue et al., 2014].

<Indirect Detection>

Alternative methods involve the detection of redox-dependent changes to cellular constituents such as proteins, DNA, lipids, or glutathione [Dickinson and Chang 2011; Wang et al. 2013; Griending et al. 2016]. However, these methods cannot generally distinguish between the oxidative species behind the changes and cannot provide good resolution for the kinetics of oxidative activity.

References

- Akai, K., et al. (2004). "Ability of ferric nitrilotriacetate complex with three pH-dependent conformations to induce lipid peroxidation." *Free Radic Res. Sep*;38(9):951-62. doi: 10.1080/1071576042000261945
- Ashoka, A. H., et al. (2020). "Recent Advances in Fluorescent Probes for Detection of HOCl and HNO." *ACS omega*, 5(4), 1730-1742. doi:10.1021/acsomega.9b03420
- B.H. Park, S.M. Fikrig, E.M. Smithwick Infection and nitroblue tetrazolium reduction by neutrophils: a diagnostic aid *Lancet*, 2 (1968), pp. 532-534
- Bedard, Karen, and Karl-Heinz Krause. 2007. "The NOX Family of ROS-Generating NADPH Oxidases: Physiology and Pathophysiology." *Physiological Reviews* 87 (1): 245-313.
- Bisht, Shilpa, Muneeb Faiq, Madhuri Tolahunase, and Rima Dada. 2017. "Oxidative Stress and Male Infertility." *Nature Reviews. Urology* 14 (8): 470-85.
- Brieger, K., S. Schiavone, F. J. Miller Jr, and K-H Krause. 2012. "Reactive Oxygen Species: From Health to Disease." *Swiss Medical Weekly* 142 (August): w13659.
- Calcerrada, P., et al. (2011). "Nitric oxide-derived oxidants with a focus on peroxynitrite: molecular targets, cellular responses and therapeutic implications." *Curr Pharm Des* 17(35): 3905-3932.
- Chattopadhyay, Sukumar, et al. "Apoptosis and necrosis in developing brain cells due to arsenic toxicity and protection with antioxidants." *Toxicology letters* 136.1 (2002): 65-76.
- Chowdhury, A. R., et al. (2020). "Mitochondria-targeted paraquat and metformin mediate ROS production to induce multiple pathways of retrograde signaling: A dose-dependent phenomenon." *Redox Biol.* doi: 10.1016/j.redox.2020.101606. PMID: 32604037; PMCID: PMC7327929.
- Dickinson, B. C. and Chang C. J. (2011). "Chemistry and biology of reactive oxygen species in signaling or stress responses." *Nature chemical biology* 7(8): 504-511.
- Drew, Barry, and Christiaan Leeuwenburgh. 2002. "Aging and the Role of Reactive Nitrogen Species." *Annals of the New York Academy of Sciences* 959 (April): 66-81.
- Egea, J., et al. (2017). "European contribution to the study of ROS: A summary of the findings and prospects for the future from the COST action BM1203 (EU-ROS)." *Redox biology* 13: 94-162.
- Flaherty, R. L., et al. (2017). "Glucocorticoids induce production of reactive oxygen species/reactive nitrogen species and DNA damage through an iNOS mediated pathway in breast cancer." *Breast Cancer Research*, 19(1), 1-13. <https://doi.org/10.1186/s13058-017-0823-8>
- Foot CS. Definition of type I and type II photosensitized oxidation. *Photochem Photobiol.* 1991;54:659.
- Fuloria, S., et al. (2021). "Comprehensive Review of Methodology to Detect Reactive Oxygen Species (ROS) in Mammalian Species and Establish Its Relationship with Antioxidants and Cancer." *Antioxidants (Basel, Switzerland)* 10(1) 128. doi:10.3390/antiox10010128
- Go, Y. M. and Jones, D. P. (2013). "The redox proteome." *J Biol Chem* 288(37): 26512-26520.

- Goud, Anuradha P., Pravin T. Goud, Michael P. Diamond, Bernard Gonik, and Husam M. Abu-Soud. 2008. "Reactive Oxygen Species and Oocyte Aging: Role of Superoxide, Hydrogen Peroxide, and Hypochlorous Acid." *Free Radical Biology & Medicine* 44 (7): 1295–1304.
- Granger, D. N. and Kvietys, P. R. (2015). "Reperfusion injury and reactive oxygen species: The evolution of a concept" *Redox Biol.* doi: 10.1016/j.redox.2015.08.020. PMID: 26484802; PMCID: PMC4625011.
- Griendling, K. K., et al. (2016). "Measurement of Reactive Oxygen Species, Reactive Nitrogen Species, and Redox-Dependent Signaling in the Cardiovascular System: A Scientific Statement From the American Heart Association." *Circulation research* 119(5): e39-75.
- Griendling, Kathy K., Rhian M. Touyz, Jay L. Zweier, Sergey Dikalov, William Chilian, Yeong-Renn Chen, David G. Harrison, Aruni Bhatnagar, and American Heart Association Council on Basic Cardiovascular Sciences. 2016. "Measurement of Reactive Oxygen Species, Reactive Nitrogen Species, and Redox-Dependent Signaling in the Cardiovascular System: A Scientific Statement From the American Heart Association." *Circulation Research* 119 (5): e39-75.
- ICH. ICH Guideline S10 Guidance on Photosafety Evaluation of Pharmaceuticals.: International Council on Harmonisation of Technical Requirements for Registration of Pharmaceuticals for Human Use; 2014.
- Itziou, A., et al. (2011). "In vivo and in vitro effects of metals in reactive oxygen species production, protein carbonylation, and DNA damage in land snails *Eobania vermiculata*." *Archives of Environmental Contamination and Toxicology*, 60(4), 697–707. <https://doi.org/10.1007/s00244-010-9583-5>
- Ji, W. O., et al. "Quantitation of the ROS production in plasma and radiation treatments of biotargets." *Sci Rep.* 2019 Dec 27;9(1):19837. doi: 10.1038/s41598-019-56160-0. PMID: 31882663; PMCID: PMC6934759.
- Kruk, J. and Aboul-Enein, H. Y. (2017). "Reactive Oxygen and Nitrogen Species in Carcinogenesis: Implications of Oxidative Stress on the Progression and Development of Several Cancer Types." *Mini-Reviews in Medicinal Chemistry*, 17:11. doi:10.2174/1389557517666170228115324
- Lee, D. Y., et al. (2020). "PEGylated Bilirubin-coated Iron Oxide Nanoparticles as a Biosensor for Magnetic Relaxation Switching-based ROS Detection in Whole Blood." *Theranostics*, 10(5), 1997-2007. doi:10.7150/tno.39662
- Li, Z., et al. (2020). "Inhibition of MiR-25 attenuates doxorubicin-induced apoptosis, reactive oxygen species production and DNA damage by targeting pten." *International Journal of Medical Sciences*, 17(10), 1415-1427. <https://doi.org/10.7150/ijms.41980>
- Liou, G. Y. and Storz, P. "Reactive oxygen species in cancer." *Free Radic Res.* 2010 May;44(5):479-96. doi:10.3109/10715761003667554. PMID: 20370557; PMCID: PMC3880197.
- Lu, Y., et al. (2010). "Phosphatidylinositol-3-kinase/akt regulates bleomycin-induced fibroblast proliferation and collagen production." *American journal of respiratory cell and molecular biology*, 42(4), 432-441. <https://doi.org/10.1165/rcmb.2009-0002OC>
- Onoue, S., et al. (2013). "Establishment and intra-/inter-laboratory validation of a standard protocol of reactive oxygen species assay for chemical photosafety evaluation." *J Appl Toxicol.* 33(11):1241-50. doi: 10.1002/jat.2776. Epub 2012 Jun 13. PMID: 22696462.
- Onoue S, Hosoi K, Toda T, Takagi H, Osaki N, Matsumoto Y, et al. Intra-/inter-laboratory validation study on reactive oxygen species assay for chemical photosafety evaluation using two different solar simulators. *Toxicology in vitro : an international journal published in association with BIBRA*. 2014;28:515-23.
- Onoue S, Hosoi K, Wakuri S, Iwase Y, Yamamoto T, Matsuoka N, et al. Establishment and intra-/inter-laboratory validation of a standard protocol of reactive oxygen species assay for chemical photosafety evaluation. *Journal of applied toxicology : JAT*. 2013;33:1241-50.
- Onoue S, Kawamura K, Igarashi N, Zhou Y, Fujikawa M, Yamada H, et al. Reactive oxygen species assay-based risk assessment of drug-induced phototoxicity: classification criteria and application to drug candidates. *J Pharm Biomed Anal.* 2008;47:967-72.
- Onoue S, Seto Y, Gandy G, Yamada S. Drug-induced phototoxicity; an early *in vitro* identification of phototoxic potential of new drug entities in drug discovery and development. *Current drug safety.* 2009;4:123-36.
- Onoue S, Tsuda Y. Analytical studies on the prediction of photosensitive/phototoxic potential of pharmaceutical substances. *Pharmaceutical research.* 2006;23:156-64.
- Ozcan, Ayla, and Metin Ogun. 2015. "Biochemistry of Reactive Oxygen and Nitrogen Species." In *Basic Principles and Clinical Significance of Oxidative Stress*, edited by Sivakumar Joghi Thatha Gowder. Rijeka: IntechOpen.
- Parrish, A. R. 2010. "2.27 - Hypoxia/Ischemia Signaling." In *Comprehensive Toxicology (Second Edition)*, edited by Charlene A. McQueen, 529–42. Oxford: Elsevier.
- PCPC. PCPC 2014 safety evaluation guidelines; Chapter 7: Evaluation of Photoirritation and Photoallergy potential. Personal Care Products Council; 2014.
- Ramos, M. F. P., et al. (2018). "Xanthine oxidase inhibitors and sepsis." *Int J Immunopathol Pharmacol.* 32:2058738418772210. doi:10.1177/2058738418772210
- Ravanat, J. L., et al. (2014). "Radiation-mediated formation of complex damage to DNA: a chemical aspect overview." *Br J Radiol* 87(1035): 20130715.
- Schutzendubel, A. and Polle, A. (2002). "Plant responses to abiotic stresses: heavy metal-induced oxidative stress and protection by mycorrhization." *Journal of Experimental Botany*, 53(372), 1351–1365. <https://doi.org/10.1093/jexbot/53.372.1351>
- Seto Y, Kato M, Yamada S, Onoue S. Development of micellar reactive oxygen species assay for photosafety evaluation of poorly water-soluble chemicals. *Toxicology in vitro : an international journal published in association with BIBRA*. 2013;27:1838-46.
- Sharma, Gunjan, Nishant Kumar Rana, Priya Singh, Pradeep Dubey, Daya Shankar Pandey, and Biplob Koch. 2017. "p53 Dependent Apoptosis and Cell Cycle Delay Induced by Heteroleptic Complexes in Human Cervical Cancer Cells." *Biomedicine & Pharmacotherapy = Biomedecine & Pharmacotherapie* 88 (April): 218–31.
- Silva, R., et al. (2019). "Light exposure during growth increases riboflavin production, reactive oxygen species accumulation and DNA damage in *Ashbya gossypii* riboflavin-overproducing strains." *FEMS Yeast Research*, 19(1), 1–7. <https://doi.org/10.1093/femsyr/foy114>
- Tsuchiya K, et al. (2005). "Oxygen radicals photo-induced by ferric nitrilotriacetate complex." *Biochim Biophys Acta.* 1725(1):111-9. doi:10.1016/j.bbagen.2005.05.001
- Wang, J., et al. (2017). "Glucocorticoids Suppress Antimicrobial Autophagy and Nitric Oxide Production and Facilitate Mycobacterial Survival in Macrophages." *Scientific reports*, 7(1), 982. <https://doi.org/10.1038/s41598-017-01174-9>
- Wang, X., et al. (2013). "Imaging ROS signaling in cells and animals." *Journal of molecular medicine* 91(8): 917-927.
- Yen, Cheng Chien, et al. "Inorganic arsenic causes cell apoptosis in mouse cerebrum through an oxidative stress-regulated signaling pathway." *Archives of toxicology* 85 (2011): 565-575.

Yuan, Yan, et al. "Cadmium-induced apoptosis in primary rat cerebral cortical neurons culture is mediated by a calcium signaling pathway." PloS one 8.5 (2013): e64330.

Zhang, Z., et al. (2011). "Reactive oxygen species mediate arsenic induced cell transformation and tumorigenesis through Wnt/ β -catenin pathway in human colorectal adenocarcinoma DLD1 cells. " Toxicology and Applied Pharmacology, 256(2), 114-121. doi:10.1016/j.taap.2011.07.016

List of Key Events in the AOP

Event: 1754: Increase, porcupine-induced Wnt secretion and Wnt signaling activation

Short Name: Increase, porcupine-induced Wnt secretion and Wnt signaling activation

Key Event Component

Process	Object	Action
Wnt protein secretion	protein-serine O-palmitoleoyltransferase porcupine	increased

AOPs Including This Key Event

AOP ID and Name	Event Type
Aop:298 - Increase in reactive oxygen species (ROS) leading to human treatment-resistant gastric cancer	KeyEvent

Stressors

Name
Radiation

Biological Context

Level of Biological Organization
Cellular

Cell term

Cell term
cell

Organ term

Organ term
organ

Domain of Applicability

Taxonomic Applicability

Term	Scientific Term	Evidence	Links
Homo sapiens	Homo sapiens	High	NCBI
Mus musculus	Mus musculus	High	NCBI

Life Stage Applicability

Life Stage	Evidence
All life stages	Moderate

Sex Applicability

Sex	Evidence
Unspecific	High

Oligomerization of FZD and low-density lipoprotein receptor-related protein 5/6 (LRP5/6) activates Wnt/ β -catenin signaling in *Homo sapiens* ([Hua et al., 2018](#)).

Key Event Description

Porcupine, which is a trans-membrane endoplasmic reticulum O-acyl transferase, is important for the secretion of Wnt ligands ([Saha et al., 2016a](#)). WNTs are secreted proteins that contain 22-24 conserved cysteine residues ([Foulquier et al., 2018](#)). The WNT molecules consist of molecular families including WNT1, WNT2, WNT2B/WNT13, WNT3, WNT4, WNT5A, WNT5B, WNT6, WNT7A, WNT7B, WNT8A, WNT8B, WNT10B, WNT11, and WNT16. ([Clevers & Nusse, 2012](#); [M. Katoh, 2001](#); [Kusserow et al., 2005](#))

Wnt proteins consist of 350-400 amino acids ([Saito-Diaz et al., 2013](#)).

WNT ligands are known to trigger at least three different downstream signaling cascades including canonical WNT/beta-catenin signaling pathway, non-canonical WNT/Ca²⁺ pathway, and planar cell polarity (PCP) pathway (De, 2011; Lai, Chien, & Moon, 2009; Willert & Nusse, 2012). WNTs bind to Frizzled proteins, which are seven-pass transmembrane receptors with an extracellular N-terminal cysteine-rich domain (Bhanot et al., 1996; Clevers, 2006). Wnt signaling begins with the binding of Wnt ligand towards the Frizzled receptors (Mohammed et al., 2016).

Wnt ligands bind to Frizzled (FZD) receptors which are seven transmembrane-domain protein receptors (Nile, Mukund, Stanger, Wang, & Hannoush, 2017). At least 10 FZD receptors are identified in human cells. FZD receptor is activated by Wnt ligand binding (MacDonald, Tamai, & He, 2009).

How it is Measured or Detected

- Secretion of WNT requires a number of other dedicated factors including the sortin receptor Wntless (WLS), which binds to Wnt and escorts it to the cell surface (Banziger et al., 2006; Ching & Nusse, 2006).
- Wnt signaling is activated by the gene mutations of the signaling components (Ziv et al., 2017).
- Wnt1, Wnt3a, and Wnt5a protein expression are measured by immunoblotting using antibodies for Wnt1, Wnt3a, and Wnt5a, respectively (L. Du et al., 2016; B. Wang et al., 2017).
- WNT2, of which expression is detected by quantitative PCR, immunoblotting, and immunohistochemistry, induces EMT (Zhou et al., 2016).
- Frizzled receptor protein level on the cell surface is measured by flow cytometry with pan-FZD antibody (Jiang et al., 2015; Zeng et al., 2018).
- DVL protein level is measured by immunoblotting with anti-DVL2 antibodies (Zeng et al., 2018).
- Fzd mRNA level is measured by quantitative reverse transcription-polymerase chain reaction (RT-PCR) (Zeng et al., 2018).
- The up-regulation of WNT ligand expression occurs in *Homo sapiens* (B. Wang et al., 2017).
- The Wnt genes play an important role in the secretion from cells, glycosylation, and tight association with the cell surface and extracellular matrix in *Drosophila melanogaster* (Willert & Nusse, 2012).

References

- Banziger, C., Soldini, D., Schutt, C., Zipperlen, P., Hausmann, G., & Basler, K. (2006). Wntless, a conserved membrane protein dedicated to the secretion of Wnt proteins from signaling cells. *Cell*, 125(3), 509-522. doi:10.1016/j.cell.2006.02.049
- Bhanot, P., Brink, M., Samos, C. H., Hsieh, J.-C., Wang, Y., Macke, J. P., . . . Nusse, R. (1996). A new member of the frizzled family from *Drosophila* functions as a Wingless receptor. *Nature*, 382, 225. doi:10.1038/382225a0
- Ching, W., & Nusse, R. (2006). A dedicated Wnt secretion factor. *Cell*, 125(3), 432-433. doi:10.1016/j.cell.2006.04.018
- Clevers, H. (2006). Wnt/beta-catenin signaling in development and disease. *Cell*, 127(3), 469-480. doi:10.1016/j.cell.2006.10.018
- Clevers, H., & Nusse, R. (2012). Wnt/beta-catenin signaling and disease. *Cell*, 149(6), 1192-1205. doi:10.1016/j.cell.2012.05.012
- De, A. (2011). Wnt/Ca²⁺ signaling pathway: a brief overview. *Acta Biochim Biophys Sin (Shanghai)*, 43(10), 745-756. doi:10.1093/abbs/gmr079
- Du, J., Zu, Y., Li, J., Du, S., Xu, Y., Zhang, L., . . . Yang, C. (2016). Extracellular matrix stiffness dictates Wnt expression through integrin pathway. *Sci Rep*, 6, 20395. doi:10.1038/srep20395
- Foulquier, S., Daskalopoulos, E. P., Lluri, G., Hermans, K. C. M., Deb, A., & Blankesteijn, W. M. (2018). WNT Signaling in Cardiac and Vascular Disease. *Pharmacol Rev*, 70(1), 68-141. doi:10.1124/pr.117.013896
- Hua, Y., Yang, Y., Li, Q., He, X., Zhu, W., Wang, J., & Gan, X. (2018). Oligomerization of Frizzled and LRP5/6 protein initiates intracellular signaling for the canonical WNT/beta-catenin pathway. *J Biol Chem*, 293(51), 19710-19724. doi:10.1074/jbc.RA118.004434
- Jiang, X., Charlat, O., Zamponi, R., Yang, Y., & Cong, F. (2015). Dishevelled promotes Wnt receptor degradation through recruitment of ZNRF3/RNF43 E3 ubiquitin ligases. *Mol Cell*, 58(3), 522-533. doi:10.1016/j.molcel.2015.03.015
- Katoh, M. (2001). Molecular cloning and characterization of human WNT3. *International journal of oncology*, 19(5), 977-982. Retrieved from <https://www.ncbi.nlm.nih.gov/pubmed/11604997>
- Kusserow, A., Pang, K., Sturm, C., Hrouda, M., Lentfer, J., Schmidt, H. A., . . . Holstein, T. W. (2005). Unexpected complexity of the Wnt gene family in a sea anemone. *Nature*, 433(7022), 156-160. doi:10.1038/nature03158
- Lai, S. L., Chien, A. J., & Moon, R. T. (2009). Wnt/Fz signaling and the cytoskeleton: potential roles in tumorigenesis. *Cell Res*, 19(5), 532-545. doi:10.1038/cr.2009.41
- MacDonald, B. T., Tamai, K., & He, X. (2009). Wnt/beta-catenin signaling: components, mechanisms, and diseases. *Dev Cell*, 17(1), 9-26. doi:10.1016/j.devcel.2009.06.016
- Mohammed, M. K., Shao, C., Wang, J., Wei, Q., Wang, X., Collier, Z., . . . Lee, M. J. (2016). Wnt/beta-catenin signaling plays an ever-expanding role in stem cell self-renewal, tumorigenesis and cancer chemoresistance. *Genes Dis*, 3(1), 11-40. doi:10.1016/j.gendis.2015.12.004
- Nile, A. H., Mukund, S., Stanger, K., Wang, W., & Hannoush, R. N. (2017). Unsaturated fatty acyl recognition by Frizzled receptors mediates dimerization upon Wnt ligand binding. *Proc Natl Acad Sci U S A*, 114(16), 4147-4152. doi:10.1073/pnas.1618293114
- Saha, S., Aranda, E., Hayakawa, Y., Bhanja, P., Atay, S., Brodin, N. P., . . . Pollard, J. W. (2016). Macrophage-derived extracellular vesicle-packaged WNTs rescue intestinal stem cells and enhance survival after radiation injury. *Nature Communications*, 7(1), 13096. doi:10.1038/ncomms13096
- Saito-Diaz, K., Chen, T. W., Wang, X., Thorne, C. A., Wallace, H. A., Page-McCaw, A., & Lee, E. (2013). The way Wnt works: components and mechanisms. *Growth Factors*, 31(1), 1-31. doi:10.3109/08977194.2012.752737
- Wang, B., Tang, Z., Gong, H., Zhu, L., & Liu, X. (2017). Wnt5a promotes epithelial-to-mesenchymal transition and metastasis in non-small-cell lung cancer. *Biosci Rep*, 37(6). doi:10.1042/BSR20171092
- Willert, K., & Nusse, R. (2012). Wnt proteins. *Cold Spring Harb Perspect Biol*, 4(9), a007864. doi:10.1101/cshperspect.a007864
- Zeng, H., Lu, B., Zamponi, R., Yang, Z., Wetzel, K., Loureiro, J., . . . Cong, F. (2018). mTORC1 signaling suppresses Wnt/beta-catenin signaling through DVL-dependent regulation of Wnt receptor FZD level. *Proc Natl Acad Sci U S A*, 115(44), E10362-E10369. doi:10.1073/pnas.1808575115
- Zhou, Y., Huang, Y., Cao, X., Xu, J., Zhang, L., Wang, J., . . . Zheng, M. (2016). WNT2 Promotes Cervical Carcinoma Metastasis and Induction of Epithelial-Mesenchymal Transition. *PLoS One*, 11(8), e0160414. doi:10.1371/journal.pone.0160414
- Ziv, E., Yarmohammadi, H., Boas, F. E., Petre, E. N., Brown, K. T., Solomon, S. B., . . . Erinjeri, J. P. (2017). Gene Signature Associated with Upregulation of the Wnt/beta-Catenin Signaling Pathway Predicts Tumor Response to Transarterial Embolization. *J Vasc Interv Radiol*, 28(3), 349-355 e341. doi:10.1016/j.jvir.2016.11.004

Event: 1755: beta-catenin activation

Short Name: beta-catenin activation

Key Event Component

Process	Object	Action
regulation of beta-catenin-TCF complex assembly	beta-catenin-TCF complex	occurrence

AOPs Including This Key Event

AOP ID and Name				Event Type
Aop:298 - Increase in reactive oxygen species (ROS) leading to human treatment-resistant gastric cancer				KeyEvent
Biological Context				
Level of Biological Organization				
Cellular				
Cell term				
Cell term				
cell				
Organ term				
Organ term				
organ				
Domain of Applicability				
Taxonomic Applicability				
Term	Scientific Term	Evidence	Links	
Homo sapiens	Homo sapiens	High	NCBI	
Life Stage Applicability				
Life Stage	Evidence			
All life stages	Moderate			
Sex Applicability				
Sex	Evidence			
Unspecific	High			
Beta-catenin is stabilized and translocated into nucleus in <i>Homo sapiens</i> (Huang et al., 2019).				
Beta-catenin is activated in <i>Homo sapiens</i> (Huang et al., 2019) (Naujok et al., 2014).				
Key Event Description				
Upon the Wnt signaling activation, beta-catenin is stabilized and activated via inhibition of the phosphorylation by GSK3beta (Huang et al., 2019). Once the beta-catenin is stabilized, it translocates into the nucleus and enhances the expression of target genes of Wnt/beta-catenin signaling pathway (Huang et al., 2019). Beta-catenin activation is related to cancer (Tanabe, 2014).				
Dishevelled (DVL), a positive regulator of Wnt signaling, forms the complex with FZD and leads to trigger the Wnt signaling together with Wnt coreceptor low-density lipoprotein (LDL) receptor-related protein 6 (LRP6) (Clevers & Nusse, 2012; Jiang, et al., 2015). DVL, however, has a controversial role to promote Wnt receptor degradation (Jiang et al., 2015). Meanwhile, DVL-dependent regulation of FZD level is involved in mTORC1 signaling suppression via Wnt/beta-catenin signaling (Zeng et al., 2018). The recruitment of Axin to the DVL-FZD complex induces the beta-catenin stabilization and activation. The stabilized beta-catenin translocates into the nucleus, which forms the complex with TCF to induce the up-regulated expression of proliferation-related genes.				
How it is Measured or Detected				
The beta-catenin level in nucleus is measured by immunoblotting with anti-beta-catenin antibody(Huang et al., 2019).				
The beta-catenin nuclear translocation is measured by immunofluorescence assay(Huang et al., 2019).				
Activity of beta-catenin is measured by Wnt/beta-catenin activity assay, in which the vector containing the firefly luciferase gene controlled by TCF/LEF binding sites is transfected in the cells (Naujok et al., 2014).				
References				
Clevers, H., & Nusse, R. (2012). Wnt/beta-catenin signaling and disease. <i>Cell</i> , 145(6), 1192-1205. doi:10.1016/j.cell.2012.05.012				
Huang, J. Q., Wei, F. K., Xu, X. L., Ye, S. X., Song, J. W., Ding, P. K., . . . Gong, L. Y. (2019). SOX9 drives the epithelial-mesenchymal transition in non-small-cell lung cancer through the Wnt/beta-catenin pathway. <i>J Transl Med</i> , 17(1), 143. doi:10.1186/s12967-019-1895-2				
Jiang, X., Charlat, O., Zamponi, R., Yang, Y., & Cong, F. (2015). Dishevelled promotes Wnt receptor degradation through recruitment of ZNRF3/RNF43 E3 ubiquitin ligases. <i>Mol Cell</i> , 58(3), 522-533. doi:10.1016/j.molcel.2015.03.015				
Naujok, O., Lentjes, J., Diekmann, U., Davenport, C., & Lenzen, S. (2014). Cytotoxicity and activation of the Wnt/beta-catenin pathway in mouse embryonic stem cells treated with four GSK3 inhibitors. <i>BMC Res Notes</i> , 7, 273. doi:10.1186/1756-0500-7-273				
Tanabe, S. (2014). Role of mesenchymal stem cells in cell life and their signaling. <i>World journal of stem cells</i> , 6(1), 24-32. doi:10.4252/wjsc.v6.i1.24				

Zeng, H., Lu, B., Zamponi, R., Yang, Z., Wetzel, K., Loureiro, J., . . . Cong, F. (2018). mTORC1 signaling suppresses Wnt/beta-catenin signaling through DVL-dependent regulation of Wnt receptor FZD level. *Proc Natl Acad Sci U S A*, *115*(44), E10362-E10369. doi:10.1073/pnas.1808575115

Event: 1457: Epithelial Mesenchymal Transition

Short Name: EMT

Key Event Component

Process	Object	Action
epithelial to mesenchymal transition	Epithelial cell	occurrence

AOPs Including This Key Event

AOP ID and Name	Event Type
Aop:241 - Latent Transforming Growth Factor beta1 activation leads to pulmonary fibrosis	KeyEvent
Aop:206 - Peroxisome proliferator-activated receptors γ inactivation leading to lung fibrosis	KeyEvent
Aop:280 - α-diketone-induced bronchiolitis obliterans	KeyEvent
Aop:347 - Toll-like receptor 4 activation and peroxisome proliferator-activated receptor gamma inactivation leading to pulmonary fibrosis	KeyEvent
Aop:414 - Aryl hydrocarbon receptor activation leading to lung fibrosis through TGF-β dependent fibrosis toxicity pathway	KeyEvent
Aop:415 - Aryl hydrocarbon receptor activation leading to lung fibrosis through IL-6 toxicity pathway	KeyEvent
Aop:443 - DNA damage and mutations leading to Metastatic Breast Cancer	KeyEvent
Aop:298 - Increase in reactive oxygen species (ROS) leading to human treatment-resistant gastric cancer	KeyEvent
Aop:452 - Adverse outcome pathway of PM-induced respiratory toxicity	KeyEvent
Aop:474 - Succinate dehydrogenase inactivation leads to cancer by promoting EMT	KeyEvent

Biological Context

Level of Biological Organization

Cellular

Cell term

Cell term
epithelial cell

Organ term

Organ term
organ

Domain of Applicability

Taxonomic Applicability

Term	Scientific Term	Evidence	Links
humans	Homo sapiens	High	NCBI

Life Stage Applicability

Life Stage	Evidence
Not Otherwise Specified	Not Specified

Sex Applicability

Sex	Evidence
Unspecific	Not Specified

The key event is applicable in *Homo sapiens*:

- Wnt5a expression leads to epithelial-mesenchymal transition (EMT) and metastasis in non-small-cell lung cancer in *Homo sapiens* (Wang et al., 2017).
- WNT2 expression lead to EMT induction in *Homo sapiens* (Zhou et al., 2016).
- EMT is induced in cancer and involved in cancer metastasis in *Homo sapiens* (Suarez-Carmona, Lesage, Cataldo, & Gilles, 2017) (Du & Shim, 2016).

Regulation of miRNA expression by DNA replication, damage and repair responses, transcription and translation has been proved in animals like mice, canine and cell line experiments.

Key Event Description

Epithelial-mesenchymal transition (EMT) is a phenomenon in which the cells transit from epithelial-like into mesenchymal-like phenotypes (Huan et al., 2022; Tanabe, 2017; Tanabe et al., 2015). In cancer, cells exhibiting EMT features contribute to metastasis and drug resistance.

It is known that D-2-hydroxyglutrate induces EMT (Guerra et al., 2017; Jia et al., 2018; Mishra et al., 2018; Sciacovelli & Frezza, 2017). D-2-hydroxyglutrate, an inhibitor of Jumonji-family histone demethylase, increased the trimethylation of histone H3 lysine 4 (H3K4) in the promoter region of the zinc finger E-box-binding homeobox 1 (ZEB1), followed by the induction of EMT (Colvin et al., 2016).

Wnt5a induces EMT and metastasis in non-small-cell lung cancer (Wang et al., 2017).

EMT is related to Wnt/beta-catenin signaling and is important for treatment-resistant cancer (Tanabe et al., 2016).

TGFbeta induces EMT (Wendt et al., 2010).

ZEB is one of the critical transcription factors for EMT regulation (Zhang et al., 2015).

SNAIL (Snail) is an important transcription factor for cell differentiation and survival. The phosphorylation and nuclear localization of Snail1 induced by Wnt signaling pathways are critical for the regulation of EMT (Kaufhold & Bonavida, 2014).

Transcription factors SNAIL and TWIST1 induce EMT (Hodge et al., 2018) (Mani et al., 2008).

It is suggested that Sp1, a transcription factor involved in cell growth and metastasis, is induced by cytochrome P450 1B1 (CYP1B1), and promotes EMT, which leads to cell proliferation and metastasis (Kwon et al., 2016).

Biological state

An epithelial-mesenchymal transition (EMT) is a biologic process in which epithelial cells are polarized, interact through their basal surface with basement membrane, and undergo biochemical changes to assume a mesenchymal cell phenotype.

This phenotypic transformation has various characters such as enhanced migratory capacity, high invasiveness, elevated resistance to apoptosis, and greatly increased production of ECM components (Kalluri, R., and Neilson, E.G. 2003). The completion of an EMT is signalled by the degradation of the underlying basement membrane and the formation of a mesenchymal cell that can migrate away from the epithelial layer in which it originated.

EMT has a number of distinct molecular processes like activation of transcription factors, expression of specific cell surface proteins, reorganization and expression of cytoskeletal proteins, production of ECM-degrading enzymes, and changes in the expression of specific microRNAs. These factors are used as biomarkers to demonstrate the passage of a cell through an EMT.

Biological compartment

Cellular

Role in General Biology:

Excessive proliferation of epithelial cells and angiogenesis mark the initiation and early growth of primary epithelial cancers. (Hanahan, D., and Weinberg, R.A. 2000). The subsequent acquisition of invasiveness, initially manifest by invasion through the basement membrane, is thought to herald the onset of the last stages of the multi-step process that leads eventually to metastatic dissemination, with life-threatening consequences. There has been an intense research going on in the genetic controls and biochemical mechanisms underlying the acquisition of the invasive phenotype and the subsequent systemic spread of the cancer cell. Activation of an EMT program has been proposed as the critical mechanism for the acquisition of malignant phenotypes by epithelial cancer cells (Thiery, J.P. 2002).

Pre-clinical experiments such as mice models and cell culture experiments has demonstrated that carcinoma cells can acquire a mesenchymal phenotype and express mesenchymal markers such as α -SMA, FSP1, vimentin, and desmin (Yang, J., and Weinberg, R.A. 2008). These cells are seen at the invasive front of primary tumors and are considered to be the cells that eventually enter into subsequent steps of the invasion-metastasis cascade, i.e., intravasation, transport through the circulation, extravasation, formation of micro metastases, and ultimately colonization (the growth of small colonies into macroscopic metastases) (Thiery, J.P. 2002, Fidler, I.J., and Poste, G. 2008, Brabletz, T., et al. 2001).

An apparent paradox comes from the observation that the EMT-derived migratory cancer cells typically establish secondary colonies at distant sites that resemble, at the histopathological level, the primary tumor from which they arose; accordingly, they no longer exhibit the mesenchymal phenotypes ascribed to metastasizing carcinoma cells. Reconciling this behaviour with the proposed role of EMT as a facilitator of metastatic dissemination requires the additional notion that metastasizing cancer cells must shed their mesenchymal phenotype via a MET during the course of secondary tumor formation (Zeisberg, M et al 2005). The tendency of disseminated cancer cells to undergo EMT likely reflects the local microenvironments that they encounter after extravasation into the parenchyma of a distant organ, quite possibly the absence of the heterotypic signals they experienced in the primary tumor that were responsible for inducing the EMT in the first place (Thiery, J.P. 2002, Jechlinger, M et al 2002, Bissell, M.J et

al 2002). These evidences indicate that induction of an EMT is likely to be a centrally important mechanism for the progression of carcinomas to a metastatic stage and implicates MET during the subsequent colonization process. However, many steps of this mechanistic model still require direct experimental validation. It remains unclear at present whether these phenomena and molecular mechanisms relate to and explain the metastatic dissemination of non-epithelial cancer cells.

The entire spectrum of signaling agents that contribute to EMTs of carcinoma cells remains unclear. One theory suggests that the genetic and epigenetic alterations undergone by cancer cells during the course of primary tumor formation render them especially responsive to EMT-inducing heterotypic signals originating in the tumor-associated stroma. Oncogenes induce senescence, and recent studies suggest that cancer cell EMTs may also play a role in preventing senescence induced by oncogenes, thereby facilitating subsequent aggressive dissemination (Smit, M.A., and Peeper, D.S. 2008, Ansieau, S., et al. 2008, Weinberg, R.A. 2008). In the case of many carcinomas, EMT-inducing signals emanating from the tumor-associated stroma, notably HGF, EGF, PDGF, and TGF- β , appear to be responsible for the induction or functional activation in cancer cells of a series of EMT-inducing transcription factors, notably Snail, Slug, zinc finger E-box binding homeobox 1 (ZEB1), Twist, Goosecoid, and FOXC2 (Thiery, J.P. 2002, Jechlinger, M. et al 2002, Shi, Y., and Massague, J. 2003, Niessen, K., et al. 2008, Medici, D et al 2008, Kokudo, T., et al. 2008). Once expressed and activated, each of these transcription factors can act pleiotropically to choreograph the complex EMT program, more often than not with the help of other members of this cohort of transcription factors. The actual implementation by these cells of their EMT program depends on a series of intracellular signaling networks involving, among other signal-transducing proteins, ERK, MAPK, PI3K, Akt, Smads, RhoB, β -catenin, lymphoid enhancer binding factor (LEF), Ras, and c-Fos as well as cell surface proteins such as β 4 integrins, α 5 β 1 integrin, and α V β 6 integrin (Tse, J.C., and Kalluri, R. 2007). Activation of EMT programs is also facilitated by the disruption of cell-cell adherens junctions and the cell-ECM adhesions mediated by integrins (Yang, J., and Weinberg, R.A. 2008, Weinberg, R.A. 2008, Gupta, P.B. et al 2005, Yang, J. et al 2006, Mani, S.A., et al. 2007, Mani, S.A., et al. 2008, Hartwell, K.A., et al. 2006, Taki, M et al 2006)..

How it is Measured or Detected

Loss of [E-cadherin](#) and cell polarity is considered to be a fundamental event in epithelial-mesenchymal transition. The simultaneous expression of epithelial (e.g. E-cadherin) and mesenchymal markers (e.g. N-cadherin and vimentin) within the airway epithelium are indicative for ongoing transition (Borthwick et al. 2009, 2010).

	Method/ measurement referenc	Reliability	Strength of evidence	Assay fit for purpose	Repeatability/ reproducibility	Direct measure
Human cell line	qRT-PCR, cell viability assay, Western blotting, EdU incorporation assay	+	Strong	Yes	Yes	Yes
Human	IHC, micro array, qPCR, SNP array	+	Moderate	Yes	Yes	Yes

- EMT can be detected by immunostaining with pro-surfactant protein-C (pro-SPC) and N-cadherin in idiopathic pulmonary fibrosis (IPF) lung *in vivo* (Kim et al., 2006).
- EMT can be detected by immunostaining with vimentin in lung alveola *in vivo* (Kim et al., 2006).
- EMT can be detected as the increased level of the transcription factors, zinc finger E-box-binding homeobox (ZEB), Twist and Snail (Huang et al., 2022).

References

- Borthwick, L. A., Parker, S. M., Brougham, K. A., Johnson, G. E., Gorowiec, M. R., Ward, C., ... Fisher, A. J. (2009). Epithelial to mesenchymal transition (EMT) and airway remodelling after human lung transplantation. *Thorax*, 64(9), 770–777. <https://doi.org/10.1136/thx.2008.104133>
- Borthwick, L. A., McIlroy, E. I., Gorowiec, M. R., Brodlie, M., Johnson, G. E., Ward, C., ... Fisher, A. J. (2010). Inflammation and epithelial to mesenchymal transition in lung transplant recipients: Role in dysregulated epithelial wound repair. *American Journal of Transplantation*, 10(3), 498–509. <https://doi.org/10.1111/j.1600-6143.2009.02953.x>
- Al Saleh, S., Al Mulla, F., & Luqmani, Y. A. (2011). Estrogen receptor silencing induces epithelial to mesenchymal transition in human breast cancer cells. *PloS one*, 6(6), e20610.
- Bissell, M. J., Radisky, D. C., Rizki, A., Weaver, V. M., & Petersen, O. W. (2002). The organizing principle: microenvironmental influences in the normal and malignant breast. *Differentiation*, 70(9-10), 537–546.
- Bouris, P., Skandalis, S. S., Piperigkou, Z., Afratis, N., Karamanou, K., Aletras, A. J., ... & Karamanos, N. K. (2015). Estrogen receptor alpha mediates epithelial to mesenchymal transition, expression of specific matrix effectors and functional properties of breast cancer cells. *Matrix Biology*, 43, 42–60.
- Brabletz, T., Jung, A., Reu, S., Porzner, M., Hlubek, F., Kunz-Schughart, L. A., ... & Kirchner, T. (2001). Variable β -catenin expression in colorectal

- cancers indicates tumor progression driven by the tumor environment. *Proceedings of the National Academy of Sciences*, 98(18), 10356-10361.
- Brabletz, T., Jung, A., Reu, S., Porzner, M., Hlubek, F., Kunz-Schughart, L. A., ... & Kirchner, T. (2001). Variable β -catenin expression in colorectal cancers indicates tumor progression driven by the tumor environment. *Proceedings of the National Academy of Sciences*, 98(18), 10356-10361.
- Colvin, H., Nishida, N., Konno, M., Haraguchi, N., Takahashi, H., Nishimura, J., . . . Ishii, H. (2016). Oncometabolite D-2-Hydroxyglurate Directly Induces Epithelial-Mesenchymal Transition and is Associated with Distant Metastasis in Colorectal Cancer. *Sci Rep*, 6, 36289. doi:10.1038/srep36289
- Du, B., & Shim, J. S. (2016). Targeting Epithelial-Mesenchymal Transition (EMT) to Overcome Drug Resistance in Cancer. *Molecules*, 21(7). doi:10.3390/molecules21070965
- Fang, C. X., Ma, C. M., Jiang, L., Wang, X. M., Zhang, N., Ma, J. N., . . . Zhao, Y. D. (2018). p38 MAPK is Crucial for Wnt1- and LiCl-Induced Epithelial Mesenchymal Transition. *Curr Med Sci*, 38(3), 473-481. doi:10.1007/s11596-018-1903-4
- Fidler, I. J., & Poste, G. (2008). The “seed and soil” hypothesis revisited. *The lancet oncology*, 9(8), 808.
- Guerra, F., Guaragnella, N., Arbini, A. A., Bucci, C., Giannattasio, S., & Moro, L. (2017). Mitochondrial Dysfunction: A Novel Potential Driver of Epithelial-to-Mesenchymal Transition in Cancer. *Front Oncol*, 7, 295. doi:10.3389/fonc.2017.00295
- Gupta, P. B., Mani, S., Yang, J., Hartwell, K., & Weinberg, R. A. (2005, January). The evolving portrait of cancer metastasis. In *Cold Spring Harbor symposia on quantitative biology* (Vol. 70, pp. 291-297). Cold Spring Harbor Laboratory Press.
- Hanahan, D., and Weinberg, R.A. (2000). The hall- marks of cancer. *Cell*. 100:57-70.
- Hartwell, K. A., Muir, B., Reinhardt, F., Carpenter, A. E., Sgroi, D. C., & Weinberg, R. A. (2006). The Spemann organizer gene, Goosecoid, promotes tumor metastasis. *Proceedings of the National Academy of Sciences*, 103(50), 18969-18974.
- Jechlinger, M., Grünert, S., & Beug, H. (2002). Mechanisms in epithelial plasticity and metastasis: insights from 3D cultures and expression profiling. *Journal of mammary gland biology and neoplasia*, 7(4), 415-432.
- Hodge, D. Q., Cui, J., Gamble, M. J., & Guo, W. (2018). Histone Variant MacroH2A1 Plays an Isoform-Specific Role in Suppressing Epithelial-Mesenchymal Transition. *Sci Rep*, 8(1), 841. doi:10.1038/s41598-018-19364-4
- Huan, Z., Zhang, Z., Zhou, C., Liu, L., Huang, C. (2022). Epithelial-mesenchymal transition: The history, regulatory mechanism, and cancer therapeutic opportunities. *MedComm*. 2022 May 18;3(2):e144. doi: 10.1002/mco2.144
- Jia, D., Park, J. H., Jung, K. H., Levine, H., & Kaiparettu, B. A. (2018). Elucidating the Metabolic Plasticity of Cancer: Mitochondrial Reprogramming and Hybrid Metabolic States. *Cells*, 7(3). doi:10.3390/cells7030021
- Kalluri, R., & Neilson, E. G. (2003). Epithelial-mesenchymal transition and its implications for fibrosis. *The Journal of clinical investigation*, 112(12), 1776-1784.
- Kaufhold, S., & Bonavida, B. (2014). Central role of Snail1 in the regulation of EMT and resistance in cancer: a target for therapeutic intervention. *J Exp Clin Cancer Res*, 33, 62. doi:10.1186/s13046-014-0062-0
- Kim, K. K., Kugler, M. C., Wolters, P. J., Robillard, L., Galvez, M. G., Brumwell, A. N., . . . Chapman, H. A. (2006). Alveolar epithelial cell mesenchymal transition develops in vivo during pulmonary fibrosis and is regulated by the extracellular matrix. *PNAS*, 103(35), 13180-13185. doi:10.1073/pnas.0605669103
- Kwon, Y. J., Baek, H. S., Ye, D. J., Shin, S., Kim, D., & Chun, Y. J. (2016). CYP1B1 Enhances Cell Proliferation and Metastasis through Induction of EMT and Activation of Wnt/beta-Catenin Signaling via Sp1 Upregulation. *PLoS One*, 11(3), e0151598. doi:10.1371/journal.pone.0151598
- Kokudo, T., Suzuki, Y., Yoshimatsu, Y., Yamazaki, T., Watabe, T., & Miyazono, K. (2008). Snail is required for TGF β -induced endothelial-mesenchymal transition of embryonic stem cell-derived endothelial cells. *Journal of cell science*, 121(20), 3317-3324.
- Lin, H. Y., Liang, Y. K., Dou, X. W., Chen, C. F., Wei, X. L., Zeng, D., ... & Zhang, G. J. (2018). Notch3 inhibits epithelial-mesenchymal transition in breast cancer via a novel mechanism, upregulation of GATA-3 expression. *Oncogenesis*, 7(8), 1-15.
- Liu, Y., Liu, R., Fu, P., Du, F., Hong, Y., Yao, M., ... & Zheng, S. (2015). N1-Guanyl-1, 7-diaminoheptane sensitizes estrogen receptor negative breast cancer cells to doxorubicin by preventing epithelial-mesenchymal transition through inhibition of eukaryotic translation initiation factor 5A2 activation. *Cellular Physiology and Biochemistry*, 36(6), 2494-2503.
- Mani, S. A., Yang, J., Brooks, M., Schwaninger, G., Zhou, A., Miura, N., ... & Weinberg, R. A. (2007). Mesenchyme Forkhead 1 (FOXC2) plays a key role in metastasis and is associated with aggressive basal-like breast cancers. *Proceedings of the National Academy of Sciences*, 104(24), 10069-10074.
- Mani, S. A., Guo, W., Liao, M. J., Eaton, E. N., Ayyanan, A., Zhou, A. Y., ... & Weinberg, R. A. (2008). The epithelial-mesenchymal transition generates cells with properties of stem cells. *Cell*, 133(4), 704-715.
- Medici, D., Hay, E. D., & Olsen, B. R. (2008). Snail and Slug promote epithelial-mesenchymal transition through β -catenin-T-cell factor-4-dependent expression of transforming growth factor- β 3. *Molecular biology of the cell*, 19(11), 4875-4887.
- Mishra, P., Tang, W., Putluri, V., Dorsey, T. H., Jin, F., Wang, F., . . . Ambis, S. (2018). ADHFE1 is a breast cancer oncogene and induces metabolic reprogramming. *J Clin Invest*, 128(1), 323-340. doi:10.1172/JCI93815
- Niessen, K., Fu, Y., Chang, L., Hoodless, P. A., McFadden, D., & Karsan, A. (2008). Slug is a direct Notch target required for initiation of cardiac cushion cellularization. *The Journal of cell biology*, 182(2), 315-325.
- Sciacovelli, M., & Frezza, C. (2017). Metabolic reprogramming and epithelial-to-mesenchymal transition in cancer. *FEBS J*, 284(19), 3132-3144. doi:10.1111/febs.14090
- Shi, Y., & Massagué, J. (2003). Mechanisms of TGF- β signaling from cell membrane to the nucleus. *cell*, 113(6), 685-700.
- Smit, M. A., & Peeper, D. S. (2008). Deregulating EMT and senescence: double impact by a single twist. *Cancer cell*, 14(1), 5-7.
- Suarez-Carmona, M., Lesage, J., Cataldo, D., & Gilles, C. (2017). EMT and inflammation: inseparable actors of cancer progression. *Mol Oncol*, 11(7), 805-823. doi:10.1002/1878-0261.12095
- Sun, J., Yang, X., Zhang, R., Liu, S., Gan, X., Xi, X., . . . Sun, Y. (2017). GOLPH3 induces epithelial-mesenchymal transition via Wnt/beta-catenin signaling pathway in epithelial ovarian cancer. *Cancer Med*, 6(4), 834-844. doi:10.1002/cam4.1040
- Taki, M., Verschuere, K., Yokoyama, K., Nagayama, M., & Kamata, N. (2006). Involvement of Ets-1 transcription factor in inducing matrix metalloproteinase-2 expression by epithelial-mesenchymal transition in human squamous carcinoma cells. *International journal of oncology*, 28(2), 487-496.

Tanabe, S. (2017). Molecular markers and networks for cancer and stem cells. *J Embryol Stem Cell Res*, 1(1).

Tanabe, S., Kawabata, T., Aoyagi, K., Yokozaki, H., & Sasaki, H. (2016). Gene expression and pathway analysis of CTNNB1 in cancer and stem cells. *World J Stem Cells*, 8(11), 384-395. doi:10.4252/wjsc.v8.i11.384

Tanabe, S., Komatsu, M., Kazuhiko, A., Yokozaki, H., & Sasaki, H. (2015). Implications of epithelial-mesenchymal transition in gastric cancer. *Translational Gastrointestinal Cancer*, 4(4), 258-264

Thiery, J. P. (2002). Epithelial-mesenchymal transitions in tumour progression. *Nature reviews cancer*, 2(6), 442-454.

Tse, J. C., & Kalluri, R. (2007). Mechanisms of metastasis: epithelial-to-mesenchymal transition and contribution of tumor microenvironment. *Journal of cellular biochemistry*, 101(4), 816-829.

Wang, B., Tang, Z., Gong, H., Zhu, L., & Liu, X. (2017). Wnt5a promotes epithelial-to-mesenchymal transition and metastasis in non-small-cell lung cancer. *Biosci Rep*, 37(6). doi:10.1042/BSR20171092

Weinberg, R. A. (2008). Twisted epithelial-mesenchymal transition blocks senescence. *Nature cell biology*, 10(9), 1021-1023.

Wendt, M. K., Smith, J. A., & Schiemann, W. P. (2010). Transforming growth factor-beta-induced epithelial-mesenchymal transition facilitates epidermal growth factor-dependent breast cancer progression. *Oncogene*, 29(49), 6485-6498. doi:10.1038/onc.2010.377

Wik, E., Ræder, M. B., Krakstad, C., Trovik, J., Birkeland, E., Hoivik, E. A., ... & Salvesen, H. B. (2013). Lack of estrogen receptor- α is associated with epithelial-mesenchymal transition and PI3K alterations in endometrial carcinoma. *Clinical Cancer Research*, 19(5), 1094-1105.

Yang, J., & Weinberg, R. A. (2008). Epithelial-mesenchymal transition: at the crossroads of development and tumor metastasis. *Developmental cell*, 14(6), 818-829.

Yang, J., Mani, S. A., & Weinberg, R. A. (2006). Exploring a new twist on tumor metastasis. *Cancer research*, 66(9), 4549-4552.

Ye, Y., Xiao, Y., Wang, W., Yearsley, K., Gao, J. X., Shetuni, B., & Barsky, S. H. (2010). ER α signaling through slug regulates E-cadherin and EMT. *Oncogene*, 29(10), 1451-1462.

Zeisberg, M., Shah, A. A., & Kalluri, R. (2005). Bone morphogenic protein-7 induces mesenchymal to epithelial transition in adult renal fibroblasts and facilitates regeneration of injured kidney. *Journal of Biological Chemistry*, 280(9), 8094-8100.

Zeng, Q., Zhang, P., Wu, Z., Xue, P., Lu, D., Ye, Z., ... & Yan, X. (2014). Quantitative proteomics reveals ER- α involvement in CD146-induced epithelial-mesenchymal transition in breast cancer cells. *Journal of proteomics*, 103, 153-169.

Zhang, P., Sun, Y., & Ma, L. (2015). ZEB1: at the crossroads of epithelial-mesenchymal transition, metastasis and therapy resistance *Cell Cycle*, 14(4), 481-487. doi:10.1080/15384101.2015.1006048

Zhou, Y., Huang, Y., Cao, X., Xu, J., Zhang, L., Wang, J., ... Zheng, M. (2016). WNT2 Promotes Cervical Carcinoma Metastasis and Induction of Epithelial-Mesenchymal Transition. *PLoS One*, 11(8), e0160414. doi:10.1371/journal.pone.0160414

List of Adverse Outcomes in this AOP

Event: 1651: Treatment-resistant gastric cancer

Short Name: Resistant gastric cancer

Key Event Component

Process	Object	Action
regulation of cellular response to drug		occurrence

AOPs Including This Key Event

AOP ID and Name	Event Type
Aop:298 - Increase in reactive oxygen species (ROS) leading to human treatment-resistant gastric cancer	AdverseOutcome

Biological Context

Level of Biological Organization

Tissue

Organ term

Organ term

organ

Domain of Applicability

Taxonomic Applicability

Term	Scientific Term	Evidence	Links
Homo sapiens	Homo sapiens	High	NCBI

Life Stage Applicability

Life Stage	Evidence
------------	----------

Life Stage Evidence

All life stages High

Sex Applicability

Sex Evidence

Unspecific High

Drug resistance occurs in *Homo sapiens* (Du & Shim, 2016).

Key Event Description

It is known that diffuse-type gastric cancer, which has a poor prognosis, is treatment-resistant and more malignant compared to intestinal-type gastric cancer (Tanabe et al., 2014). Drug resistance is involved in EMT, which is an important phenomenon exhibiting features similar to cancer stem cells (CSCs) (Du & Shim, 2016).

EMT is involved in metastasis and cancer therapy resistance (Smith & Bhowmick, 2016).

How it is Measured or Detected

Treatment-resistant gastric cancer and EMT can be detected with biomarkers (Zeisberg & Neilson, 2009).

Treatment-resistant gastric cancer which exhibits EMT phenotype can be detected as the increased level of the transcription factors, zinc finger E-box-binding homeobox 1/2 (ZEB1/2), SNAIL1/2, and TWIST2 which are associated with the activation of EMT-related genes (Tanabe et al., 2022a and 2022b).

Regulatory Significance of the AO

Drug resistance is very important in cancer treatment since cancer metastasis and recurrence are some of the main obstacles to treating cancer. Cancer stem cells that share the phenotype of EMT may be targeted in anti-cancer drug development.

References

- Du, B., & Shim, J. S. (2016). Targeting Epithelial-Mesenchymal Transition (EMT) to Overcome Drug Resistance in Cancer. *Molecules*, 21(7). doi:10.3390/molecules21070965
- Smith, B. N., & Bhowmick, N. A. (2016). Role of EMT in Metastasis and Therapy Resistance. *J Clin Med*, 5(2). doi:10.3390/jcm5020017
- Tanabe, S., Aoyagi, K., Yokozaki, H., Sasaki, H. (2014). Gene expression signatures for identifying diffuse-type gastric cancer associated with epithelial-mesenchymal transition. *International journal of oncology*, 44(6), 1955-1970. doi:10.3892/ijo.2014.2387
- Tanabe, S., Quader, S., Cabral, H., Ono, R. (2020a). Interplay of EMT and CSC in Cancer and the Potential Therapeutic Strategies. *Front Pharmacol*, 11, 904. doi:10.3389/fphar.2020.00904
- Tanabe S, Quader S, Ono R, Cabral H, Aoyagi K, Hirose A, Yokozaki H., Sasaki, H. (2020b). Molecular Network Profiling in Intestinal- and Diffuse-Type Gastric Cancer. *Cancers (Basel)*, 12(12), 3833. doi:10.3390/cancers12123833
- Zeisberg, M., & Neilson, E. G. (2009). Biomarkers for epithelial-mesenchymal transitions. *J Clin Invest*, 119(6), 1429-1437. doi:10.1172/JCI36183

Appendix 2**List of Key Event Relationships in the AOP****List of Adjacent Key Event Relationships**

Relationship: 3505: Increase, ROS leads to Increase, porcupine-induced Wnt secretion and Wnt signaling activation

AOPs Referencing Relationship

AOP Name	Adjacency	Weight of Evidence	Quantitative Understanding
Increase in reactive oxygen species (ROS) leading to human treatment-resistant gastric cancer	adjacent	Moderate	Moderate

Evidence Supporting Applicability of this Relationship**Taxonomic Applicability**

Term	Scientific Term	Evidence	Links
Homo sapiens	Homo sapiens	Moderate	NCBI

Life Stage Applicability

Life Stage	Evidence
All life stages	Moderate

Life Stage Evidence**Sex Applicability****Sex Evidence**

Unspecific Moderate

Prolonged ROS induces inflammation and tissue damage in *Homo sapiens* (Vallée & Lecarpentier, 2018).

Key Event Relationship Description

ROS production causes tissue damage (Gao, Zhou, Lin, Paus, & Yue, 2019). ROS production is involved in Wnt-driven tumorigenesis (Myant et al., 2013). The prolonged ROS induces inflammation leading to carcinogenesis (Vallée & Lecarpentier, 2018). Injury causes the Porcupine-induced Wnt secretion (Saha et al., 2016).

Evidence Supporting this KER**Biological Plausibility**

Sustained ROS increase caused by/causes DNA damage, which will alter several signaling pathways including Wnt signaling.

Macrophages accumulate into injured tissue to recover the tissue damage, which may be followed by porcupine-induced Wnt secretion. ROS stimulate inflammatory factor production and Wnt/beta-catenin signaling (Vallée & Lecarpentier, 2018).

Empirical Evidence

Production of ROS by DNA double-strand break causes tissue damages (Gao et al., 2019).

ROS signaling induces Wnt/beta-catenin signaling (Pérez, Taléns-Visconti, Rius-Pérez, Finamor, & Sastre, 2017).

Uncertainties and Inconsistencies

The balance of ROS signaling is important, and dual effects of ROS should be taken in consideration. The ROS may enhance Wnt/beta-catenin proliferating pathways to promote tumorigenesis, while ROS may disrupt tumor progression by different pro-apoptotic mechanisms (Pérez et al., 2017). It is also known that Wnt signaling induces ROS signaling (Cheung et al., 2016). Wnt/beta-catenin signaling control by ROS needs to be further investigated (Caliceti, Nigro, Rizzo, & Ferrari, 2014).

Quantitative Understanding of the Linkage**Response-response relationship**

ROS induces inflammatory responses (Bhattacharyya, Chattopadhyay, Mitra, & Crowe, 2014). Oxidant induces ROS generation and p38 MAPK activation in macrophages (Conway & Kinter, 2006). ROS induce tissue damage in cardiac myocytes (Miller & Cheung, 2016; Yang et al., 2006).

Time-scale

For the colony formation assay, cells were treated with 400 microM/L H2O2 for 1 week, where the medium was changed every three days (Wang et al., 2019).

Known modulating factors

GPX2, an activator of Wnt/beta-catenin signaling, is identified as a key regulator of intracellular H2O2 levels and an inhibitor of apoptosis (Wang et al., 2019).

Known Feedforward/Feedback loops influencing this KER

The reduction in ROS levels in the human serum albumin-treated cerebral ischemia/reperfusion-induced injury may be mediated by Wnt/betacatenin signaling (Tang, Shen, Zhang, Yang, & Liu, 2019).

References

- Bhattacharyya, A., Chattopadhyay, R., Mitra, S., & Crowe, S. E. (2014). Oxidative stress: an essential factor in the pathogenesis of gastrointestinal mucosal diseases. *Physiological reviews*, 94(2), 329-354. doi:10.1152/physrev.00040.2012
- Caliceti, C., Nigro, P., Rizzo, P., & Ferrari, R. (2014). ROS, Notch, and Wnt signaling pathways: crosstalk between three major regulators of cardiovascular biology. *BioMed research international*, 2014, 318714-318714. doi:10.1155/2014/318714
- Cheung, E. C., Lee, P., Ceteci, F., Nixon, C., Blyth, K., Sansom, O. J., & Vousden, K. H. (2016). Opposing effects of TIGAR- and RAC1-derived ROS on Wnt driven proliferation in the mouse intestine. *Genes & development*, 30(1), 52-63. doi:10.1101/gad.271130.115
- Conway, J. P., & Kinter, M. (2006). Dual role of peroxiredoxin I in macrophage-derived foam cells. *The Journal of biological chemistry*, 281(38), 27991-28001. doi:10.1074/jbc.M605026200
- Gao, Q., Zhou, G., Lin, S.-J., Paus, R., & Yue, Z. (2019). How chemotherapy and radiotherapy damage the tissue: Comparative biology lessons from feather and hair models. *Experimental dermatology*, 28(4), 413-418. doi:10.1111/exd.13846
- Miller, B. A., & Cheung, J. Y. (2016). TRPM2 protects against tissue damage following oxidative stress and ischaemia-reperfusion. *The Journal of physiology*, 594(15), 4181-4191. doi:10.1113/jp270934
- Myant, K. B., Cammareri, P., McGhee, E. J., Ridgway, R. A., Huels, D. J., Cordero, J. B., . . . Sansom, O. J. (2013). ROS production and NF-κB activation triggered by RAC1 facilitate WNT-driven intestinal stem cell proliferation and colorectal cancer initiation. *Cell stem cell*, 12(6), 761-773. doi:10.1016/j.stem.2013.04.006
- Pérez, S., Taléns-Visconti, R., Rius-Pérez, S., Finamor, I., & Sastre, J. (2017). Redox signaling in the gastrointestinal tract. *Free radical biology & medicine*, 104, 75-103. doi:10.1016/j.freeradbiomed.2016.12.048
- Saha, S., Aranda, E., Hayakawa, Y., Bhanja, P., Atay, S., Brodin, N. P., . . . Pollard, J. W. (2016). Macrophage-derived extracellular vesicle-packaged WNTs rescue intestinal stem cells and enhance survival after radiation injury. *Nature Communications*, 7, 13096-13096. doi:10.1038/ncomms13096

Tang, Y., Shen, J., Zhang, F., Yang, F.-Y., & Liu, M. (2019). Human serum albumin attenuates global cerebral ischemia/reperfusion-induced brain injury in a Wnt/ β -Catenin/ROS signaling-dependent manner in rats. *Biomedicine & pharmacotherapy = Biomedecine & pharmacotherapie*, 115, 108871-108871. doi:10.1016/j.biopha.2019.108871

Vallée, A., & Lecarpentier, Y. (2018). Crosstalk Between Peroxisome Proliferator-Activated Receptor Gamma and the Canonical WNT/ β -Catenin Pathway in Chronic Inflammation and Oxidative Stress During Carcinogenesis. *Frontiers in immunology*, 9, 745-745. doi:10.3389/fimmu.2018.00745

Wang, Y., Cao, P., Alshwmi, M., Jiang, N., Xiao, Z., Jiang, F., . . . Li, S. (2019). GPX2 suppression of H(2)O(2) stress regulates cervical cancer metastasis and apoptosis via activation of the β -catenin-WNT pathway. *OncoTargets and therapy*, 12, 6639-6651. doi:10.2147/OTT.S208781

Yang, K. T., Chang, W. L., Yang, P. C., Chien, C. L., Lai, M. S., Su, M. J., & Wu, M. L. (2006). Activation of the transient receptor potential M2 channel and poly(ADP-ribose) polymerase is involved in oxidative stress-induced cardiomyocyte death. *Cell Death & Differentiation*, 13(10), 1815-1826. doi:10.1038/sj.cdd.4401813

Relationship: 2070: Increase, porcupine-induced Wnt secretion and Wnt signaling activation leads to beta-catenin activation

AOPs Referencing Relationship

AOP Name	Adjacency	Weight of Evidence	Quantitative Understanding
Increase in reactive oxygen species (ROS) leading to human treatment-resistant gastric cancer	adjacent	Moderate	Moderate

Evidence Supporting Applicability of this Relationship

Taxonomic Applicability

Term	Scientific Term	Evidence	Links
Homo sapiens	Homo sapiens	High	NCBI

Life Stage Applicability

Life Stage	Evidence
All life stages	High

Sex Applicability

Sex	Evidence
Unspecific	High

Wnt/beta-catenin signaling, which regulates key cellular functions including proliferation, is a highly conserved pathway through evolution (Pai et al., 2017).

Key Event Relationship Description

Secreted Wnt ligand stimulates Wnt/beta-catenin signaling, in which beta-catenin is activated. Wnt ligand binds to Frizzled receptor, which leads to GSK3 β inactivation. GSK3 β inactivation leads to beta-catenin dephosphorylation, which avoids the ubiquitination of the beta-catenin and stabilizes the beta-catenin (Clevers & Nusse, 2012). The translocation of stabilized beta-catenin induces the transcription of genes involved in proliferation (Pai et al., 2017).

Evidence Supporting this KER

Biological Plausibility

Canonical Wnt pathway consists of Wnt, GSK3 β , and beta-catenin cascade (Clevers & Nusse, 2012; Hatsell, Rowlands, Hiremath, & Cowin, 2003).

GSK3 β recruitment to LRP6 leads to form un-phosphorylated beta-catenin inducing the stabilization and translocation of the beta-catenin (MacDonald, Tamai, & He, 2009).

Stabilized beta-catenin accumulates in cytosol and translocates into the nucleus leading to beta-catenin activation (MacDonald et al., 2009).

Empirical Evidence

[Incidence concordance]

Dishevelled (DVL), a positive regulator of Wnt signaling, form the complex with FZD and lead to trigger the Wnt signaling together with Wnt coreceptor low-density lipoprotein (LDL) receptor-related protein 6 (LRP6) (Clevers & Nusse, 2012; Jiang, Charlat, Zamponi, Yang, & Cong, 2015). Wnt binds to FZD and activates the Wnt signaling (Clevers & Nusse, 2012; Janda, Waghay, Levin, Thomas, & Garcia, 2012; Nile, Mukund, Stanger, Wang, & Hannoush, 2017). Wnt binding towards FZD induces the formation of the protein complex with LRP5/6 and DVL, leading to the downstream signaling activation including beta-catenin (Clevers & Nusse, 2012).

Uncertainties and Inconsistencies

Some Wnt ligands bind to FZD, leading to Wnt/beta-catenin signaling inactivation. DVL, a positive regulator of Wnt signaling, has a controversial role to promote Wnt receptor degradation (Jiang et al., 2015). DVL-dependent regulation of FZD level is involved in mTORC1 signaling suppression via Wnt/beta-catenin signaling (Zeng et al., 2018)

GSK3 β phosphorylates LRP6 as well as remaining GSK3 β phosphorylates beta-catenin which would be ubiquitinated and degraded (MacDonald et al., 2009).

Quantitative Understanding of the Linkage

Response-response relationship

Wnt3 promotes proliferation and survival in HUVECs (Shen et al., 2018).

GSK3beta inhibition by 1 uM of SB216763 or 5 uM of BRD3731 results in the decreased phosphorylation and stabilization of beta-catenin (Stump et al., 2019). The level of beta-catenin is increased by the inhibition of GSK3beta kinase activity (Stump et al., 2019). GSK3beta inhibition by small interference RNA (siRNA) of GSK3beta results in the decreased phosphorylation and increased expression of beta-catenin (Stump et al., 2019).

Time-scale

FZD7 enhances the activity of canonical Wnt/beta-catenin signaling with the treatment of WNT3A for 1 to 6 hr (Cao et al., 2017). The treatment with SB216763 or BRD3731, GSK3beta inhibitors, decreases phosphorylated beta-catenin and increased beta-catenin expression in 48 hours (Stump et al., 2019). The cells are treated with GSK3beta small interference RNA (siRNA) for 48 hours to silence the expression of GSK3beta, which results in the activation of beta-catenin pathway (Stump et al., 2019).

Known modulating factors

FZD5 can activate WNT3A/beta-catenin signaling in a dose-dependent manner (Hua et al., 2018). The increase in FZD5 protein enhances cell response to WNT3A. (Hua et al., 2018). LRP5 can augment WNT3A/beta-catenin signaling in a dose-dependent manner (Hua et al., 2018). The binding of Wnt and FZD induce the formation of the protein complex with the Dvl, Axin, CK1 GSK3, beta-catenin and APC to induce the beta-catenin translocation into the nucleus (Clevers & Nusse, 2012).

Known Feedforward/Feedback loops influencing this KER

Beta-catenin is required and sufficient for the sequestration of GSK3 in acidic cytoplasmic endosomes (Taelman et al., 2010). Beta-catenin, of which level increases in Wnt signaling, facilitates GSK3 sequestration leading to feed-forward loop formation (Taelman et al., 2010). The Wnt ligand is antagonized with secreted Frizzled-related proteins (sFRPs) and Wnt inhibitory protein (WIF), both of which can bind Wnts and inhibit interactions between WNT and FZD (Bovolenta, Esteve, Ruiz, Cisneros, & Lopez-Rios, 2008; Clevers & Nusse, 2012). The Dickkopf 1 (DKK1) can disrupts Wnt-induced FZD-LRP6 complex formation (Clevers & Nusse, 2012; Ellwanger et al., 2008; Semenov, Zhang, & He, 2008).

References

- Bovolenta, P., Esteve, P., Ruiz, J. M., Cisneros, E., & Lopez-Rios, J. (2008). Beyond Wnt inhibition: new functions of secreted Frizzled-related proteins in development and disease. *Cell Sci*, 121(Pt 6), 737-746. doi:10.1242/jcs.026096
- Cao, T. T., Xiang, D., Liu, B. L., Huang, T. X., Tan, B. B., Zeng, C. M., . . . Fu, L. (2017). FZD7 is a novel prognostic marker and promotes tumor metastasis via WNT and EMT signaling pathways in esophageal squamous cell carcinoma. *Oncotarget*, 8(39), 65957-65968. doi:10.18632/oncotarget.19586
- Clevers, H., & Nusse, R. (2012). Wnt/beta-catenin signaling and disease. *Cell*, 149(6), 1192-1205. doi:10.1016/j.cell.2012.05.012
- Ellwanger, K., Saito, H., Clement-Lacroix, P., Maltry, N., Niedermeyer, J., Lee, W. K., . . . Niehrs, C. (2008). Targeted disruption of the Wnt regulator Kremen induces limb defects and high bone density. *Mol Cell Biol*, 28(15), 4875-4882. doi:10.1128/MCB.00222-08
- Hatsell, S., Rowlands, T., Hiremath, M., & Cowin, P. (2003). Beta-catenin and Tcfs in mammary development and cancer. *Mammary Gland Biol Neoplasia*, 8(2), 145-158. Retrieved from <https://www.ncbi.nlm.nih.gov/pubmed/14635791>
- Hua, Y., Yang, Y., Li, Q., He, X., Zhu, W., Wang, J., & Gan, X. (2018). Oligomerization of Frizzled and LRP5/6 protein initiates intracellular signaling for the canonical WNT/beta-catenin pathway. *J Biol Chem*, 293(51), 19710-19724. doi:10.1074/jbc.RA118.004434
- Janda, C. Y., Waghray, D., Levin, A. M., Thomas, C., & Garcia, K. C. (2012). Structural basis of Wnt recognition by Frizzled. *Science*, 337(6090), 59-64. doi:10.1126/science.1222879
- Jiang, X., Charlat, O., Zamponi, R., Yang, Y., & Cong, F. (2015). Dishevelled promotes Wnt receptor degradation through recruitment of ZNRF3/RNF43 E3 ubiquitin ligases. *Mol Cell*, 58(3), 522-533. doi:10.1016/j.molcel.2015.03.015
- MacDonald, B. T., Tamai, K., & He, X. (2009). Wnt/beta-catenin signaling: components, mechanisms, and diseases. *Dev Cell*, 17(1), 9-26. doi:10.1016/j.devcel.2009.06.016
- Nile, A. H., Mukund, S., Stanger, K., Wang, W., & Hannoush, R. N. (2017). Unsaturated fatty acyl recognition by Frizzled receptors mediates dimerization upon Wnt ligand binding. *Proc Natl Acad Sci U S A*, 114(16), 4147-4152. doi:10.1073/pnas.1618293114
- Pai SG, Carneiro BA, Mota JM, Costa R, Leite CA, Barroso-Sousa R, Kaplan JB, Chae YK, Giles FJ. Wnt/beta-catenin pathway: modulating anticancer immune response. *J Hematol Oncol*. 2017 May 5;10(1):101. doi: 10.1186/s13045-017-0471-6. PMID: 28476164; PMCID: PMC5420131.
- Semenov, M. V., Zhang, X., & He, X. (2008). DKK1 antagonizes Wnt signaling without promotion of LRP6 internalization and degradation. *J Biol Chem*, 283(31), 21427-21432. doi:10.1074/jbc.M800014200
- Shen, M., Bai, D., Liu, B., Lu, X., Hou, R., Zeng, C., . . . Yin, T. (2018). Dysregulated Txnip-ROS-Wnt axis contributes to the impaired ischemic heart repair in diabetic mice. *Biochimica et biophysica acta. Molecular basis of disease*, 1864(12), 3735-3745. doi:10.1016/j.bbdis.2018.09.029
- Stump, B., Shrestha, S., Lamattina, A. M., Louis, P. H., Cho, W., Perrella, M. A., . . . El-Chemaly, S. (2019). Glycogen synthase kinase 3-beta inhibition induces lymphangiogenesis through beta-catenin-dependent and mTOR-independent pathways. *PLoS One*, 14(4), e0213831. doi:10.1371/journal.pone.0213831
- Taelman, V. F., Dobrowolski, R., Plouhinec, J. L., Fuentealba, L. C., Vorwald, P. P., Gumper, I., . . . De Robertis, E. M. (2010). Wnt signaling requires sequestration of glycogen synthase kinase 3 inside multivesicular endosomes. *Cell*, 143(7), 1136-1148. doi:10.1016/j.cell.2010.11.034
- Zeng, H., Lu, B., Zamponi, R., Yang, Z., Wetzel, K., Loureiro, J., . . . Cong, F. (2018). mTORC1 signaling suppresses Wnt/beta-catenin signaling through DVL-dependent regulation of Wnt receptor FZD level. *Proc Natl Acad Sci U S A*, 115(44), E10362-E10369. doi:10.1073/pnas.1808575115

Relationship: 3199: beta-catenin activation leads to EMT

AOPs Referencing Relationship

AOP Name	Adjacency	Weight of Evidence	Quantitative Understanding
Increase in reactive oxygen species (ROS) leading to human treatment-resistant gastric cancer	adjacent	Moderate	Moderate

Evidence Supporting Applicability of this Relationship

Taxonomic Applicability

Term	Scientific Term	Evidence	Links
Homo sapiens	Homo sapiens	High	NCBI

Life Stage Applicability

Life Stage	Evidence
------------	----------

Life Stage Evidence	
All life stages	High
Sex Applicability	
Sex	Evidence
Unspecific	High
- The inhibition of c-MET decreases the expression of beta-catenin and Snail in human diffuse-type gastric cancer (<i>Homo sapiens</i>) (Sohn et al., 2019). - The treatment with garcinol decreases the expression of beta-catenin and ZEB1/ZEB2 in human breast cancer cells (<i>Homo sapiens</i>) (Ahmad et al., 2012). - Zeb1 activation leads to EMT via Prex1 activation in NCH421k, NCH441, and NCH644 human glioblastoma model cells (<i>Homo sapiens</i>) (Rosmaninho et al., 2018). - Zeb1 siRNA induced the suppression of EMT in SGC-7901 human gastric cancer cell line (<i>Homo sapiens</i>) (Xue et al., 2019). Snail induces EMT in SAS and HSC-4 human head and neck squamous cancer cells (<i>Homo sapiens</i>) (Ota et al., 2016). - Snail induces EMT in B16-F10 murine melanoma cells (<i>Mus musculus</i>) (Kudo-Saito, Shirako, Takeuchi, & Kawakami, 2009; Wang, Shi, Chai, Ying, & Zhou, 2013). - Twist1 is related to EMT in MCF-7 and MDA-MB-231 human breast cancer cell lines (<i>Homo sapiens</i>) (Menendez-Menendez et al., 2019). - Twist induces EMT in Huh7 human hepatocellular carcinoma cell lines (<i>Homo sapiens</i>) (Hu et al., 2019).	
Key Event Relationship Description	
Beta-catenin activation, of which mechanism include the stabilization of the dephosphorylated beta-catenin and translocation of beta-catenin into the nucleus, induce the formation of beta-catenin-TCF complex and transcription of transcription factors such as Snail, Zeb and Twist (Clevers & Nusse, 2012) (Ahmad et al., 2012; Pearlman, Montes de Oca, Pal, & Afaq, 2017; Sohn et al., 2019; Yang et al., 2019). EMT-related transcription factors including Snail, ZEB and Twist are up-regulated in cancer cells (Diaz, Vinas-Castells, & Garcia de Herreros, 2014). The transcription factors such as Snail, ZEB and Twist bind to E-cadherin (CDH1) promoter and inhibit the CDH1 transcription via the consensus E-boxes (5'-CACCTG-3' or 5'-CAGGTG-3'), which leads to EMT (Diaz et al., 2014).	
Evidence Supporting this KER	
Biological Plausibility	
The treatment of human gastric cancer cells with INC280, which inhibits c-MET overexpressed in diffuse-type gastric cancer with poor prognosis, shows downregulation in beta-catenin and Snail expression, (Sohn et al., 2019). The treatment with garcinol, a polyisoprenylated benzophenone derivative that is obtained from <i>Garcinia indica</i> extract, induced ZEB1 and ZEB2 down-regulation, increase in phosphorylated beta-catenin, and decrease in nuclear beta-catenin in human breast cancer cells (Ahmad et al., 2012). Sortilin, a member of the Vps10p sorting receptor family which is highly expressed in high-grade malignant glioma, positively regulates GSK-3beta/beta-catenin/Twist signaling pathway in glioblastoma (Yang et al., 2019). TM4SF1 promotes EMT via Wnt/beta-catenin/SOX2 pathway in colorectal cancer (Yang et al., 2020). The transcription factors such as Snail, Zeb, and Twist inhibit the CDH1 expression through their binding towards the promoter of CDH1, which leads to inhibition of cell adhesion and EMT (Diaz et al., 2014)	
Empirical Evidence	
[Dose concordance]	
The inhibition of sortilin by AF38469 (a sortilin inhibitor) or small interference RNA (siRNA) results in a decrease in beta-catenin and Twist expression in human glioblastoma cells (Yang et al., 2019).	
[Time concordance]	
The complex of beta-catenin and TCF4 induces epithelial-mesenchymal transition (EMT)-activator ZEB (Sanchez-Tillo E et al., 2011).	
[Incidence concordance]	
The inhibition of c-MET, which is overexpressed in diffuse-type gastric cancer, induced an increase in phosphorylated beta-catenin, decrease in beta-catenin and Snail (Sohn et al., 2019).	
The garcinol, which has an anti-cancer effect, increases phosphorylated beta-catenin, decreases beta-catenin and ZEB1/ZEB2, and inhibits EMT (Ahmad et al., 2012).	
Histone deacetylase inhibitors affect EMT-related transcription factors including ZEB, Twist, and Snail (Wawruszak et al., 2019).	
Snail and Zeb induces EMT and suppress E-cadherin (CDH1) (Batlle et al., 2000; Diaz et al., 2014; Peinado, Olmeda, & Cano, 2007).	
Uncertainties and Inconsistencies	
It is possible that the inhibition of ZEB1 and ZEB2 by garcinol treatment is caused by down-regulation of NFkappaB and Wnt/beta-catenin signaling (Ahmad et al., 2012).	
The EMT is induced different transcription factors other than Zeb, Twist, and Snail, which includes E47 and KLF4 (Diaz et al., 2014).	
Zeb, Twist, and Snail may activate or inactivate different genes or molecules to induce phenomena related to EMT and other phenomena other than EMT (Li & Balazsi, 2018).	
Quantitative Understanding of the Linkage	
Response-response relationship	
The treatment with AF38469, a sortilin inhibitor, in 0, 100, 200, 400, 800, and 1600 nM concentration inhibited beta-catenin and Twist (EMT regulator) expression dose-dependently in human glioblastoma cells (Yang et al., 2019).	
Snail (SNAIL, a key transcription factor of EMT induced by beta-catenin) mRNA is methylated, and N^6 -methyladenosine (m^6A) in its coding region	

(CDS) and 3' untranslated region (3'UTR) are significantly enriched during EMT progression (Lin et al., 2019). The m⁶A enrichment fold of *SNAI1* mRNA in EMT cells is about 2.3-fold greater than in control cells (Lin et al., 2019).

Time-scale

Nuclear accumulation of beta-catenin induces endogenous ZEB1 in 15 and 30 min (Sanchez-Tillo E et al., 2011).

The treatment with 25 uM of garcinol for 48 hours induced an increase in phosphorylated beta-catenin and decreased nuclear beta-catenin protein and ZEB1/ZEB2 mRNA in human breast cancer cells (Ahmad et al., 2012).

The treatment with AF38469, a sortilin inhibitor, for 0, 2, 4, 8, 16, or 24 hours shows that the expression of beta-catenin and Twist decrease in 8 hours followed by the subsequent decrease in 16 and 24 hours in human glioblastoma cells (Yang et al., 2019).

Snail (SNAI1) transfection for 48 hours induces the repression of E-cadherin (CDH1) protein expression (Lin et al., 2019).

SNAI1 mRNA in polysome is up-regulated in EMT-undergoing HeLa cells treated with 10 ng/ml of TGF-beta for 3 days compared with control cells (Lin et al., 2019).

Known modulating factors

The proto-oncogene MET regulates beta-catenin and Snail expression (Sohn et al., 2019).

The inhibition of GSK3beta by SB216763 induced expression of beta-catenin and Twist, as well as mesenchymal markers such as N-cadherin, vimentin, and MMP9 (Yang et al., 2019).

The decrease in E-cadherin (CDH1), a cell adhesion molecule, is related to EMT (Diaz et al., 2014).

Methyltransferase-like 3 (METTL3) modulates methylation of Snail (SNAI1) mRNA and EMT (Lin et al., 2019).

The binding of beta-catenin to members of the TCF/LEF family transcription factors increase gene expression related to EMT such as Twist and decrease E-cadherin protein expression (Qualtrough, Rees, Speight, Williams, & Paraskeva, 2015).

Known Feedforward/Feedback loops influencing this KER

The inhibited expression of phosphorylated GSK3beta, beta-catenin, and Twist by sortilin inhibition is reversed by GSK3beta inhibition. Furthermore, twist overexpression by lentivirus increased the inhibited expression of N-cadherin, MMP9, and vimentin and reverses the inhibitory effect of AF38469 on sortilin, which suggests that sortilin induces glioblastoma invasion mainly via GSK3beta/beta-catenin/Twist induced mesenchymal transition (Yang et al., 2019).

The inhibition of Hedgehog signaling pathway with cyclopamine reduces beta-catenin-TCF transcriptional activity, decreases the Twist expression, induces E-cadherin expression, and inhibits EMT (Qualtrough et al., 2015).

References

- Ahmad, A., Sarkar, S. H., Bitar, B., Ali, S., Aboukameel, A., Sethi, S., . . . Sarkar, F. H. (2012). Garcinol regulates EMT and Wnt signaling pathways in vitro and in vivo, leading to anticancer activity against breast cancer cells. *Mol Cancer Ther*, *11*(10), 2193-2201. doi:10.1158/1535-7163.MCT-12-0232-T
- Battle, E., Sancho, E., Francí, C., Domínguez, D., Monfar, M., Baulida, J., & García de Herreros, A. (2000). The transcription factor Snail is a repressor of E-cadherin gene expression in epithelial tumour cells. *Nature Cell Biology*, *2*(2), 84-89. doi:10.1038/35000034
- Clevers, H., & Nusse, R. (2012). Wnt/beta-catenin signaling and disease. *Cell*, *149*(6), 1192-1205. doi:10.1016/j.cell.2012.05.012
- Diaz, V. M., Vinas-Castells, R., & García de Herreros, A. (2014). Regulation of the protein stability of EMT transcription factors *Cell Adh Migr*, *8*(4), 418-428. doi:10.4161/19336918.2014.969998
- Hu, B., Cheng, J. W., Hu, J. W., Li, H., Ma, X. L., Tang, W. G., . . . Yang, X. R. (2019). KPNA3 Confers Sorafenib Resistance to Advanced Hepatocellular Carcinoma via TWIST Regulated Epithelial-Mesenchymal Transition. *Journal of Cancer*, *10*(17), 3914-3925. doi:10.7150/jca.31448
- Kudo-Saito, C., Shirako, H., Takeuchi, T., & Kawakami, Y. (2009). Cancer Metastasis Is Accelerated through Immunosuppression during Snail-Induced EMT of Cancer Cells. *Cancer Cell*, *15*(3), 195-206. doi:10.1016/j.ccr.2009.01.023
- Li, C., & Balazsi, G. (2018). A landscape view on the interplay between EMT and cancer metastasis *NPJ Syst Biol Appl*, *4*, 34. doi:10.1038/s41540-018-0068-x
- Lin, X., Chai, G., Wu, Y., Li, J., Chen, F., Liu, J., . . . Wang, H. (2019). RNA m(6A) methylation regulates the epithelial mesenchymal transition of cancer cells and translation of Snail. *Nat Commun*, *10*(1), 2065. doi:10.1038/s41467-019-09865-9
- Menendez-Menendez, J., Hermida-Prado, F., Granda-Diaz, R., Gonzalez, A., Garcia-Pedrero, J. M., Del-Rio-Ibáñez, N., . . . Martínez-Campa, C. (2019). Deciphering the Molecular Basis of Melatonin Protective Effects on Breast Cells Treated with Doxorubicin: TWIST1 a Transcription Factor Involved in EMT and Metastasis, a Novel Target of Melatonin. *Cancers (Basel)*, *11*(7). doi:10.3390/cancers11071011
- Ota, I., Masui, T., Kurihara, M., Yook, J. I., Mikami, S., Kimura, T., . . . Kitahara, T. (2016). Snail-induced EMT promotes cancer stem cell-like properties in head and neck cancer cells. *Oncol Rep*, *35*(1), 261-266. doi:10.3892/or.2015.4348
- Pearlman, R. L., Montes de Oca, M. K., Pal, H. C., & Afaq, F. (2017). Potential therapeutic targets of epithelial-mesenchymal transition in melanoma. *Cancer Lett*, *391*, 125-140. doi:10.1016/j.canlet.2017.01.029
- Peinado, H., Olmeda, D., & Cano, A. (2007). Snail, Zeb and bHLH factors in tumour progression: an alliance against the epithelial phenotype *Nat Rev Cancer*, *7*(6), 415-428. doi:10.1038/nrc2131
- Qualtrough, D., Rees, P., Speight, B., Williams, A. C., & Paraskeva, C. (2015). The Hedgehog Inhibitor Cyclopamine Reduces beta-Catenin-Tcf Transcriptional Activity, Induces E-Cadherin Expression, and Reduces Invasion in Colorectal Cancer Cells. *Cancers (Basel)*, *7*(3), 1885-1899. doi:10.3390/cancers7030867
- Rosmaninho, P., Mükusch, S., Piscopo, V., Teixeira, V., Raposo, A. A., Warta, R., . . . Castro, D. S. (2018). Zeb1 potentiates genome-wide gene transcription with Lef1 to promote glioblastoma cell invasion. *The EMBO Journal*, *37*(15), e97115. doi:10.15252/embj.201797115
- Sanchez-Tillo E, de Barrios O, Siles L, Cuatrecasas M, Castells A, Postigo A. beta-catenin/TCF4 complex induces the epithelial-to-mesenchymal transition (EMT)-activator ZEB1 to regulate tumor invasiveness. *Proc Natl Acad Sci U S A*, 2011;108(48):19204-9.
- Sohn, S. H., Kim, B., Sul, H. J., Kim, Y. J., Kim, H. S., Kim, H., . . . Zang, D. Y. (2019). INC280 inhibits Wnt/beta-catenin and EMT signaling pathways and its induce apoptosis in diffuse gastric cancer positive for c-MET amplification. *BMC Res Notes*, *12*(1), 125. doi:10.1186/s13104-019-4163-x

Tang Q, Chen J, Di Z, Yuan W, Zhou Z, Liu Z, Han S, Liu Y, Ying G, Shu X, Di M. TM4SF1 promotes EMT and cancer stemness via the Wnt/ β -catenin/SOX2 pathway in colorectal cancer. *J Exp Clin Cancer Res*. 2020 Nov 5;39(1):232. doi: 10.1186/s13046-020-01690-z. PMID: 33153498; PMCID: PMC7643364.

Wang, Y., Shi, J., Chai, K., Ying, X., & Zhou, B. P. (2013). The Role of Snail in EMT and Tumorigenesis *Current cancer drug targets*, 13(9), 963-972. doi: 10.2174/15680096113136660102

Wawruszak, A., Kalafut, J., Okon, E., Czapinski, J., Halasa, M., Przybyszewska, A., . . . Stepulak, A. (2019). Histone Deacetylase Inhibitors and Phenotypical Transformation of Cancer Cells. *Cancers (Basel)*, 11(2). doi:10.3390/cancers11020148

Xue, Y., Zhang, L., Zhu, Y., Ke, X., Wang, Q., & Min, H. (2019). Regulation of Proliferation and Epithelial-to-Mesenchymal Transition (EMT) of Gastric Cancer by ZEB1 via Modulating Wnt5a and Related Mechanisms. *Medical science monitor : international medical journal of experimental and clinical research*, 25, 1663-1670. doi:10.12659/MSM.912338

Yang, W., Wu, P. F., Ma, J. X., Liao, M. J., Wang, X. H., Xu, L. S., . . . Yi, L. (2019). Sortilin promotes glioblastoma invasion and mesenchymal transition through GSK-3 β /catenin/twist pathway. *Cell Death Dis*, 10(3), 208. doi:10.1038/s41419-019-1449-9

Relationship: 3200: EMT leads to Resistant gastric cancer

AOPs Referencing Relationship

AOP Name	Adjacency	Weight of Evidence	Quantitative Understanding
Increase in reactive oxygen species (ROS) leading to human treatment-resistant gastric cancer	adjacent	Moderate	Moderate

Evidence Supporting Applicability of this Relationship

Taxonomic Applicability

Term	Scientific Term	Evidence	Links
Homo sapiens	Homo sapiens	High	NCBI

Life Stage Applicability

Life Stage	Evidence
All life stages	High

Sex Applicability

Sex	Evidence
Unspecific	High

EMT induces cancer invasion, metastasis (*Homo sapiens*) ([P. Zhang et al., 2015](#)).

EMT is related to cancer drug resistance in MCF-7 human breast cancer cells (*Homo sapiens*) ([B. Du & Shim, 2016](#)).

Key Event Relationship Description

Some population of the cells exhibiting EMT demonstrates the feature of cancer stem cells (CSCs), which are related to cancer malignancy ([Shibue & Weinberg, 2017](#); [Shihori Tanabe, 2015a, 2015b](#); [Tanabe, Aoyagi, Yokozaki, & Sasaki, 2015](#)).

EMT phenomenon is related to cancer metastasis and cancer therapy resistance ([Smith & Bhowmick, 2016](#); [Tanabe, 2013](#)). The increased expression of enzymes that degrade the extracellular matrix components and the decrease in adhesion to the basement membrane in EMT induces the cell to escape from the basement membrane and metastasis ([Smith & Bhowmick, 2016](#)). Morphological changes observed during EMT are associated with therapy resistance ([Smith & Bhowmick, 2016](#)).

Evidence Supporting this KER

Biological Plausibility

The morphological and physiological changes associated with EMT are involved in invasiveness and drug resistance ([Shibue & Weinberg, 2017](#)). The EMT-activated particular carcinoma cells in primary tumors invade the surrounding stroma ([Shibue & Weinberg, 2017](#)). The EMT-activated carcinoma cells interact with the surrounding extracellular matrix protein to induce focal adhesion kinase and extracellular signal-related kinase activation, followed by the transforming growth factor-beta (TGF β) and canonical and/or noncanonical Wnt pathways to induce cancer stem cell (CSC) properties which contribute to the drug resistance ([Shibue & Weinberg, 2017](#)).

EMT-associated down-regulation of multiple apoptotic signaling pathways induces drug efflux and slows cell proliferation to induce the general resistance of carcinoma cells to anti-cancer drugs ([Shibue & Weinberg, 2017](#)).

Snail, an EMT-related transcription factor, induces the expression of the AXL receptor tyrosine kinase, which enables the cancer cells to survive by the activation of AXL signaling triggered by the binding of its ligand growth arrest-specific protein 6 (GAS6) ([Shibue & Weinberg, 2017](#)).

The EMT-activated cells evade the lethal effect of cytotoxic T cells, which include the elevated expression of programmed cell death 1 ligand (PD-L1) which binds to the programmed cell death protein 1 (PD-1) inhibitory immune-checkpoint receptor on the cell surface of cytotoxic T cells ([Shibue & Weinberg, 2017](#)).

Empirical Evidence

Incidence concordance

Slug/Snai2, a *ces-1*-related zinc finger transcription factor gene, confers resistance to p53-mediated apoptosis of hematopoietic progenitors by repressing *PUMA* (also known as *BBC3*, encoding Bcl-2-binding component 3) ([Inukai et al., 1999](#); [Shibue & Weinberg, 2017](#); [W.-S. Wu et al., 2005](#)).

EMT activation induces the expression of multiple members of the ATP-binding cassette (ABC) transporter family, which results in the resistance to doxorubicin ([Saxena, Stephens, Pathak, & Rangarajan, 2011](#); [Shibue & Weinberg, 2017](#)).

TGFbeta-1 induced EMT results in the acquisition of cancer stem cell (CSC) like properties [Pirozzi et al., 2011](#); [Shibue & Weinberg, 2017](#)).

Snail-induced EMT induces cancer metastasis and resistance to dendritic cell-mediated immunotherapy [Kudo-Saito, Shirako, Takeuchi, & Kawakami, 2009](#)).

Zinc finger E-box-binding homeobox (ZEB1)-induced EMT results in the relief of miR-200-mediated repression of programmed cell death 1 ligand (PD-L1) expression, a major inhibitory ligand for the programmed cell death protein (PD-1) immune-checkpoint protein on CD8+ cytotoxic T lymphocyte (CTL), subsequently the CD8+ T cell immunosuppression and metastasis [Chen et al., 2014](#)).

Uncertainties and Inconsistencies

The reversing process of EMT, which names as a mesenchymal-epithelial transition (MET), maybe one of the candidates for the anti-cancer therapy, where the plasticity of the cell phenotype is of importance and under investigation [Shibue & Weinberg, 2017](#)).

Quantitative Understanding of the Linkage

Response-response relationship

Induction of EMT by TGFbeta and Twist increases the gene expression of EMT markers such as Snail, Vimentin, N-cadherin, and ABC transporters including ABCA3, ABCC1, ABCC3, and ABCC10 [Saxena et al., 2011](#)).

Human mammary epithelial cells (HMLE) stably expressing Twist, FOXC2 or Snail demonstrates the increased cell viability compared to control HMLE in the treatment with about 0.3, 3, 30 mM of doxorubicin, dose-dependently [Saxena et al., 2011](#)).

Time-scale

The treatment with doxorubicin for 48 hours demonstrates the increase in the cell viability in Twist/FOXC2/Snail overexpressed HMLE compared to control HMLE [Saxena et al., 2011](#)).

The inhibition of Twist or Zeb1 with small interference RNA (siRNA) induced the inhibition of cell viability compared to control MDAMB231 cells treated with doxorubicin for 48 hours [Saxena et al., 2011](#)).

Known modulating factors

ABC transporters that are related to drug resistance are overexpressed in the EMT-activated cells [Saxena et al., 2011](#)). The expression of PD-L1, which binds to the PD-1 on the cytotoxic T cells, is up-regulated in EMT-activated cells, which results in the inhibition of cancer immunity and the resistance to cancer therapy [Shibue & Weinberg, 2017](#)).

Known Feedforward/Feedback loops influencing this KER

The investigation of EMT-CSC relations is important to understand the relationship between EMT and cancer malignancy. Non-CSCs in cancer can spontaneously undergo EMT and dedifferentiate into new CSC, subsequently induce the regeneration of tumorigenic potential [Marjanovic, Weinberg, & Chaffer, 2013](#); [Shibue & Weinberg, 2017](#)).

The plastic CSC theory demonstrates the bidirectional conversions between non-CSCs and CSCs, which may contribute to the acquisition of cancer malignancy in EMT-activated cells [Marjanovic et al., 2013](#)).

References

- Chen, L., Gibbons, D. L., Goswami, S., Cortez, M. A., Ahn, Y.-H., Byers, L. A., . . . Qin, F. X.-F. (2014). Metastasis is regulated via microRNA-200/ZEB1 axis control of tumour cell PD-L1 expression and intratumoral immunosuppression. *Nature communications*, 5, 5241-5241. doi:10.1038/ncomms6241
- Du, B., & Shim, J. S. (2016). Targeting Epithelial-Mesenchymal Transition (EMT) to Overcome Drug Resistance in Cancer *Molecules*, 21(7). doi:10.3390/molecules21070965
- Inukai, T., Inoue, A., Kurosawa, H., Goi, K., Shinjyo, T., Ozawa, K., . . . Look, A. T. (1999). SLUG, a ces-1-Related Zinc Finger Transcription Factor Gene with Antiapoptotic Activity, Is a Downstream Target of the E2A-HLF Oncoprotein. *Molecular Cell*, 4(3), 343-352. doi:[https://doi.org/10.1016/S1097-2765\(00\)80336-6](https://doi.org/10.1016/S1097-2765(00)80336-6)
- Kudo-Saito, C., Shirako, H., Takeuchi, T., & Kawakami, Y. (2009). Cancer Metastasis Is Accelerated through Immunosuppression during Snail-Induced EMT of Cancer Cells. *Cancer Cell*, 15(3), 195-206. doi:<https://doi.org/10.1016/j.ccr.2009.01.023>
- Marjanovic, N. D., Weinberg, R. A., & Chaffer, C. L. (2013). Cell plasticity and heterogeneity in cancer. *Clinical chemistry*, 59(1), 168-179. doi:10.1373/clinchem.2012.184655
- Pirozzi, G., Tirino, V., Camerlingo, R., Franco, R., La Rocca, A., Liguori, E., . . . Rocco, G. (2011). Epithelial to mesenchymal transition by TGFβ-1 induction increases stemness characteristics in primary non small cell lung cancer cell line. *PLoS One*, 6(6), e21548-e21548. doi:10.1371/journal.pone.0021548
- Saxena, M., Stephens, M. A., Pathak, H., & Rangarajan, A. (2011). Transcription factors that mediate epithelial-mesenchymal transition lead to multidrug resistance by upregulating ABC transporters. *Cell death & disease*, 2(7), e179-e179. doi:10.1038/cddis.2011.61
- Shibue, T., & Weinberg, R. A. (2017). EMT, CSCs, and drug resistance: the mechanistic link and clinical implications *Nat Rev Clin Oncol*, 14(10), 611-629. doi:10.1038/nrclinonc.2017.44
- Smith, B. N., & Bhowmick, N. A. (2016). Role of EMT in Metastasis and Therapy Resistance *J Clin Med*, 5(2). doi:10.3390/jcm5020017
- Tanabe, S. (2013). Perspectives of gene combinations in phenotype presentation. *World journal of stem cells*, 5(3), 61-67. doi:10.4252/wjsc.v5.i3.61
- Tanabe, S. (2015a). Origin of cells and network information. *World journal of stem cells*, 7(3), 535-540. doi:10.4252/wjsc.v7.i3.535
- Tanabe, S. (2015b). Signaling involved in stem cell reprogramming and differentiation. *World journal of stem cells*, 7(7), 992-998. doi:10.4252/wjsc.v7.i7.992
- Tanabe, S., Aoyagi, K., Yokozaki, H., & Sasaki, H. (2015). Regulated genes in mesenchymal stem cells and gastric cancer *World journal of stem cells*, 7(1), 208-222. doi:10.4252/wjsc.v7.i1.208
- Wu, W.-S., Heinrichs, S., Xu, D., Garrison, S. P., Zambetti, G. P., Adams, J. M., & Look, A. T. (2005). Slug Antagonizes p53-Mediated Apoptosis of Hematopoietic Progenitors by Repressing puma. *Cell*, 123(4), 641-653. doi:<https://doi.org/10.1016/j.cell.2005.09.029>
- Zhang, P., Sun, Y., & Ma, L. (2015). ZEB1: at the crossroads of epithelial-mesenchymal transition, metastasis and therapy resistance *Cell Cycle*, 14(4), 481-487. doi:10.1080/15384101.2015.1006048

