


AOP ID and Title:

AOP 306: Androgen receptor (AR) antagonism leading to short anogenital distance (AGD) in male (mammalian) offspring

Short Title: AR antagonism leading to short AGD

Graphical Representation**Authors**

Sofie Christiansen; National Food Institute, Technical University of Denmark, Kongens Lyngby, 2800 Denmark

Monica Kam Draskau; National Food Institute, Technical University of Denmark, Kongens Lyngby, 2800 Denmark

Terje Svingen; National Food Institute, Technical University of Denmark, Kongens Lyngby, 2800 Denmark

Status

Author status	OECD status	OECD project	SAAOP status
Under development: Not open for comment. Do not cite	Under Development	1.90	Included in OECD Work Plan

	Author status	OECD status	OECD project	SAAOP status																																																								
Abstract This AOP links Androgen receptor antagonism during fetal life with short anogenital distance (AGD) in male offspring. A short AGD around birth is a marker for feminization of male fetuses and is associated with male reproductive disorders, including reduced fertility in adulthood. Although a short AGD is not necessarily ‘adverse’ from a human health perspective, it is considered an ‘adverse outcome’ in OECD test guidelines; AGD measurements are mandatory in specific tests for developmental and reproductive toxicity in chemical risk assessment (TG 443, TG 421/422, TG 414). The AR is a nuclear receptor involved in the transcriptional regulation of various target genes during development and adulthood across species. Its main ligand is testosterone and dihydrotestosterone (DHT). Under normal physiological conditions, testosterone produced mainly by the testicles, is converted in peripheral tissues by 5α-reductase into DHT, which in turn binds AR and activates downstream target genes. AR signaling is necessary for normal masculinization of the developing fetus, including differentiation of the levator ani/bulbocavernosus (LABC) muscle complex in male fetuses. The LABC complex does not develop in the absence, or low levels of, androgen signaling, as in female fetuses. The key events in this pathway is antagonism of the AR in target cells of the primitive perineal region, which leads to inactivation of the AR and failure to properly masculinize the perineum/LABC complex. In this instance, the local levels of testosterone or DHT may be normal, but prevented from binding the AR. Summary of the AOP Events Molecular Initiating Events (MIE), Key Events (KE), Adverse Outcomes (AO) <table><tr><th>Sequence</th><th>Type</th><th>Event ID</th><th>Title</th><th>Short name</th></tr><tr><td></td><td>MIE</td><td>26</td><td>Antagonism, Androgen receptor</td><td>Antagonism, Androgen receptor</td></tr><tr><td></td><td>KE</td><td>1614</td><td>Decrease, androgen receptors (AR) activation</td><td>Decrease, AR activation</td></tr><tr><td></td><td>KE</td><td>1687</td><td>decrease, transcription of genes by AR</td><td>decrease, transcription of genes by AR</td></tr><tr><td></td><td>AO</td><td>1688</td><td>decrease, male anogenital distance</td><td>short male AGD</td></tr></table> Key Event Relationships <table><tr><th>Upstream Event</th><th>Relationship Type</th><th>Downstream Event</th><th>Evidence</th><th>Quantitative Understanding</th></tr><tr><td>Antagonism, Androgen receptor</td><td>adjacent</td><td>Decrease, androgen receptors (AR) activation</td><td>High</td><td>High</td></tr><tr><td>Decrease, androgen receptors (AR) activation</td><td>adjacent</td><td>decrease, transcription of genes by AR</td><td>High</td><td>Moderate</td></tr><tr><td>decrease, transcription of genes by AR</td><td>adjacent</td><td>decrease, male anogenital distance</td><td>Moderate</td><td>Low</td></tr><tr><td>Antagonism, Androgen receptor</td><td>non-adjacent</td><td>decrease, male anogenital distance</td><td>Moderate</td><td>Low</td></tr></table> Stressors <table><tr><th>Name</th><th>Evidence</th></tr><tr><td>Finasteride</td><td>High</td></tr><tr><td>Flutamide</td><td>High</td></tr></table> Finasteride Intrauterine exposure in rats can result in shorter male AGD in male offspring as reported in: Bowman et al (2003), Toxicol Sci 74:393-406; doi: 10.1093/toxsci/kfg128					Sequence	Type	Event ID	Title	Short name		MIE	26	Antagonism, Androgen receptor	Antagonism, Androgen receptor		KE	1614	Decrease, androgen receptors (AR) activation	Decrease, AR activation		KE	1687	decrease, transcription of genes by AR	decrease, transcription of genes by AR		AO	1688	decrease, male anogenital distance	short male AGD	Upstream Event	Relationship Type	Downstream Event	Evidence	Quantitative Understanding	Antagonism, Androgen receptor	adjacent	Decrease, androgen receptors (AR) activation	High	High	Decrease, androgen receptors (AR) activation	adjacent	decrease, transcription of genes by AR	High	Moderate	decrease, transcription of genes by AR	adjacent	decrease, male anogenital distance	Moderate	Low	Antagonism, Androgen receptor	non-adjacent	decrease, male anogenital distance	Moderate	Low	Name	Evidence	Finasteride	High	Flutamide	High
Sequence	Type	Event ID	Title	Short name																																																								
	MIE	26	Antagonism, Androgen receptor	Antagonism, Androgen receptor																																																								
	KE	1614	Decrease, androgen receptors (AR) activation	Decrease, AR activation																																																								
	KE	1687	decrease, transcription of genes by AR	decrease, transcription of genes by AR																																																								
	AO	1688	decrease, male anogenital distance	short male AGD																																																								
Upstream Event	Relationship Type	Downstream Event	Evidence	Quantitative Understanding																																																								
Antagonism, Androgen receptor	adjacent	Decrease, androgen receptors (AR) activation	High	High																																																								
Decrease, androgen receptors (AR) activation	adjacent	decrease, transcription of genes by AR	High	Moderate																																																								
decrease, transcription of genes by AR	adjacent	decrease, male anogenital distance	Moderate	Low																																																								
Antagonism, Androgen receptor	non-adjacent	decrease, male anogenital distance	Moderate	Low																																																								
Name	Evidence																																																											
Finasteride	High																																																											
Flutamide	High																																																											

Christiansen et al (2009), Environ Health Perspect 117:1839-1846; doi: 10.1289/ehp.0900689

Schwartz et al (2019), Toxicol Sci 169:303-311; doi: 10.1093/toxsci/kfz046

Flutamide

Finasteride is a selective androgen receptor (AR) antagonist (Simard et al 1986) that has been shown to induce shorter male AGD in rats after in utero exposure (Foster & Harris 2005; Hass et al 2007; Kita et al 2016; McIntyre et al 2001; Mylchreest et al 1999; Scott et al 2007; Welsh et al 2007).

References:

Foster & Harris (2005), Toxicol Sci 85:1024-1032; doi: 10.1093/toxsci/kfi159

Hass et al (2007), Environ Health Perspect 115(suppl 1):122-128; doi: 10.1289/ehp.0360

Kita et al (2016), Toxicology 368-369:152-161; doi: 10.1016/j.tox.2016.08.021

McIntyre et al (2001), Toxicol Sci 62:236-249; doi: 10.1093/toxsci/62.2.236

Mylchreest et al (1999), Toxicol Appl Pharmacol 156:81-95; doi: 10.1006/taap.1999.8643

Scott et al (2007), Endocrinology 148:2027-2036; doi: 10.1210/en.2006-1622

Simard et al (1986), Mol Cell Endocrinol 44:261-270; doi: 10.1016/0303-7207(86)90132-2

Overall Assessment of the AOP

Domain of Applicability

Life Stage Applicability

Life Stage Evidence

Pregnancy High

Taxonomic Applicability

Term	Scientific Term	Evidence	Links
human	Homo sapiens	Moderate	NCBI
rat	Rattus norvegicus	High	NCBI
mouse	Mus musculus	Moderate	NCBI

Sex Applicability

Sex Evidence

Male High

References

1. Schwartz CL, Christiansen S, Vinggaard AM, Axelstad M, Hass U and **Svingen T** (2019), Anogenital distance as a toxicological or clinical marker for fetal androgen action and risk for reproductive disorders. *Arch Toxicol* 93: 253-272.

Appendix 1

List of MIEs in this AOP

[Event: 26: Antagonism, Androgen receptor](#)

Short Name: Antagonism, Androgen receptor

AOPs Including This Key Event

AOP ID and Name	Event Type
Aop:306 - Androgen receptor (AR) antagonism leading to short anogenital distance (AGD) in male (mammalian) offspring	MolecularInitiatingEvent
Aop:344 - Androgen receptor (AR) antagonism leading to nipple retention (NR) in male (mammalian) offspring	MolecularInitiatingEvent
Aop:345 - Androgen receptor (AR) antagonism leading to decreased fertility in females	MolecularInitiatingEvent
Aop:372 - Androgen receptor antagonism leading to testicular cancer	MolecularInitiatingEvent

Stressors

Name

Mercaptobenzole
 Triticonazole
 Flusilazole
 Epoxiconazole
 Prochloraz
 Propiconazole
 Tebuconazole
 Flutamide
 Cyproterone acetate
 Vinclozolin

Biological Context

Level of Biological Organization

Molecular

Cell term

Cell term

eukaryotic cell

Evidence for Perturbation by Stressor

Overview for Molecular Initiating Event

A large number of drugs and chemicals have been shown to antagonise the AR using various AR reporter gene assays. The AR is specifically targeted in AR-sensitive cancers, for example the use of the anti-androgenic drug flutamide in treating prostate cancer ([Alapi & Fischer, 2006](#)). Flutamide has also been used in several rodent in vivo studies showing anti-androgenic effects (feminization of male offspring) evident by e.g. short anogenital distance (AGD) in males ([Foster & Harris, 2005](#); [Hass et al. 2007](#); [Kita et al. 2016](#)). QSAR models can predict AR antagonism for a wide range of chemicals, many of which have shown in vitro antagonistic potential ([Vinggaard et al. 2008](#)).

Triticonazole

Using hAR-EcoScreen Assay, triticonazole showed a LOEC for antagonisms of 0.2 uM and an IC50 of 0.3 (±0.01) uM ([Draskau et al. 2019](#))

Flusilazole

Using hAR-EcoScreen Assay, flusilazole showed a LOEC for antagonisms of 0.8 μM and an IC₅₀ of 2.8 (± 0.1) μM ([Draskau et al. 2019](#)). ►

Epoxiconazole

Using transiently AR-transfected CHO cells, epoxiconazole showed a LOEC of 1.6 μM and an IC₅₀ of 10 μM ([Kjærstad et al. 2010](#))

Prochloraz

Using transiently AR-transfected CHO cells, prochloraz showed a LOEC of 6.3 μM and an IC₅₀ of 13 μM ([Kjærstad et al. 2010](#))

Propiconazole

Using transiently AR-transfected CHO cells, propiconazole showed a LOEC of 12.5 μM and an IC₅₀ of 18 μM ([Kjærstad et al. 2010](#))

Tebuconazole

Using transiently AR-transfected CHO cells, tebuconazole showed a LOEC of 3.1 μM and an IC₅₀ of 8.1 μM ([Kjærstad et al. 2010](#))

Flutamide

Using the AR-CALUX reporter assay in antagonism mode, flutamide showed an IC₅₀ of 1.3 μM ([Sonneveld et al. 2005](#)).

Cyproterone acetate

Using the AR-CALUX reporter assay in antagonism mode, cyproterone acetate showed an IC₅₀ of 7.1 nM ([Sonneveld et al. 2005](#)).

Vinclozolin

Using the AR-CALUX reporter assay in antagonism mode, vinclozolin showed an IC₅₀ of 1.0 μM ([Sonneveld et al. 2005](#)).

Domain of Applicability**Taxonomic Applicability**

Term	Scientific Term	Evidence	Links
human	Homo sapiens	High	NCBI
mouse	Mus musculus	High	NCBI
rat	Rattus norvegicus	High	NCBI
human and other cells in culture	human and other cells in culture	High	NCBI

Life Stage Applicability

Life Stage	Evidence
Foetal	High
Embryo	Moderate

During development and at adulthood High

Sex Applicability**Sex Evidence**

Mixed High

Both the DNA-binding and ligand-binding domains of the AR are highly evolutionary conserved, whereas the transactivation domain show more divergence which may affect AR-mediated gene regulation across species ([Davey & Grossmann, 2016](#)). Despite certain inter-species differences, AR function mediated through gene expression is highly conserved, with mutations studies from both humans and rodents showing strong correlation for AR-dependent development and function ([Walters et al. 2010](#)).

This KE is applicable for both sexes, across developmental stages into adulthood, in numerous cells and tissues and across taxa

Key Event Description

The androgen receptor (AR) and its function

Development of the male reproductive system and secondary male characteristics is dependent on androgens (foremost testosterone (T) and dihydrotestosterone (DHT)). T and the more biologically active DHT act by binding to the AR ([MacLean et al. 1993](#); [MacLeod et al. 2010](#); [Schwartz et al. 2019](#)), with human AR mutations and mouse knock-out models having established its pivotal role in masculinization and spermatogenesis ([Walters et al. 2010](#)). The AR is a ligand-activated transcription factor belonging to the steroid hormone nuclear receptor family ([Davey & Grossmann. 2016](#)). The AR has three domains; the N-terminal domain, the DNA-binding domain and the ligand-binding domain, with the latter being most evolutionary conserved. Apart from the essential role AR plays for male reproductive development and function ([Walters et al. 2010](#)), the AR is also expressed in many other tissues and organs such as bone, muscles, ovaries and the immune system ([Rana et al. 2014](#)).

AR antagonism as Key Event

The main function of the AR is to activate gene transcription in cells. Canonical signaling occurs by ligands (androgens) binding to AR in the cytoplasm which results in translocation to the cell nucleus, receptor dimerization and binding to specific regulatory DNA sequences ([Heemers & Tindall. 2007](#)). The gene targets regulated by AR activation depends on cell/tissue type and what stage of development activation occur, and is, for instance, dependent on available co-factors. Apart from the canonical signaling pathway, AR can also function through non-genomic modalities, for instance rapid change in cell function by ion transport changes ([Heinlein & Chang. 2002](#)). However, with regard to this specific KE the canonical signaling pathway is what is referred to.

How it is Measured or Detected

AR antagonism can be measured in vitro by transient or stable transactivation assays to evaluate nuclear receptor activation. There is already a validated assay for AR (ant)agonism adopted by the OECD, Test No. 458: *Stably Transfected Human Androgen Receptor Transcriptional Activation Assay for Detection of Androgenic Agonist and Antagonist Activity of Chemicals* ([OECD. 2016](#)). The stably transfected AR-EcoScreen™ cells ([Sato et al. 2004](#)) should be used for the assay and is freely available for the Japanese Collection of Research Bioresources (JCRB) Cell Bank under reference number JCRB1328.

Other assays include the AR-CALUX reporter gene assay that is derived from human U2-OS cells stably transfected with the human AR and an AR responsive reporter gene ([van der Burg et al. 2010](#)), various transiently transfected reporter cell lines ([Körner et al. 2004](#)), and more.

Recently developed AR dimerization assay may soon be included in TGs for its improved ability to measure potential stressor-mediated dimerization/activation ([Lee et al. 2021](#)).

References

- Alapi EM, Fischer J (2006) Table of Selected Analogue Classes. In *Analogue-based Drug Discovery*, Fischer J, Ganellin CR (eds), p 515. Weinheim: Wiley-VCH Verlag GmbH & Co
- Davey RA, Grossmann M (2016) Androgen Receptor Structure, Function and Biology: From Bench to Bedside. *Clin Biochem Rev* **37**: 3-15
- Draskau MK, Boberg J, Taxvig C, Pedersen M, Frandsen HL, Christiansen S, Svingen T (2019) In vitro and in vivo endocrine disrupting effects of the azole fungicides triticonazole and flusilazole. *Environ Pollut* **255**: 113309
- Foster PM, Harris MW (2005) Changes in androgen-mediated reproductive development in male rat offspring following exposure to a single oral dose of flutamide at different gestational ages. *Toxicol Sci* **85**: 1024-1032
- Hass U, Scholze M, Christiansen S, Dalgaard M, Vinggaard AM, Axelstad M, Metzдорff SB, Kortenkamp A (2007) Combined exposure to anti-androgens exacerbates disruption of sexual differentiation in the rat. *Environ Health Perspect* **115 Suppl. 1**: 122-128
- Heemers HV, Tindall DJ (2007) Androgen receptor (AR) coregulators: a diversity of functions converging on and regulating the AR transcriptional complex. *Endocr Rev* **28**: 778-808
- Heinlein CA, Chang C (2002) The roles of androgen receptors and androgen-binding proteins in nongenomic androgen actions. *Mol Endocrinol* **16**: 2181-2187
- Kita DH, Meyer KB, Venturelli AC, Adams R, Machado DL, Morais RN, Swan SH, Gennings C, Martino-Andrade AJ (2016) Manipulation of pre and postnatal androgen environments and anogenital distance in rats. *Toxicology* **368-369**: 152-161
- Kjærstad MB, Taxvig C, Nellemann C, Vinggaard AM, Andersen HR (2010) Endocrine disrupting effects in vitro of conazole antifungals used as pesticides and pharmaceuticals. *Reprod Toxicol* **30**: 573-582
- Körner W, Vinggaard AM, Térouanne B, Ma R, Wieloch C, Schlumpf M, Sultan C, Soto AM (2004) Interlaboratory comparison of four in vitro assays for assessing androgenic and antiandrogenic activity of environmental chemicals. *Environ Health Perspect* **112**: 695-702
- Lee SH, Hong KY, Seo H, Lee HS, Park Y (2021) Mechanistic insight into human androgen receptor-mediated endocrine-disrupting potentials by a stable bioluminescence resonance energy transfer-based dimerization assay. *Chem Biol Interact* **349**: 109655
- MacLean HE, Chu S, Warne GL, Zajac JD (1993) Related individuals with different androgen receptor gene deletions. *J Clin Invest* **91**: 1123-

1128

MacLeod DJ, Sharpe RM, Welsh M, Fiskens M, Scott HM, Hutchison GR, Drake AJ, van den Driesche S (2010) Androgen action in the masculinization programming window and development of male reproductive organs. *Int J Androl* **33**: 279-287

OECD. (2016) Test No. 458: Stably Transfected Human Androgen Receptor Transcriptional Activation Assay for Detection of Androgenic Agonist and Antagonist Activity of Chemicals. *OECD Guidelines for the Testing of Chemicals, Section 4*, Paris.

Rana K, Davey RA, Zajac JD (2014) Human androgen deficiency: insights gained from androgen receptor knockout mouse models. *Asian J Androl* **16**: 169-177

Satoh K, Ohyama K, Aoki N, Iida M, Nagai F (2004) Study on anti-androgenic effects of bisphenol A diglycidyl ether (BADGE), bisphenol F diglycidyl ether (BFDGE) and their derivatives using cells stably transfected with human androgen receptor, AR-EcoScreen. *Food Chem Toxicol* **42**: 983-993

Schwartz CL, Christiansen S, Vinggaard AM, Axelstad M, Hass U, Svingen T (2019) Anogenital distance as a toxicological or clinical marker for fetal androgen action and risk for reproductive disorders. *Arch Toxicol* **93**: 253-272

Sonneveld E, Jansen HJ, Riteco JA, Brouwer A, van der Burg B (2005) Development of androgen- and estrogen-responsive bioassays, members of a panel of human cell line-based highly selective steroid-responsive bioassays. *Toxicol Sci* **83**: 136-148

van der Burg B, Winter R, Man HY, Vangenechten C, Berckmans P, Weimer M, Witters H, van der Linden S (2010) Optimization and prevalidation of the in vitro AR CALUX method to test androgenic and antiandrogenic activity of compounds. *Reprod Toxicol* **30**: 18-24

Vinggaard AM, Niemelä J, Wedebye EB, Jensen GE (2008) Screening of 397 chemicals and development of a quantitative structure-activity relationship model for androgen receptor antagonism. *Chem Res Toxicol* **21**: 813-823

Walters KA, Simanainen U, Handelsman DJ (2010) Molecular insights into androgen actions in male and female reproductive function from androgen receptor knockout models. *Hum Reprod Update* **16**: 543-558

List of Key Events in the AOP

Event: 1614: Decrease, androgen receptors (AR) activation

Short Name: Decrease, AR activation

AOPs Including This Key Event

AOP ID and Name	Event Type
Aop:288 - Inhibition of 17α-hydrolase/C 10.20-lyase (Cyp17A1) activity leads to birth reproductive defects (cryptorchidism) in male (mammals)	KeyEvent
Aop:305 - 5α-reductase inhibition leading to short anogenital distance (AGD) in male (mammalian) offspring	KeyEvent
Aop:306 - Androgen receptor (AR) antagonism leading to short anogenital distance (AGD) in male (mammalian) offspring	KeyEvent
Aop:307 - Decreased testosterone synthesis leading to short anogenital distance (AGD) in male (mammalian) offspring	KeyEvent
Aop:344 - Androgen receptor (AR) antagonism leading to nipple retention (NR) in male (mammalian) offspring	KeyEvent
Aop:372 - Androgen receptor antagonism leading to testicular cancer	KeyEvent

Biological Context

Level of Biological Organization

Cellular

Key Event Description

Androgen receptor activation is regulated by the binding of androgens. AR activity can be decreased by either a lack of steroidal ligands (testosterone, DHT) or the presence of antagonist compounds.¹²

How it is Measured or Detected

Significance of AR signaling in fetal development can be studied through a conditional deletion of the androgen receptor using a Cre/loxP approach. The recommended animal model for reproductive study is the mouse.³

Also, epidemiological case-studies following mouse or humans expressing a complete androgen insensitivity allow to directly assess the effects of a lack of AR activation on the development.⁴

Enzyme immunoassay (ELISA) kits for in vitro quantitative measurement of AR activity are available. Androgen receptors activity can be measured using bioassay such as the (Anti-)Androgen Receptor CALUX reporter gene assay.⁵

References

- ¹ Davey R.A and Grossmann M. (2016) Androgen Receptor Structure, Function and Biology: From Bench to Bedside. Clinical Biochemist Reviews, 37(1): 3-15. PCM4810760
- ² Gao W., Bohl C.E. and Dalton J.T. (2005) Chemistry and Structural Biology of Androgen Receptor. Chemical Reviews 105(9): 3352-3370 <https://doi.org/10.1021/cr020456u>
- ³ Kaftanovskaya E.M., Huang Z., Barbara A.M., De Gendt K., Verhoeven G., Ivan P. Gorlov, and Agoulnik A.I. (2012) Cryptorchidism in Mice with an Androgen Receptor Ablation in Gubernaculum Testis. Molecular Endocrinology, 26(4): 598-607. <https://doi.org/10.1210/me.2011-1283>
- ⁴ Hutson J.M. (1985) A biphasic model for the hormonal control of testicular descent. Lancet, 24;2(8452): 419-21 [http://dx.doi.org/10.1016/S0140-6736\(85\)92739-4](http://dx.doi.org/10.1016/S0140-6736(85)92739-4)
- ⁵ van der Burg B., Winter R., Man HY., Vangenechten C., Berckmans P., Weimer M., Witters M. and van der Linden S. (2010) Optimization and prevalidation of the in vitro AR CALUX method to test androgenic and antiandrogenic activity of compounds. Reproductive Toxicology, 30(1):18-24 <https://doi.org/10.1016/j.reprotox.2010.04.012>

Event: 1687: decrease, transcription of genes by AR

Short Name: decrease, transcription of genes by AR

AOPs Including This Key Event

AOP ID and Name	Event Type
Aop:306 - Androgen receptor (AR) antagonism leading to short anogenital distance (AGD) in male (mammalian) offspring	KeyEvent
Aop:372 - Androgen receptor antagonism leading to testicular cancer	KeyEvent

Biological Context

Level of Biological Organization

Cellular

List of Adverse Outcomes in this AOP

Event: 1688: decrease, male anogenital distance

Short Name: short male AGD

Key Event Component

Process	Object	Action
androgen receptor signaling pathway	Musculature of male perineum	disrupted

AOPs Including This Key Event

AOP ID and Name	Event Type
Aop:305 - 5α-reductase inhibition leading to short anogenital distance (AGD) in male (mammalian) offspring	AdverseOutcome
Aop:306 - Androgen receptor (AR) antagonism leading to short anogenital distance (AGD) in male (mammalian) offspring	AdverseOutcome
Aop:307 - Decreased testosterone synthesis leading to short anogenital distance (AGD) in male (mammalian) offspring	AdverseOutcome

Stressors

Name

Butylparaben
 p,p'-DDE
 Bis(2-ethylhexyl) phthalate
 Dexamethasone
 Fenitrothion
 Finasteride
 Flutamide
 Ketoconazole
 Linuron
 Prochloraz
 Procymidone
 Triticonazole
 Vinclozolin
 di-n-hexyl phthalate
 Dicyclohexyl phthalate
 butyl benzyl phthalate
 monobenzyl phthalate
 di-n-heptyl phthalate

Biological Context

Level of Biological Organization

Tissue

Organ term

Organ term

perineum

Evidence for Perturbation by Stressor

Butylparaben

Butylparaben has been shown to cause decreased male AGD in rats following intrauterine exposure to 500 and 1000 mg/kg bw/day ([Boberg et al. 2016](#); [Zhang et al. 2014](#)). A separate study using 600 mg/kg bw/day did not see an effect on male AGD ([Boberg et al. 2008](#)).

p,p'-DDE

p,p,DDE has been shown to cause decreased male AGD in rats following intrauterine exposure to 100-200 mg/kg bw/day ([Loeffler & Peterson, 1999](#); [Wolf et al. 1999](#)).

Bis(2-ethylhexyl) phthalate

DEHP has been shown to cause decreased male AGD in rats following intrauterine exposure to 300-1500 mg/kg bw/day ([Christiansen et al. 2010](#); [Gray et al. 2000](#); [Howdeshell et al. 2007](#); [Jarfelt et al. 2005](#); [Kita et al. 2016](#); [Li et al. 2013](#); [Lin et al. 2009](#); [Moore et al. 2001](#); [Nardelli et al. 2017](#); [Saillenfait et al. 2009](#); [Wolf et al. 1999](#)).

Dexamethasone

Dexamethasone has been shown to cause decreased male AGD in rats following intrauterine exposure to 0.1 mg/kg bw/day ([Van den Driesche et al. 2012](#)).

Fenitrothion

Fenitrothion has been shown to cause decreased male AGD in rats following intrauterine exposure to 25 mg/kg bw/day ([Turner et al. 2002](#)).

Finasteride

Finasteride has been shown to cause decreased male AGD in rats following intrauterine exposure to 100 mg/kg bw/day ([Bowman et al. 2003](#)).

Flutamide

Flutamide has been shown to cause decreased male AGD in rats following intrauterine exposure to doses between 16-100 mg/kg bw/day ([Foster & Harris, 2005](#); [Hass et al. 2007](#); [Kita et al. 2016](#); [McIntyre et al. 2001](#); [Mylchreest et al. 1999](#); [Scott et al. 2007](#); [Welsh et al. 2007](#)).

Ketoconazole

Ketoconazole has been shown to cause decreased male AGD in rats following intrauterine exposure to 50 mg/kg bw/day in one study ([Taxvig et al. 2008](#)), but no effect in another study using same dose ([Wolf et al. 1999](#)).

Linuron

Linuron has been shown to cause decreased male AGD in rats following intrauterine exposure to 50-100 mg/kg bw/day ([Hotchkiss et al. 2004](#); [McIntyre et al. 2002](#); [Wolf et al. 1999](#)).

Prochloraz

Prochloraz has been shown to cause decreased male AGD in rats following intrauterine exposure to 150-250 mg/kg bw/day ([Laier et al. 2006](#); [Noriega et al. 2005](#)).

Procymidone

Procymidone has been shown to cause decreased male AGD in rats following intrauterine exposure to doses between 50-150 mg/kg bw/day ([Hass et al. 2012](#); [Hass et al. 2007](#); [Wolf et al. 1999](#)).

Triticonazole

Triticonazole has been shown to cause decreased male AGD in rats following intrauterine exposure to 150 and 450 mg/kg bw/day ([Draskau et al. 2019](#)).

Vinclozolin

Vinclozolin has been shown to cause decreased male AGD in rats following intrauterine exposure to doses between 50-200 mg/kg bw/day ([Christiansen et al. 2009](#); [Gray et al. 1994](#); [Hass et al. 2007](#); [Matsuura et al. 2005](#); [Ostby et al. 1999](#); [Schneider et al. 2011](#); [Wolf et al. 2004](#)).

di-n-hexyl phthalate

DnHP has been shown to cause decreased male AGD in rats following intrauterine exposure to 500-750 mg/kg bw/day ([Saillenfait et al. 2009a](#); [Saillenfait et al. 2009b](#)).

Dicyclohexyl phthalate

DCHP has been shown to cause decreased male AGD in rats following intrauterine exposure to 350-750 mg/kg bw/day ([Aydoğan Ahbab & Barlas. 2015](#); [Hoshino et al. 2005](#); [Saillenfait et al. 2009a](#)).

butyl benzyl phthalate

BBP has been shown to cause decreased male AGD in rats following intrauterine exposure to 500-1000 mg/kg bw/day ([Ema & Miyawaki. 2002](#); [Gray et al. 2000](#); [Hotchkiss et al. 2004](#); [Nagao et al. 2000](#); [Tyl et al. 2004](#)).

monobenzyl phthalate

MBP has been shown to cause decreased male AGD in rats following intrauterine exposure to 375 mg/kg bw/day ([Ema et al. 2003](#)).

di-n-heptyl phthalate

DHPP has been shown to cause decreased male AGD in rats following intrauterine exposure to 1000 mg/kg bw/day ([Saillenfait et al. 2011](#)).

Domain of Applicability

Taxonomic Applicability

Term	Scientific Term	Evidence	Links
human	Homo sapiens	Moderate	NCBI
rat	Rattus norvegicus	High	NCBI
mouse	Mus musculus	High	NCBI

Life Stage Applicability

Life Stage Evidence

Foetal	High
--------	------

Sex Applicability

Sex Evidence

Male	High
------	------

A short AGD in male offspring is a marker of insufficient androgen action during critical fetal developmental stages ([Schwartz et al. 2019](#); [Welsh et al. 2008](#)). A short AGD is thus a sign of undervirilization, which is also associated with a series of male reproductive disorders, including genital malformations and infertility in humans ([Juul et al. 2014](#); [Skakkebaek et al. 2001](#)).

There are numerous human epidemiological studies showing associations with intrauterine exposure to anti-androgenic chemicals and short AGD in newborn boys alongside other reproductive disorders ([Schwartz et al. 2019](#)). This underscores the human relevance of this AO. However, in reproductive toxicity studies and chemical risk assessment, rodents (rats and mice) are what is tested on. The list of chemicals inducing short male AGD in male rat offspring is extensive, as evidenced by the 'stressor' list and reviewed by ([Schwartz et al. 2019](#)).

Key Event Description

The anogenital distance (AGD) refers to the distance between anus and the external genitalia. In rodents and humans, the male AGD is approximately twice the length as the female AGD ([Salazar-Martinez et al. 2004](#); [Schwartz et al. 2019](#)). This sexual dimorphism is a consequence of sex hormone-dependent development of secondary sexual characteristics ([Schwartz et al. 2019](#)). In males, it is believed that androgens (primarily DHT) activate AR-positive cells in non-myotoc cells in the fetal perineum region to initiate differentiation of the perineal *levator ani* and *bulbocavernosus* (LABC) muscle complex ([Ipulan et al. 2014](#)). This AR-dependent process occurs within a critical window of development, around gestational days 15-18 in rats ([MacLeod et al. 2010](#)). In females, the absence of DHT prevents this masculinization effect from occurring.

The involvement of androgens in masculinization of the male fetus, including the perineum, has been known for a very long time

([Jost, 1953](#)), and AGD has historically been used to, for instance, sex newborn kittens. It is now well established that the AGD in newborns is a proxy readout for the intrauterine sex hormone milieu the fetus was developing. Too low androgen levels in XY fetuses makes the male AGD shorter, whereas excess (ectopic) androgen levels in XX fetuses makes the female AGD longer, in humans and rodents ([Schwartz et al, 2019](#)).

How it is Measured or Detected

The AGD is a morphometric measurement carried out by trained technicians (rodents) or medical staff (humans).

In rodent studies AGD is assessed as the distance between the genital papilla and the anus, and measured using a stereomicroscope with a micrometer eyepiece. The AGD index (AGDi) is often calculated by dividing AGD by the cube root of the body weight. It is important in statistical analysis to use litter as the statistical unit. This is done when more than one pup from each litter is examined. Statistical analyses is adjusted using litter as an independent, random and nested factor. AGD are analysed using body weight as covariate as recommended in Guidance Document 151 ([OECD, 2013](#)).

Regulatory Significance of the AO

In regulatory toxicology, the AGD is mandatory inclusions in OECD test guidelines used to test for developmental and reproductive toxicity of chemicals. Guidelines include 'TG 443 extended one-generation study', 'TG 421/422 reproductive toxicity screening studies' and 'TG 414 developmental toxicity study'.

References

- Aydoğar Ahabab M, Barlas N (2015) Influence of in utero di-n-hexyl phthalate and dicyclohexyl phthalate on fetal testicular development in rats. *Toxicol Lett* **233**: 125-137
- Boberg J, Axelstad M, Svingen T, Mandrup K, Christiansen S, Vinggaard AM, Hass U (2016) Multiple endocrine disrupting effects in rats perinatally exposed to butylparaben. *Toxicol Sci* **152**: 244-256
- Boberg J, Metzдорff S, Wortziger R, Axelstad M, Brokken L, Vinggaard AM, Dalgaard M, Nellemann C (2008) Impact of diisobutyl phthalate and other PPAR agonists on steroidogenesis and plasma insulin and leptin levels in fetal rats. *Toxicology* **250**: 75-81
- Bowman CJ, Barlow NJ, Turner KJ, Wallace DG, Foster PM (2003) Effects of in utero exposure to finasteride on androgen-dependent reproductive development in the male rat. *Toxicol Sci* **74**: 393-406
- Christiansen S, Boberg J, Axelstad M, Dalgaard M, Vinggaard AM, Metzдорff SB, Hass U (2010) Low-dose perinatal exposure to di(2-ethylhexyl) phthalate induces anti-androgenic effects in male rats. *Reprod Toxicol* **30**: 313-321
- Christiansen S, Scholze M, Dalgaard M, Vinggaard AM, Axelstad M, Kortenkamp A, Hass U (2009) Synergistic disruption of external male sex organ development by a mixture of four antiandrogens. *Environ Health Perspect* **117**: 1839-1846
- Draskau MK, Boberg J, Taxvig C, Pedersen M, Frandsen HL, Christiansen S, Svingen T (2019) In vitro and in vivo endocrine disrupting effects of the azole fungicides triticonazole and flusilazole. *Environ Pollut* **255**: 113309
- Ema M, Miyawaki E (2002) Effects on development of the reproductive system in male offspring of rats given butyl benzyl phthalate during late pregnancy. *Reprod Toxicol* **16**: 71-76
- Ema M, Miyawaki E, Hirose A, Kamata E (2003) Decreased anogenital distance and increased incidence of undescended testes in fetuses of rats given monobenzyl phthalate, a major metabolite of butyl benzyl phthalate. *Reprod Toxicol* **17**: 407-412
- Foster PM, Harris MW (2005) Changes in androgen-mediated reproductive development in male rat offspring following exposure to a single oral dose of flutamide at different gestational ages. *Toxicol Sci* **85**: 1024-1032
- Gray LE, Jr., Ostby J, Furr J, Price M, Veeramachaneni DN, Parks L (2000) Perinatal exposure to the phthalates DEHP, BBP, and DINP, but not DEP, DMP, or DOTP, alters sexual differentiation of the male rat. *Toxicol Sci* **58**: 350-365
- Gray LEJ, Ostby JS, Kelce WR (1994) Developmental effects of an environmental antiandrogen: the fungicide vinclozolin alters sex differentiation of the male rat. *Toxicol Appl Pharmacol* **129**: 46-52
- Hass U, Boberg J, Christiansen S, Jacobsen PR, Vinggaard AM, Taxvig C, Poulsen ME, Herrmann SS, Jensen BH, Petersen A, Clemmensen LH, Axelstad M (2012) Adverse effects on sexual development in rat offspring after low dose exposure to a mixture of endocrine disrupting pesticides. *Reprod Toxicol* **34**: 261-274
- Hass U, Scholze M, Christiansen S, Dalgaard M, Vinggaard AM, Axelstad M, Metzдорff SB, Kortenkamp A (2007) Combined exposure to anti-androgens exacerbates disruption of sexual differentiation in the rat. *Environ Health Perspect* **115 Suppl. 1**: 122-128

- Hoshino N, Iwai M, Okazaki Y (2005) A two-generation reproductive toxicity study of dicyclohexyl phthalate in rats. *J Toxicol Sci* **30 Spec No**: 79-96
- Hotchkiss AK, Parks-Saldutti LG, Ostby JS, Lambright C, Furr J, Vandenberg JG, Gray LEJ (2004) A mixture of the "antiandrogens" linuron and butyl benzyl phthalate alters sexual differentiation of the male rat in a cumulative fashion. *Biol Reprod* **71**: 1852-1861
- Howdeshell KL, Furr J, Lambright CR, Rider CV, Wilson VS, Gray LE, Jr. (2007) Cumulative effects of dibutyl phthalate and diethylhexyl phthalate on male rat reproductive tract development: altered fetal steroid hormones and genes. *Toxicol Sci* **99**: 190-202
- Ipulan LA, Suzuki K, Sakamoto Y, Murashima A, Imai Y, Omori A, Nakagata N, Nishinakamura R, Valasek P, Yamada G (2014) Nonmyocytic androgen receptor regulates the sexually dimorphic development of the embryonic bulbocavernosus muscle. *Endocrinology* **155**: 2467-2479
- Jarfelt K, Dalgaard M, Hass U, Borch J, Jacobsen H, Ladefoged O (2005) Antiandrogenic effects in male rats perinatally exposed to a mixture of di(2-ethylhexyl) phthalate and di(2-ethylhexyl) adipate. *Reprod Toxicol* **19**: 505-515
- Jost A (1953) Problems of fetal endocrinology: The gonadal and hypophyseal hormones. *Recent Prog Horm Res* **8**: 379-418
- Juul A, Almstrup K, Andersson AM, Jensen TK, Jorgensen N, Main KM, Rajpert-De Meyts E, Toppari J, Skakkebaek NE (2014) Possible fetal determinants of male infertility. *Nat Rev Endocrinol* **10**: 553-562
- Kita DH, Meyer KB, Venturelli AC, Adams R, Machado DL, Morais RN, Swan SH, Gennings C, Martino-Andrade AJ (2016) Manipulation of pre and postnatal androgen environments and anogenital distance in rats. *Toxicology* **368-369**: 152-161
- Laier P, Metzdorff SB, Borch J, Hagen ML, Hass U, Christiansen S, Axelstad M, Kledal T, Dalgaard M, McKinnell C, Brokken LJ, Vinggaard AM (2006) Mechanisms of action underlying the antiandrogenic effects of the fungicide prochloraz. *Toxicol Appl Pharmacol* **213**: 2
- Li M, Qiu L, Zhang Y, Hua Y, Tu S, He Y, Wen S, Wang Q, Wei G (2013) Dose-related effect by maternal exposure to di-(2-ethylhexyl) phthalate plasticizer on inducing hypospadiac male rats. *Environ Toxicol Pharmacol* **35**: 55-60
- Lin H, Lian QQ, Hu GX, Jin Y, Zhang Y, Hardy DO, Chen GR, Lu ZQ, Sottas CM, Hardy MP, Ge RS (2009) In utero and lactational exposures to diethylhexyl-phthalate affect two populations of Leydig cells in male Long-Evans rats. *Biol Reprod* **80**: 882-888
- Loeffler IK, Peterson RE (1999) Interactive effects of TCDD and p,p'-DDE on male reproductive tract development in in utero and lactationally exposed rats. *Toxicol Appl Pharmacol* **154**: 28-39
- MacLeod DJ, Sharpe RM, Welsh M, Fisk M, Scott HM, Hutchison GR, Drake AJ, van den Driesche S (2010) Androgen action in the masculinization programming window and development of male reproductive organs. *Int J Androl* **33**: 279-287
- Matsuura I, Saitoh T, Ashina M, Wako Y, Iwata H, Toyota N, Ishizuka Y, Namiki M, Hoshino N, Tsuchitani M (2005) Evaluation of a two-generation reproduction toxicity study adding endpoints to detect endocrine disrupting activity using vinclozolin. *J Toxicol Sci* **30 Spec No**: 163-168
- McIntyre BS, Barlow NJ, Foster PM (2001) Androgen-mediated development in male rat offspring exposed to flutamide in utero: permanence and correlation of early postnatal changes in anogenital distance and nipple retention with malformations in androgen-dependent tissues. *Toxicol Sci* **62**: 236-249
- McIntyre BS, Barlow NJ, Sar M, Wallace DG, Foster PM (2002) Effects of in utero linuron exposure on rat Wolffian duct development. *Reprod Toxicol* **16**: 131-139
- Melching-Kollmuss S, Fussell KC, Schneider S, Buesen R, Groeters S, Strauss V, van Ravenzwaay B (2017) Comparing effect levels of regulatory studies with endpoints derived in targeted anti-androgenic studies: example prochloraz. *Arch Toxicol* **91**: 143-162
- Moore RW, Rudy TA, Lin TM, Ko K, Peterson RE (2001) Abnormalities of sexual development in male rats with in utero and lactational exposure to the antiandrogenic plasticizer Di(2-ethylhexyl) phthalate. *Environ Health Perspect* **109**: 229-237
- Mylchreest E, Sar M, Cattley RC, Foster PM (1999) Disruption of androgen-regulated male reproductive development by di(n-butyl) phthalate during late gestation in rats is different from flutamide. *Toxicol Appl Pharmacol* **156**: 81-95
- Nagao T, Ohta R, Marumo H, Shindo T, Yoshimura S, Ono H (2000) Effect of butyl benzyl phthalate in Sprague-Dawley rats after gavage administration: a two-generation reproductive study. *Reprod Toxicol* **14**: 513-532
- Nardelli TC, Albert O, Lalancette C, Culty M, Hales BF, Robaire B (2017) In utero and lactational exposure study in rats to identify replacements for di(2-ethylhexyl) phthalate. *Sci Rep* **7**: 3862
- Noriega NC, Ostby J, Lambright C, Wilson VS, Gray LE, Jr. (2005) Late gestational exposure to the fungicide prochloraz delays the onset of parturition and causes reproductive malformations in male but not female rat offspring. *Biol Reprod* **72**: 1324-1335

OECD. (2013) Guidance document in support of the test guideline on the extended one generation reproductive toxicity study No. 151.

Ostby J, Kelce WR, Lambright C, Wolf CJ, Mann P, Gray CLJ (1999) The fungicide procymidone alters sexual differentiation in the male rat by acting as an androgen-receptor antagonist in vivo and in vitro. *Toxicol Ind Health* **15**: 80-93

Saillenfait AM, Gallissot F, Sabaté JP (2009a) Differential developmental toxicities of di-n-hexyl phthalate and dicyclohexyl phthalate administered orally to rats. *J Appl Toxicol* **29**: 510-521

Saillenfait AM, Roudot AC, Gallissot F, Sabaté JP (2011) Prenatal developmental toxicity studies on di-n-heptyl and di-n-octyl phthalates in Sprague-Dawley rats. *Reprod Toxicol* **32**: 268-276

Saillenfait AM, Sabaté JP, Gallissot F (2009b) Effects of in utero exposure to di-n-hexyl phthalate on the reproductive development of the male rat. *Reprod Toxicol* **28**: 468-476

Salazar-Martinez E, Romano-Riquer P, Yanez-Marquez E, Longnecker MP, Hernandez-Avila M (2004) Anogenital distance in human male and female newborns: a descriptive, cross-sectional study. *Environ Health* **3**: 8

Schneider S, Kaufmann W, Strauss V, van Ravenzwaay B (2011) Vinclozolin: a feasibility and sensitivity study of the ILSI-HESI F1-extended one-generation rat reproduction protocol. *Regulatory Toxicology and Pharmacology* **59**: 91-100

Schwartz CL, Christiansen S, Vinggaard AM, Axelstad M, Hass U, Svingen T (2019) Anogenital distance as a toxicological or clinical marker for fetal androgen action and risk for reproductive disorders. *Arch Toxicol* **93**: 253-272

Scott HM, Hutchison GR, Mahood IK, Hallmark N, Welsh M, De Gendt K, Verhoeven H, O'Shaughnessy P, Sharpe RM (2007) Role of androgens in fetal testis development and dysgenesis. *Endocrinology* **148**: 2027-2036

Skakkebaek NE, Rajpert-De Meyts E, Main KM (2001) Testicular dysgenesis syndrome: an increasingly common developmental disorder with environmental aspects. *Hum Reprod* **16**: 972-978

Taxvig C, Vinggaard AM, Hass U, Axelstad M, Metzдорff S, Nellemann C (2008) Endocrine-disrupting properties in vivo of widely used azole fungicides. *Int J Androl* **31**: 170-177

Turner KJ, Barlow NJ, Struve MF, Wallace DG, Gaido KW, Dorman DC, Foster PM (2002) Effects of in utero exposure to the organophosphate insecticide fenitrothion on androgen-dependent reproductive development in the Crl:CD(SD)BR rat. *Toxicol Sci* **68**: 174-183

Tyl RW, Myers CB, Marr MC, Fail PA, Seely JC, Brine DR, Barter RA, Butala JH (2004) Reproductive toxicity evaluation of dietary butyl benzyl phthalate (BBP) in rats. *Reprod Toxicol* **18**: 241-264

Van den Driesche S, Kolovos P, Platts S, Drake AJ, Sharpe RM (2012) Inter-relationship between testicular dysgenesis and Leydig cell function in the masculinization programming window in the rat. *PLoS one* **7**: e30111

Welsh M, Saunders PT, Fiskens M, Scott HM, Hutchison GR, Smith LB, Sharpe RM (2008) Identification in rats of a programming window for reproductive tract masculinization, disruption of which leads to hypospadias and cryptorchidism. *J Clin Invest* **118**: 1479-1490

Welsh M, Saunders PT, Sharpe RM (2007) The critical time window for androgen-dependent development of the Wolffian duct in the rat. *Endocrinology* **148**: 3185-3195

Wolf CJ, LeBlanc GA, Gray LE, Jr. (2004) Interactive effects of vinclozolin and testosterone propionate on pregnancy and sexual differentiation of the male and female SD rat. *Toxicol Sci* **78**: 135-143

Wolf CJJ, Lambright C, Mann P, Price M, Cooper RL, Ostby J, Gray CLJ (1999) Administration of potentially antiandrogenic pesticides (procymidone, linuron, iprodione, chlozolinate, p,p'-DDE, and ketoconazole) and toxic substances (dibutyl- and diethylhexyl phthalate, PCB 169, and ethane dimethane sulphonate) during sexual differentiation produces diverse profiles of reproductive malformations in the male rat. *Toxicol Ind Health* **15**: 94-118

Zhang L, Dong L, Ding S, Qiao P, Wang C, Zhang M, Zhang L, Du Q, Li Y, Tang N, Chang B (2014) Effects of n-butylparaben on steroidogenesis and spermatogenesis through changed E₂ levels in male rat offspring. *Environ Toxicol Pharmacol* **37**: 705-717

Appendix 2

List of Key Event Relationships in the AOP

List of Adjacent Key Event Relationships

Relationship: 2130: Antagonism, Androgen receptor leads to Decrease, AR activation

AOPs Referencing Relationship

AOP Name	Adjacency	Weight of Evidence	Quantitative Understanding
Androgen receptor (AR) antagonism leading to short anogenital distance (AGD) in male (mammalian) offspring	adjacent	High	High
Androgen receptor (AR) antagonism leading to nipple retention (NR) in male (mammalian) offspring	adjacent	High	

[Relationship: 2128: Decrease, AR activation leads to decrease, transcription of genes by AR](#)

AOPs Referencing Relationship

AOP Name	Adjacency	Weight of Evidence	Quantitative Understanding
Androgen receptor (AR) antagonism leading to short anogenital distance (AGD) in male (mammalian) offspring	adjacent	High	Moderate

Evidence Supporting Applicability of this Relationship

Taxonomic Applicability

Term	Scientific Term	Evidence	Links
human	Homo sapiens	High	NCBI
mouse	Mus musculus	High	NCBI
human and other cells in culture	human and other cells in culture	High	NCBI
rat	Rattus norvegicus	High	NCBI

Life Stage Applicability

Life Stage	Evidence
During development and at adulthood	High

Sex Applicability

Sex	Evidence
Male	High
Female	High

[Relationship: 2129: decrease, transcription of genes by AR leads to short male AGD](#)

AOPs Referencing Relationship

AOP Name	Adjacency	Weight of Evidence	Quantitative Understanding
Androgen receptor (AR) antagonism leading to short anogenital distance (AGD) in male (mammalian) offspring	adjacent	Moderate	Low

List of Non Adjacent Key Event Relationships

[Relationship: 2123: Antagonism, Androgen receptor leads to short male AGD](#)

AOPs Referencing Relationship

AOP Name	Adjacency	Weight of Evidence	Quantitative Understanding
----------	-----------	--------------------	----------------------------

Androgen receptor (AR) antagonism leading to short anogenital distance (AGD) in male (mammalian) offspring	AOP Name	non-Adverse	Moderate Weight of Evidence	Low	Quantitative Understanding
--	-----------------	-------------	-----------------------------	-----	-----------------------------------